arXiv:0801.0820v2 [math.AG] 25 Feb 2008

MODULI OF ADHM SHEAVES AND LOCAL DONALDSON-THOMAS THEORY

D.-E. DIACONESCU

Dedicated to The Memory of John Ionesei – John Prinţul

Abstract. The ADHM construction establishes a one-to-one correspondence between framed torsion free sheaves on the projective plane and stable framed representations of a quiver with relations in the category of complex vector spaces. We construct an algebraic moduli space of stable framed representations of the same quiver with relations in the abelian category of coherent sheaves on a smooth complex projective variety X. If X is a curve, we prove that this moduli space is virtually smooth and related by relative Beilinson spectral sequence to the curve counting construction via stable pairs of Pandharipande and Thomas. This yields a new conjectural construction for the local Donaldson-Thomas theory of curves as well as a natural higher rank generalization.

Contents

1. Introduction 2
2. ADHM Sheaves 7
2.1. Stable ADHM Sheaves 7
2.2. Properties of ADHM Sheaves 8
2.3. Boundedness 10
3. Moduli Space of Stable ADHM Sheaves 12
3.1. The Parameter Space 13
3.2. The Moduli Space 21
3.3. Torus Actions and Fixed Loci 23
4. Deformation Theory of ADHM Sheaves on Curves 28
4.1. Deformation Complex 28
4.2. ADHM Sheaves with Trivial Framing 40
5. Virtual Smoothness for ADHM Sheaves on Curves 43
5.1. The Tangent-Obstruction Complex 43
5.2. Virtual Smoothness 48
6. Admissible Pairs 51
7. ADHM Sheaves via Relative Beilinson Spectral Sequence 56
7.1. Relative Beilinson Monad for Admissible Pairs 56
7.2. Admissible Pairs and ADHM Sheaves 64
7.3. Isomorphism of Moduli Spaces 69

7.4. Comparison of Tangent-Obstruction Theories 70
Appendix A. Base Change Properties of Direct Images 79
References 80

1. Introduction

An ADHM quiver Q_{ADHM} is a quiver of the form

(1.1)



with one relation specified by the linear combination of paths

$$a_1a_2 - a_2a_1 + cb. \tag{1.2}$$

ADHM sheaves are framed twisted representations of an ADHM quiver in the abelian category of coherent $\mathcal{O}_X$-modules of a smooth projective variety X over $\mathbb{C}$. The twisting data consists of two invertible $\mathcal{O}_X$-modules M_1, M_2, while the framing data consists of a locally free $\mathcal{O}_X$-module E_∞ (see (2.1) for a precise definition.) In order to keep the notation short, we will denote by $\mathcal{X}$ the data (X, M_1, M_2, E_∞).

Motivated by BPS state counting problems in string theory, in this paper we study the geometry of the moduli space of ADHM sheaves on smooth projective varieties, and their relation to relative Beilinson monads for projective plane bundles over curves. Note that quiver sheaves have been previously considered in the literature in [4, 3, 30, 67, 11]. In particular ADHM sheaves have been previously studied in [67] in relation to the relative Beilinson monad for noncommutative surfaces over curves. A similar relation between quiver sheaves and relative Beilinson monad has been employed in the context of integrable systems in [11]. Moduli problems for such objects have been also previously considered in [65, 29, 66, 5], as well as [11]. We will later elaborate on the relation between some of the above papers and the present work. Let us first present out main results.

Let X be a smooth projective variety over $\mathbb{C}$ equipped with a very ample line bundle $\mathcal{O}_X(1)$. We will consider ADHM sheaves subject to a stability condition (2.2). A routine argument shows that flat families of stable ADHM sheaves with fixed Hilbert polynomial P on X form a fibered groupoid $\mathfrak{p} : \mathfrak{M}_{ADHM}(\mathcal{X}, P) \to \mathfrak{S}$ over the category $\mathfrak{S}$ of schemes of finite type over $\mathbb{C}$ (proposition 3.2). In section (3.2) we prove our first result,

Theorem 1.1. *The groupoid $\mathfrak{M}_{ADHM}(\mathcal{X}, P)$ is a separated algebraic space of finite type over $\mathbb{C}$.*

Remark 1.2. *Since $\mathfrak{M}_{ADHM}(\mathcal{X}, P)$ is an algebraic space, it is isomorphic to its coarse moduli space constructed in* [42]. *Throughout this paper we will identify $\mathfrak{M}_{ADHM}(\mathcal{X}, P)$ with its coarse moduli space. In particular for* *us a K-valued point of $\mathfrak{M}_{ADHM}(\mathcal{X}, P)$ for some field K over $\mathbb{C}$ will be an isomorphism class of stable ADHM sheaves on $X_K = X \times_{Spec(\mathbb{C})} \mathrm{Spec}(K)$.*

Remark 1.3. (i) *Note that the stability condition* (ii) *in (2.2) is very similar to the nondegeneracy condition formulated in* [67, Thm 3.1]. *However the later disallows all proper subsheaves $E' \subset E$ satisfying the conditions listed there, not only the saturated ones. It will become clear in section (7) that this difference has important consequences for applications to local Donaldson-Thomas theory.*

(ii) *The stability condition (2.2) is in fact the asymptotic form of a GIT stability condition for ADHM sheaves analogous the stability conditions introduced in* [65, 29, 66]. *Moreover, using standard GIT techniques* [56, 39, 38, 65, 29, 66] *one can prove* [22] *that the coarse moduli space of ADHM sheaves is in fact a quasi-projective scheme over $\mathbb{C}$. This is a natural generalization of previous work on moduli of decorated sheaves which includes* [68, 15, 17, 13, 14, 16]. *However the proof is very technical, and the construction of the moduli space as an algebraic space suffices for our purposes.*

Remark 1.4. *Note that the proof of theorem (1.1) also provides a presentation of the moduli space $\mathfrak{M}_{ADHM}(\mathcal{X}, P)$ as a quotient stack R/G where R is a quasi-projective scheme over $\mathbb{C}$ and $G = GL(N, \mathbb{C})$ for some $N \in \mathbb{Z}_{>0}$. Moreover, there exists a natural G-equivariant universal family of stable ADHM sheaves $\mathcal{E}_R$ on $R \times X$.*

Next, suppose X is a smooth projective curve of genus g over $\mathbb{C}$. Our next result establishes equivariant virtual smoothness of the moduli space of ADHM sheaves with respect to a natural class of algebraic torus actions. In this case the Hilbert polynomial of an ADHM sheaf $\mathcal{E}$ is determined by the rank $r \in \mathbb{Z}$, $r \geq 1$ and the degree $e \in \mathbb{Z}$. We will denote by $\mathfrak{M}_{ADHM}(\mathcal{X}, r, e)$ the moduli space of stable ADHM sheaves on X with fixed (r, e).

Theorem 1.5. *Suppose X is a smooth projective curve over $\mathbb{C}$ and suppose there exists a torus action $\mathbf{T} \times \mathfrak{M}_{ADHM}(\mathcal{X}, r, e) \to \mathfrak{M}_{ADHM}(\mathcal{X}, r, e)$ induced by a $\mathbf{T}$-action on the smooth quasi-projective atlas R so that the universal family $\mathcal{E}_R$ is $\mathbf{T}$-equivariant. Then the moduli space $\mathfrak{M}_{ADHM}(\mathcal{X}, r, e)$ has a $\mathbf{T}$-equivariant perfect tangent-obstruction theory.*

Here we employ definition [47, Def. 2.1] for a perfect tangent-obstruction theory of a Deligne-Mumford stack. Theorem (1.5) is proven in section (5.2).

Note that there are natural algebraic torus actions on $\mathfrak{M}_{ADHM}(\mathcal{X}, P)$ satisfying the conditions of theorem (1.5) presented in examples (3.12), (3.13). Proposition (3.15) shows that if X is a smooth projective curve, the fixed locus $\mathfrak{M}(\mathcal{X}, r, e)^{\mathbf{T}}$ of the torus action defined in example (3.13) is a projective scheme over $\mathbb{C}$. Then theorem (1.5), proposition (3.15) and [31] yield the following corollary, which is also proved in section (5.2).

Corollary 1.6. *Let X be a smooth projective curve over $\mathbb{C}$ and $E_\infty = \oplus_{a=1}^{r_\infty} L_a$ where $r_\infty \in \mathbb{Z}$, $r_\infty \geq 1$ and L_a, $a = 1, \ldots, r_\infty$, are line bundles* *on X. Let $\mathbf{T} \times \mathfrak{M}_{ADHM}(\mathcal{X}, r, e) \to \mathfrak{M}_{ADHM}(\mathcal{X}, r, e)$ be the torus action defined in example (3.13). Then the fixed locus $\mathfrak{M}_{ADHM}(\mathcal{X}, r, e)^{\mathbf{T}}]$ is a projective scheme over $\mathbb{C}$ equipped with a virtual cycle $[\mathfrak{M}_{ADHM}(\mathcal{X}, r, e)^{\mathbf{T}}]$ and a $\mathbf{T}$-equivariant virtual normal bundle $N^{vir}_{\mathfrak{M}_{ADHM}(\mathcal{X},r,e)^{\mathbf{T}}/\mathfrak{M}_{ADHM}(\mathcal{X},r,e)}$ defined as an element of the K-theory of coherent locally free sheaves on $\mathfrak{M}_{ADHM}(\mathcal{X}, r, e)^{\mathbf{T}}$.*

Corollary (1.6) allows us to define an equivariant ADHM theory of curves by analogy with [18, 58].

Definition 1.7. *Suppose X is a smooth projective curve over $\mathbb{C}$ and $E_\infty = \oplus_{a=1}^{r_\infty} L_a$ where $r_\infty \in \mathbb{Z}$, $r_\infty \geq 1$ and L_a, $a = 1, \ldots, r_\infty$, are line bundles on X. Let $\mathbf{T} \times \mathfrak{M}_{ADHM}(\mathcal{X}, r, e) \to \mathfrak{M}_{ADHM}(\mathcal{X}, r, e)$ be the torus action defined in example (3.13). Then the rank r equivariant ADHM theory of the data $\mathcal{X} = (X, M_1, M_2, E_\infty)$ is defined by*

$$Z_{ADHM}(\mathcal{X})_r(q) = \sum_{e\in\mathbb{Z}} q^e \int_{[\mathfrak{M}_{ADHM}(\mathcal{X},r,e)^{\mathbf{T}}]} \frac{1}{e_{\mathbf{T}}(N^{vir}_{\mathfrak{M}_{ADHM}(\mathcal{X},r,e)^{\mathbf{T}}/\mathfrak{M}_{ADHM}(\mathcal{X},r,e)})} \tag{1.3}$$

where $e_{\mathbf{T}}(N^{vir}_{\mathfrak{M}_{ADHM}(\mathcal{X},r,e)^{\mathbf{T}}/\mathfrak{M}_{ADHM}(\mathcal{X},r,e)})$ is the $\mathbf{T}$-equivariant Euler class of $N^{vir}_{\mathfrak{M}_{ADHM}(\mathcal{X},r,e)^{\mathbf{T}}/\mathfrak{M}_{ADHM}(\mathcal{X},r,e)}$.

Remark 1.8. *Note that the partition function* (1.3) *is a natural generalization of the instanton counting function defined in* [54] *to the relative setting. One can also generalize the K-theoretic partition function of* [55] *to the relative setting relying on corollary (1.6) and the results* [19, 25]. *The K-theoretic ADHM theory of curves will be treated elsewhere.*

String theoretic arguments predict that the ADHM theory of curves should be related to the local Donaldson-Thomas theory of curves defined in [50, 51, 18, 58]. In sections (6), (7) we show that the relative Beilinson spectral sequence [59] relates stable ADHM sheaves with $E_\infty = \mathcal{O}_X$ on a smooth projective curve X to stable pairs [45, 46, 61] on the projective plane bundle $Y = \mathrm{Proj}(\mathcal{O}_X \oplus M_1 \oplus M_2)$ over X. The later have been recently employed by Pandharipande and Thomas [61] as a new curve counting device on smooth projective threefolds (see [60, 62] for further results.) As mentioned above, similar relations between quiver sheaves and the relative Beilinson monad have been previously found in [67, 11]. However, the connection to the local Donaldson-Thomas theory of curves via stable pairs is new, and relies heavily on the stability condition (2.2).

In order to establish a connection between stable ADHM sheaves and stable pairs, we introduce in section (6) (see definition (6.2)) a new class of objects – called admissible pairs – which provide a natural intermediate construction between stable pairs and stable ADHM sheaves. Admissible pairs are proven to be equivalent to stable pairs in proposition (6.4) and they are also naturally related to stable ADHM sheaves with $E_\infty = \mathcal{O}_X$

by the relative Beilinson spectral sequence (proposition 7.7). Admissible pairs are characterized by a pair (d,n) of numerical invariants defined in (6.2). The main properties of admissible pairs derived from [45, 46, 61] are summarized by the following theorem.

Theorem 1.9. (i) *There exists a quasi-projective moduli space* $\mathfrak{M}_{Adm}(Y,d,n)$ *over* $\mathbb{C}$ *of admissible pairs on* Y *with fixed numerical invariants* (d,n)*.*

(ii) *There is a* $\mathbf{T} = \mathbb{C}^\times \times \mathbb{C}^\times$ *action on* $\mathfrak{M}_{Adm}(Y,d,n)$ *induced by a scaling action on* Y*, so that* $\mathfrak{M}_{Adm}(Y,d,n)$ *is equipped with a natural* $\mathbf{T}$*-equivariant perfect tangent-obstruction theory.*

(iii) *The fixed locus* $\mathfrak{M}_{Adm}(Y,d,n)^{\mathbf{T}}$ *is a projective scheme over* $\mathbb{C}$ *equipped with a virtual cycle* $[\mathfrak{M}_{Adm}(Y,d,n)^{\mathbf{T}}]$ *and an equivariant virtual normal bundle* $N^{vir}_{\mathfrak{M}_{Adm}(Y,d,n)^{\mathbf{T}}/\mathfrak{M}_{Adm}(Y,d,n)}$*.*

A proof of this theorem based on [45, 46, 61] is outlined in section (6).

Theorem (1.9) allows us to define the local equivariant Pandharipande-Thomas [61] theory of curves.

Definition 1.10. *Let* X *be a smooth projective curve of genus* g*,* M_1, M_2 *be invertible sheaves on* X *and* $Y = \mathrm{Proj}(\mathcal{O}_X \oplus M_1 \oplus M_2)$*. Then we define the degree* d *local Pandharipande-Thomas theory of the triple* (X, M_1, M_2) *to be*

$$Z_{PT}(X,M_1,M_2)_d(q) = \sum_{n\in\mathbb{Z}} q^n \int_{[\mathfrak{M}_{Adm}(Y,d,n)^{\mathbf{T}}]} \frac{1}{e_{\mathbf{T}}(N^{vir}_{\mathfrak{M}_{Adm}(Y,d,n)^{\mathbf{T}}/\mathfrak{M}_{Adm}(Y,d,n)})}.$$

Our third result is the following.

Theorem 1.11. *Let* X *be a smooth projective curve of genus* g*,* M_1, M_2 *be invertible sheaves on* X *and* $E_\infty = \mathcal{O}_X$*. Let* $Y = \mathrm{Proj}(\mathcal{O}_X \oplus M_1 \oplus M_2)$ *and let* $\mathbf{T} = \mathbb{C}^\times \times \mathbb{C}^\times$ *act on* $\mathfrak{M}_{ADHM}(\mathcal{X}, r, e)$ *as in example (3.12). Then there is a* $\mathbf{T}$*-equivariant isomorphism of algebraic moduli spaces*

$$\mathfrak{f} : \mathfrak{M}_{Adm}(Y,d,n) \simeq \mathfrak{M}_{ADHM}(\mathcal{X}, d, n + d(g-1))$$

so that the perfect tangent-obstruction theories of the two moduli spaces are compatible with respect to $\mathfrak{f}$*.*

Here compatible perfect obstruction theories is meant in the sense of [47, Def. 4.1]. Theorem (1.11) is proven in sections (7.3), (7.4).

Theorem (1.11) implies

Corollary 1.12. *Let* X *be a smooth projective curve genus* g*,* M_1, M_2 *be invertible sheaves on* X *and* $Y = \mathrm{Proj}(\mathcal{O}_X \oplus M_1 \oplus M_2)$*. Then we have*

$$Z_{PT}(X,M_1,M_2)_d(q) = q^{-d(g-1)} Z_{ADHM}(\mathcal{X})_d(q) \tag{1.4}$$

for any $d \geq 1$*, where* $\mathcal{X} = (X, M_1, M_2, \mathcal{O}_X)$*.*

According to [61, Conj. 3.3] we have the following conjectural relation between the Pandharipande-Thomas and the local Donaldson-Thomas theory of curves [50, 51, 18, 58]

$$Z_{PT}(X,M_1,M_2)_d(q) = Z'_{DT}(X,M_1,M_2)_d(q).$$

Therefore corollary (1.12) immediately yields a similar conjectural relation between the ADHM theory of curves with trivial framing and the local Donaldson-Thomas theory of curves.

For more general framing $E_\infty = \oplus_{a=1}^{r_\infty} L_a$ as in definition (1.7), we expect the ADHM theory of $Z_{ADHM}(\mathcal{X})_r(q)$ to be related to the higher rank local Donaldson-Thomas theory of curves. This problem, as well as some geometric applications will be treated elsewhere.

Note added for Version 2. Moduli spaces of twisted ADHM sheaves have also been recently studied by Marcos Jardim in [41], subject to a more general stability condition. In particular [41, Thm 24] establishes that the moduli space of stable twisted ADHM sheaves on a complex projective space is a quasi-projective variety.

Notation and Conventions. Throughout this paper, we will denote by $\mathfrak{S}$ the category of schemes of finite type over $\mathbb{C}$. For any such schemes X, S we set $X_S = S \times X$ and $X_s = \mathrm{Spec}(k(s)) \times_S X$ for any point $s \in S$, where $k(s)$ is the residual field of s. Let also $p_S : X_S \to S$, $p_X : X_S \to X$ denote the canonical projections. We will also set $F_S = p_X^* F$ for any $\mathcal{O}_X$-module F. Given a morphism $f : S' \to S$, we will denote by $f_X = f \times 1_X : X_{S'} \to X_S$. Any morphism $f : S' \to S$, yields a commutative diagram of the form

$$\begin{tikzcd}
X_{S'} \arrow[r, "p'_X"] \arrow[d, "f_X"'] & X \arrow[d, "1_X"] \\
X_S \arrow[r, "p_X"] & X.
\end{tikzcd}$$

Then for any $\mathcal{O}_X$-module F there is a canonical isomorphism $F_{S'} \simeq f_X^* F_S$ which will be implicit in the following. Finally, we willy employ the conventions of [20, Ch. 1.2, Ch. 1.3] for computations in derived categories.

Acknowledgments. The physical idea of this construction originated in discussions with Robbert Dijkgraaf in spring 2006 concerning the physics of Donaldson-Thomas theory. It was reinforced in independent discussions with Nick Halmagy in summer 2006 concerning different aspects of D-brane BPS states. I am very grateful to them for sharing their insights with me.

During the completion of this project I greatly benefited from discussions and correspondence with Richard Thomas who kindly made a preliminary version of [61] available to me prior to publication. The results of sections (6), (7) were prompted by his comments on an earlier version of the present paper. The proof of lemma (5.3) is also due to him, eventual errors being my own.

I am very grateful to Tony Pantev for patient explanations about moduli stacks and virtual cycles, as well as invaluable help in proving theorem (1.1). The proof of lemma (3.11) is also due to him, eventual errors being, again, my own.

I am also very grateful to Ionut Ciocan-Fontanine and Bumsig Kim for very helpful discussions and collaboration on a related project. In particular, I owe special thanks to Ionut Ciocan-Fontanine for pointing out references [42, 19, 25] to me.

I would also like to thank Aaron Bertram, Giulio Bonelli, Ugo Bruzzo, Ron Donagi, Bogdan Florea, Alberto Garcia-Raboso, Sheldon Katz, Alexander Kuznetzov, Y.P. Lee, Sergio Lukic, Greg Moore, Alessandro Tanzini, and especially Cumrun Vafa for very helpful conversations and encouragement.

Part of this work was done during a visit at SISSA Trieste in spring 2007, as well as during the "Derived Categories in Mathematics and Physics - Snowbird, June 17-23, 2007" workshop, the "Workshop on Algebraic Geometry and Physics - Seoul, June 25-29, 2007", the "String Theory and Quantum Geometry - Aspen 2007" workshop and the "Simons Workshop on Mathematics and Physics, Stony Brook 2007" which provided a very stimulating mathematical atmosphere.

I would also like to acknowledge the partial support of NSF grant PHY-0555374-2006.

2. ADHM Sheaves

2.1. **Stable ADHM Sheaves.**

Quiver sheaves are representations of a quiver with relations in the abelian category of modules over a scheme or more generally a ringed space [4, 3, 30]. Let X be an arbitrary scheme, M_1, M_2 be fixed invertible sheaves on X and E_∞ be a coherent $\mathcal{O}_X$-module. In this paper we will consider framed twisted representations of an ADHM quiver in the abelian category of coherent $\mathcal{O}_X$-modules. As stated in the introduction, we will denote the data (X, M_1, M_2, E_∞) by $\mathcal{X}$.

Definition 2.1. (i) *An ADHM sheaf on X is defined by the data $\mathcal{E} = (E, \Phi_{1,2}, \phi, \psi)$ where E is a coherent $\mathcal{O}_X$-module and*

$$\Phi_i : E \otimes_X M_i \to E, \quad \phi : E \otimes M_1 \otimes_X M_2 \to E_\infty, \quad \psi : E_\infty \to E,$$

with $i = 1, 2$, are morphisms of $\mathcal{O}_X$-modules satisfying

$$\Phi_1 \circ (\Phi_2 \otimes 1_{M_1}) - \Phi_2 \circ (\Phi_1 \otimes 1_{M_2}) + \psi \circ \phi = 0.$$

We will refer to M_1, M_2 as twisting data and E_∞ as framing data.

(ii) *A morphism $\xi : \mathcal{E} \to \mathcal{E}'$ of ADHM sheaves on X is a morphism of $\mathcal{O}_X$-modules $\xi : E \to E'$ so that the following diagrams are commutative*

(2.1)

$$\begin{tikzcd}
E \otimes_X M_i \arrow[r, "\Phi_i"] \arrow[d, "\xi\otimes 1_{M_i}"'] & E \arrow[d, "\xi"] \\
E' \otimes M_i \arrow[r, "\Phi_i'"] & E'
\end{tikzcd}
\qquad
\begin{tikzcd}
E_\infty \arrow[r, "\psi"] \arrow[d, "1_{E_\infty}"'] & E \arrow[d, "\xi"] \\
E_\infty \arrow[r, "\psi'"] & E'
\end{tikzcd}$$

$$\begin{tikzcd}
E \otimes_Y M_1 \otimes_X M_2 \arrow[r, "\phi"] \arrow[d, "\xi\otimes 1_{M_1 \otimes_X M_2}"'] & E_\infty \arrow[d, "1_{E_\infty}"] \\
E' \otimes_X M_1 \otimes_X M_2 \arrow[r, "\phi'"] & E_\infty
\end{tikzcd}$$

(iii) A morphism $\xi : \mathcal{E} \to \mathcal{E}'$ is an isomorphism of ADHM sheaves if $\xi : E \to E'$ is an isomorphism of $\mathcal{O}_X$-modules.

Suppose X is a smooth projective variety over an infinite field k of characteristic 0. In addition to the twisting data M_1, M_2 and framing data E_∞, we will also fix a very ample line bundle $\mathcal{O}_X(1)$ on X. Let M_{12} denote $M_1 \otimes_X M_2$.

Definition 2.2. *An ADHM sheaf $\mathcal{E} = (E, \Phi_{1,2}, \phi, \psi)$ on X with twisting data M_1, M_2 and framing data E_∞ is stable if the following conditions are satisfied*

- *(i) E is a torsion-free $\mathcal{O}_X$-module*
- *(ii) ψ is not identically zero*
- *(iii) There exists no nontrivial proper saturated subsheaf $0 \subset E' \subset E$ so that $\Phi_i(E' \otimes_X M_i) \subseteq E'$ for $i = 1, 2$ and $\mathrm{Im}(\psi) \subseteq E'$.*

The stable ADHM sheaf $\mathcal{E}$ will be said to have Hilbert polynomial P if E has Hilbert polynomial P.

2.2. **Properties of ADHM Sheaves.** Next we establish some properties of ADHM sheaves will be needed later in the paper. We will work under the same assumptions as above i.e. X is a smooth projective variety over an infinite field k of characteristic 0 equipped with a very ample line bundle $\mathcal{O}_X(1)$. M_1, M_2 will be fixed twisting data and the locally free $\mathcal{O}_X$-module E_∞ will be fixed framing data.

Lemma 2.3. (*i*) *Let $\xi : \mathcal{E} \to \mathcal{E}'$ be a morphism of stable ADHM sheaves on X, where $\mathcal{E}, \mathcal{E}'$ have identical Hilbert polynomials. Then ξ is either trivial or an isomorphism.*

(*ii*) *If the base field k is algebraically closed, the automorphism group of a stable ADHM sheaf on X is trivial.*

Proof. Let $\xi : E \to E'$ be a morphism of coherent $\mathcal{O}_X$-modules which yields a morphism of ADHM sheaves i.e. the diagrams (2.1) for $\mathcal{E}, \mathcal{E}', \xi$ are commutative. Then it follows that

$$\Phi'_i(\mathrm{Im}(\xi) \otimes_X M_i) \subseteq \mathrm{Im}(\xi), \qquad \mathrm{Im}(\psi') \subseteq \mathrm{Im}(\xi). \tag{2.2}$$

Suppose ξ is not identically zero, and kernel $\mathrm{Ker}(\xi)$ is a nontrivial proper subsheaf of E. Since E is torsion free, $\mathrm{Ker}(\xi)$ must be torsion free as well, hence $\mathrm{rk}(\mathrm{Im}(\xi)) < \mathrm{rk}(E)$. Therefore the saturation $\overline{\mathrm{Im}(\xi)}$ is a proper subsheaf of E'. Let

$$\overline{\Phi'}_i : \overline{\mathrm{Im}(\xi)} \otimes_X M_i \xrightarrow{\Phi'_i|_{\overline{\mathrm{Im}(\xi)}\otimes_X M_i}} E' \twoheadrightarrow E'/\overline{\mathrm{Im}(\xi)}$$

be morphisms induced by Φ'_i, $i = 1, 2$. For generic $x \in X$, $\mathrm{Im}(\xi)_x = \overline{\mathrm{Im}(\xi)}_x$, therefore relations (2.2) imply that $\mathrm{Im}(\overline{\Phi}_i)_x = 0$. However the quotient $E'/\overline{\mathrm{Im}(\xi)}$ is torsion free since $\overline{\mathrm{Im}(\xi)}$ is saturated, hence $\mathrm{Im}(\overline{\Phi'}_i)$ must be trivial. This implies that

$$\Phi'_i(\overline{\mathrm{Im}(\xi)} \otimes_X M_i) \subseteq \overline{\mathrm{Im}(\xi)}$$

Obviously we also have $\mathrm{Im}(\psi') \subseteq \overline{\mathrm{Im}(\xi)}$ according to (2.2). Therefore the stability condition implies that $\overline{\mathrm{Im}(\xi)}$ must be either trivial or equal to E', and we have reached a contradiction.

In conclusion, if $\xi : E \to E'$ is nontrivial, it must be injective. Then $\mathrm{Coker}(\xi)$ is a torsion sheaf on X. However, E, E' have identical Hilbert polynomials by assumption, therefore $\mathrm{Coker}(\xi)$ must be in fact trivial. Therefore if ξ is not trivial, it must be an isomorphism. This proves lemma (2.3.i).

In order to prove (2.3.ii), take $\mathcal{E} = \mathcal{E}'$. Lemma (2.3.$i$) implies that any nontrivial endomorphism $\xi : \mathcal{E} \to \mathcal{E}'$ must be an automorphism. Then a standard argument (see for example [40, Cor. 1.2.8]) implies that any such morphism must be of the form $\xi = \lambda 1_E$, $\lambda \in \mathbb{C}$ if k is algebraically closed. Moreover, since ξ is a morphism of ADHM sheaves, we must have in particular $\lambda\psi = \psi$. Since ψ is non-trivial by stability, it follows that $\lambda = 1$. This proves lemma (2.3.ii).

□

Lemma 2.4. *Let $\mathcal{E}$ be an ADHM sheaf on X (not necessarily stable). Then there exists a canonical saturated subsheaf $E_0 \subseteq E$ so that*

$$\Phi_i(E_0 \otimes_X M_i) \subseteq E_0, \qquad \mathrm{Im}(\psi) \subset E_0$$

for $i = 1, 2$.

Proof. If $\Phi_i = 0$, $i = 1, 2$, we take E_0 to be the saturation of $\mathrm{Im}(\psi)$ in E.

Now suppose Φ_1, Φ_2 are not simultaneously trivial. Let $I = (i_1, i_2, \ldots, i_k)$ be an ordered collection of indices $i_l \in \{1, 2\}$ for $l = 1, \ldots, k$. For each such ordered collection, let

$$M_I = M_{i_1} \otimes_X \cdots \otimes_X M_{i_k}$$

and

$$\Phi_I : E \otimes M_I \xrightarrow{\Phi_{i_k} \circ (\Phi_{i_{k-1}} \otimes 1_{M_k}) \circ \cdots \circ (\Phi_{i_1} \otimes 1_{M_{i_2} \otimes \cdots \otimes M_{i_k}})} E$$

To the empty collection $I = \emptyset$ we formally assign $M_I = \mathcal{O}_X$ and $\Phi_I = 1_E$. Then set

$$E_0' = \sum_I \Phi_I(\mathrm{Im}(\psi) \otimes_X M_I) \tag{2.3}$$

where the sum is over all finite collections I of arbitrary length. According to [32, 5.3.4], E_0' is a coherent submodule of E. We also have

$$\Phi_i(E_0') \subseteq E_0', \qquad \mathrm{Im}(\psi) \subseteq E_0'$$

by construction.

Let E_0 be the saturation of E_0' in E. Obviously, E_0 contains $\mathrm{Im}(\psi)$ as a subsheaf. If $E_0 = E$, then obviously $\Phi_i(E_0 \otimes_X M_i) \subseteq E_0$ for $i = 1, 2$. If $E_0 \neq E$, consider the following composition of morphisms of $\mathcal{O}_X$-modules

$$\Phi_{i,0} : E_0 \otimes_X M_i \xrightarrow{\Phi_i} E \twoheadrightarrow E/E_0.$$

for $i = 1, 2$. Since E_0 is the saturation of E_0' it follows that $\Phi_{i,0}$ vanishes generically on X, hence $\mathrm{Im}(\Phi_{i,0})$ must be a torsion $\mathcal{O}_X$-module. However E/E_0 is torsion free since E_0 is saturated, hence $\mathrm{Im}(\Phi_{i,0})$ must be trivial. Therefore $\Phi_{i,0} = 0$ and we have $\Phi_i(E_0 \otimes_X M_i) \subseteq E_0$ for $i = 1, 2$.

□

Lemma 2.5. *Suppose E_∞ is a simple locally free sheaf on X. Then any stable ADHM sheaf $\mathcal{E}$ on X must have $\phi = 0$.*

Proof. Since E_∞ is simple, the endomorphism algebra $\mathrm{End}(E_\infty)$ is isomorphic to $\mathbb{C}$. In particular, for any endomorphism $\xi_\infty : E_\infty \to E_\infty$, $\mathrm{Tr}(\xi_\infty) = 0$ implies $\xi_\infty = 0$. Then the inductive argument used in the proof of lemma [53, Lemma 2.8] applies without modification to any stable ADHM sheaf $\mathcal{E}$ with E_∞ simple. This proves that the restriction $\phi|_{E_0' \otimes_X M_{12}}$ is trivial, where E_0' is the subsheaf of E constructed in (2.3).

Then we claim that the restriction $\phi|_{E_0 \otimes_X M_{12}}$ also vanishes, where E_0 is the saturation of E_0' as in lemma (2.4). By construction, it is clear that the restriction $\phi|_{E_0 \otimes_X M_{12}}$ vanishes generically on X. Therefore its image $\mathrm{Im}\,(\phi|_{E_0 \otimes_X M_{12}})$ must be a torsion sheaf on X. This is impossible since E_∞ is locally free. Therefore $\phi|_{E_0 \otimes_X M_{12}}$ must be trivial. However, the stability condition implies that $E_0 = E$, hence $\phi = 0$.

□

2.3. **Boundedness.** Next we prove a boundedness result for stable ADHM sheaves with fixed Hilbert polynomial P. In this section we will take $k = \mathbb{C}$, hence X is a smooth complex projective variety.

Lemma 2.6. *Let E be a torsion-free sheaf of rank $r \geq 2$ on X. Suppose E is not semistable, and let*

$$0 = HN_0(E) \subset HN_1(E) \subset \ldots \subset HN_h(E) = E$$

be the Harder-Narasimhan filtration of E. Then

$$\mu(HN_1(E)) + (r-1)\mu(E/HN_{h-1}(E)) \leq r\mu(E). \tag{2.4}$$

Proof. This is a standard result used for example in the proof of [56, Prop. 3.2], [64, Thm. 3.1]. The proof relies on an inductive argument and will be left to the reader.

□

Proposition 2.7. *Let $\mathcal{A}_P$ be the set of torsion free sheaves E with fixed Hilbert polynomial P so that $(E, \Phi_{1,2}, \phi, \psi)$ is a stable ADHM sheaf for some morphisms $(\Phi_{1,2}, \phi, \psi)$. Then $\mathcal{A}_P$ is bounded.*

Proof. The proof will be based on Maruyama's theorem [49]

Theorem 2.8. (Maruyama). *A family of sheaves E with fixed Hilbert polynomial P and $\mu_{\max}(E) \leq C$ for a fixed constant C is bounded.*

Therefore it suffices to prove that there exists a constant C so that $\mu_{\max}(E) \leq C$ for all stable ADHM sheaves $\mathcal{E} = (E, \Phi_{1,2}, \phi, \psi)$ on X with Hilbert polynomial P. If E is semistable $\mu_{\max}(E) = \mu(E)$ is clearly bounded. In particular this is the case if $\mathrm{rk}(E) = 1$, hence we will assume $\mathrm{rk}(E) \geq 2$ from now on in this proof.

Suppose E is not semistable, and let

$$0 = HN_0(E) \subset HN_1(E) \subset \cdots \subset HN_h(E) = E, \tag{2.5}$$

$h \geq 2$, be the Harder-Narasimhan filtration of E. Note that the successive quotients are torsion free and semistable. Given two torsion free $\mathcal{O}_X$-modules, any extension

$$0 \to F_1 \to F \to F_2 \to 0$$

must be torsion free. Then a simple inductive argument shows that $HN_j(E)$, $j = 1, \ldots, h-1$ are saturated subsheaves of E i.e the quotients $E/HN_j(E)$ are torsion free for all $j = 1, \ldots, h-1$. In particular this implies that $h \leq r$.

Since $HN_j(E)$, $j = 1, \ldots, h-1$ are also nontrivial and proper subsheaves by construction, it follows from the stability condition that for each $j = 1, \ldots, h-1$ either

(a) there exists $i \in \{1, 2\}$ so that $\Phi_i(HN(E)_j \otimes M_i) \not\subseteq HN(E)_j$, or
(b) $\mathrm{Im}(\psi) \not\subseteq HN(E)_j$.

Let $j_\psi \in \{1, \ldots, h-1\}$ be the index characterized by

$$\begin{aligned} &\mathrm{Im}(\psi) \not\subseteq HN_j(E), && \text{for } j \leq j_\psi \\ &\mathrm{Im}(\psi) \subseteq HN_j(E), && \text{for } j \geq j_\psi + 1. \end{aligned}$$

Note that the morphism

$$\overline{\psi} : E_\infty \to E/HN_{j_\psi}(E)$$

is nontrivial and [40, Lemma 1.3.3] implies that

$$\mu_{\min}(E_\infty) \leq \mu_{\max}(E/HN_{j_\psi}(E)).$$

By construction (see the proof of [40, Thm. 1.3.4]) we have $\mu_{\max}(E/HN_{j_\psi}(E)) = \mu(HN_{j_\psi+1}(E)/HN_{j_\psi}(E))$, therefore we obtain

$$\mu_{\min}(E_\infty) \leq \mu(HN_{j_\psi+1}(E)/HN_{j_\psi}(E)). \tag{2.6}$$

Moreover, if $j_\psi = h-1$, inequality (2.6) specializes to

$$\mu_{\min}(E_\infty) \leq \mu(E/HN_{h-1}(E)).$$

which yields

$$-\mu(E/HN_{h-1}(E)) \leq -\mu_{\min}(E_\infty). \tag{2.7}$$

If $j_\psi < h-1$, the stability condition implies that for each $j \in \{j_\psi+1, \ldots, h-1\}$ there exists $i_j \in \{1, 2\}$ so that $\Phi_{i_j}(HN_j(E) \otimes M_{i_j}) \not\subseteq HN_j(E)$. Then the argument employed in the proof of [56, Prop 3.2] and [64, Thm. 3.1] shows that

$$\mu(HN_j(E)/HN_{j-1}(E)) \leq \mu(HN_{j+1}(E)/HN_j(E)) - \deg(M_{i_j}). \tag{2.8}$$

Summing inequalities (2.8) from $j = j_\psi + 1$ to $j = h - 1$ we obtain

$$\mu(HN_{j_\psi+1}(E)/HN_{j_\psi}(E)) \leq \mu(E/HN_{h-1}(E)) - \sum_{j=j_\psi+1}^{h-1} \deg(M_{i_j}).$$

Then using inequality (2.6) we obtain

$$\mu_{\min}(E_\infty) + \sum_{j=j_\psi+1}^{h-1} \deg(M_{i_j}) \leq \mu(E/HN_{h-1}(E)).$$

which further yields

$$\mu_{\min}(E_\infty) - (h-1)\max\{|\deg(M_1)|, |\deg(M_2)|\} \leq \mu(E/HN_{h-1}(E)).$$

Since we have established above that $h \leq r$, we finally obtain

$$(2.9)\quad -\mu(E/HN_{h-1}(E)) \leq -\mu_{\min}(E_\infty) + (r-1)\max\{|\deg(M_1)|, |\deg(M_2)|\}$$

Taking into account (2.7), (2.9), inequality (2.4) implies the existence of the required upper bound for $\mu(HN_1(E)) = \mu_{\max}(E)$.

□

3. Moduli Space of Stable ADHM Sheaves

The goal of this section is to prove theorem (1.1). We will take X to be a smooth complex projective variety equipped with a very ample line bundle $\mathcal{O}_X(1)$. Let M_1, M_2, E_∞ be fixed twisting, respectively framing data on X. We begin with the construction of the groupoid of stable ADHM sheaves.

Definition 3.1. *(i) A flat family of ADHM sheaves on X parameterized by S is an ADHM sheaf $\mathcal{E}_S$ on X_S with twisting data $(M_1)_S, (M_2)_S$ and framing data $(E_\infty)_S$ so that E_S is flat over S and the the restriction $\mathcal{E}_S|_{X_sS}$ is a stable ADHM sheaf on X with fixed Hilbert polynomial P for any point $s \in S$.*

(ii) Two flat families of ADHM sheaves $\mathcal{E}_S = (E_S, \Phi_{S,1,2}, \phi_S, \psi_S)$ $\mathcal{E}'_S = (E'_S, \Phi'_{S,1,2}, \phi'_S, \psi'_S)$ are isomorphic if they are isomorphic as ADHM sheaves on X_S.

For fixed X, M_1, M_2, E_∞ and fixed Hilbert polynomial P, we define a fibered category $\mathfrak{p} : \mathfrak{M}_{ADHM}(\mathcal{X}, P) \to \mathfrak{S}$ over $\mathfrak{S}$ as follows

- The objects of $\mathfrak{M}_{ADHM}(\mathcal{X}, P)$ are flat families $\mathcal{E}_S$ of stable ADHM sheaves on X with Hilbert polynomial P parameterized by schemes S of finite type over $\mathbb{C}$.
- Given two objects $\mathcal{E}_S$ and $\mathcal{E}_T$, a morphism in $\mathrm{Mor}_{\mathfrak{M}_{ADHM}(\mathcal{X},P)}(\mathcal{E}_S, \mathcal{E}_T)$ is a pair (f, ξ_S) where $f : S \to T$ is a morphism of schemes of finite type and ξ_S is an isomorphism

$$\xi_S : \mathcal{E}_S \xrightarrow{\sim} f_X^*\mathcal{E}_T.$$

The projection functor $\mathfrak{p} : \mathfrak{M}_{ADHM}(\mathcal{X}, P) \to \mathfrak{S}$ assigns the parameter scheme S to a family $\mathcal{E}_S$ and the morphism $f : S \to T$ to a pair (f, ξ_S).

Proposition 3.2. $\mathfrak{p} : \mathfrak{M}_{ADHM}(\mathcal{X}, P) \to \mathfrak{S}$ *is a category fibered in groupoids.*

Proof. Follows from general properties of the pull-back of coherent sheaves and morphisms of coherent sheaves.

□

Our next goal is to prove that $\mathfrak{M}_{ADHM}(\mathcal{X}, P)$ is an Artin stack. The main step is the construction of a parameter space for stable $ADHM$ sheaves on X with fixed Hilbert polynomial P.

3.1. **The Parameter Space.** Recall that the set $\mathcal{A}_P$ defined in proposition (2.7) is bounded. Then there exists an integer m_0 sufficiently large so that for all $m \geq m_0$

(A) $E(m)$, $E(m)\otimes_X M_1^{-1}$, $E(m)\otimes_X M_2^{-1}$ are generated by global sections and $H^k(X, E(m)) = 0$, $H^k(X, E(m)\otimes_X M_1^{-1}) = 0$, $H^k(X, E(m)\otimes_X M_2^{-1}) = 0$ for all $k \geq 1$ and for all E in $\mathcal{A}_P$.

(B) $E_\infty(m)$, $E_\infty(m) \otimes_X M_{12}^{-1}$ are generated by global sections and $H^k(X, E_\infty(m)) = 0$, $H^k(X, E_\infty(m) \otimes_X M_{12}^{-1}) = 0$ for all $k \geq 1$.

Fix $m \geq m_0$ and let V be a complex vector space of dimension $N = P(m)$. Let $V_\infty = H^0(X, E_\infty(m))$, and $V_{\infty,12} = H^0(X, E_\infty \otimes_X M_{12}^{-1})$. Note that condition ($B$) implies that we can choose a surjective morphism

$$q_\infty : V_\infty \otimes \mathcal{O}_X(-m) \twoheadrightarrow E_\infty$$

so that $H^0(q_\infty(m))$ is an isomorphism. We will fix such a morphism in the following.

Definition 3.3. *An ADHM quotient of type* (P, m) *is defined as the set of data* $(\mathcal{E}, q, \Xi_{1,2}, \xi, \eta)$ *where* $\mathcal{E}$ *is a stable ADHM sheaf on* X *with Hilbert polynomial* P*,*

$$q : V \otimes \mathcal{O}_X(-m) \twoheadrightarrow E$$

is a surjective morphism of $\mathcal{O}_X$ *modules, and*

$$\begin{aligned} \Xi_i &: V \to H^0(X, E(m) \otimes_X M_i^{-1}) \\ \xi &: V \to H^0(X, E_\infty(m) \otimes_X M_{12}^{-1}) \\ \eta &: V_\infty \to H^0(X, E(m)) \end{aligned}$$

are linear maps so that

(i) $H^0(q(m)) : V \to H^0(X, E(m))$ *is an isomorphism.*

(ii) The following factorizations hold

(3.1)

$$\begin{tikzcd}
V\otimes\mathcal{O}_X \arrow[r, "q(m)"] \arrow[d, "\Xi_i\otimes 1_{\mathcal{O}_X}"'] & E(m) \arrow[ddl, bend left, "\Phi_i(m)"] \\
H^0(X, E(m)\otimes_X M_i^{-1})\otimes\mathcal{O}_X \arrow[d, "ev"'] & \\
E(m)\otimes_X M_i^{-1} &
\end{tikzcd}
\qquad
\begin{tikzcd}
V_\infty\otimes\mathcal{O}_X \arrow[r, "q_\infty(m)"] \arrow[d, "\eta\otimes 1_{\mathcal{O}_X}"'] & E_\infty(m) \arrow[ddl, bend left, "\psi(m)"] \\
H^0(X, E(m))\otimes\mathcal{O}_X \arrow[d, "ev"'] & \\
E(m) &
\end{tikzcd}$$

$$\begin{tikzcd}
V\otimes\mathcal{O}_X \arrow[r, "q(m)"] \arrow[d, "\xi\otimes 1_{\mathcal{O}_X}"'] & E(m) \arrow[ddl, bend left, "\phi(m)"] \\
H^0(X, E_\infty(m)\otimes_X M_{12}^{-1})\otimes\mathcal{O}_X \arrow[d, "ev"'] & \\
E_\infty(m)\otimes_X M_{12}^{-1} &
\end{tikzcd}$$

for $i = 1, 2$.

Definition 3.4. *Two ADHM quotients of type (P, m) $(\mathcal{E}, q, \Xi_{1,2}, \xi, \eta)$ and $(\mathcal{E}', q', \Xi'_{1,2}, \xi', \eta')$ are isomorphic if there exists an isomorphism of ADHM sheaves $\rho : \mathcal{E}\xrightarrow{\sim}\mathcal{E}'$ so that the following diagrams are commutative*

$$\begin{tikzcd}
V\otimes\mathcal{O}_X(-m) \arrow[r, "q"] \arrow[d, "1_V\otimes 1_{\mathcal{O}_X(-m)}"'] & E \arrow[d, "\rho"] \\
V\otimes\mathcal{O}_X(-m) \arrow[r, "q'"] & E'
\end{tikzcd}
\qquad
\begin{tikzcd}
V \arrow[r, "\Xi_i"] \arrow[d, "1_V"'] & H^0(X, E(m)\otimes_X M_i^{-1}) \arrow[d, "H^0\left(\rho(m)\otimes 1_{M_i^{-1}}\right)"] \\
V \arrow[r, "\Xi'_i"] & H^0(X, E'(m)\otimes_X M_i^{-1})
\end{tikzcd}$$

$$\begin{tikzcd}
V \arrow[r, "\xi"] \arrow[d, "1_V"'] & H^0(X, E(m)\otimes_X M_{12}^{-1}) \arrow[d, "H^0\left(\rho(m)\otimes 1_{M_{12}^{-1}}\right)"] \\
V \arrow[r, "\xi'"] & H^0(X, E'(m)\otimes_X M_{12}^{-1})
\end{tikzcd}
\qquad
\begin{tikzcd}
V_\infty \arrow[r, "\eta"] \arrow[d, "1_{V_\infty}"'] & H^0(X, E(m)) \arrow[d, "H^0(\rho(m))"] \\
V_\infty \arrow[r, "\eta'"] & H^0(X, E'(m))
\end{tikzcd}$$

for $i = 1, 2$.

Definition 3.5. *A flat family of ADHM quotients of type (P, m) parameterized by a scheme S of finite type over $\mathbb{C}$ is defined as the set of data $(\mathcal{E}_S, q_S, \Xi_{S,1,2}, \xi_S, \eta_S)$ where $\mathcal{E}_S$ is a flat family of stable ADHM sheaves on X parameterized by S,*

$$q_S : V\otimes\mathcal{O}_X(-m)_S \twoheadrightarrow E_S$$

is a surjective morphism of $\mathcal{O}_{X_S}$-modules, and

$$\begin{aligned}\Xi_{S,i} &: V\otimes \mathcal{O}_S \to p_{S*}(E_S\otimes_{X_S}(\mathcal{O}_X(m)\otimes_X M_i^{-1})_S)\\ \xi_S &: V\otimes \mathcal{O}_S \to p_{S*}((E_\infty(m)\otimes_X M_{12}^{-1})_S)\\ \eta_S &: V_\infty\otimes \mathcal{O}_S \to p_{S*}(E_S\otimes_{X_S}\mathcal{O}_X(m)_S)\end{aligned}$$

are morphisms of $\mathcal{O}_S$-modules so that the restriction $(\mathcal{E}_S, q_S, \Xi_{S,1,2}, \xi_S, \eta_S)|_{X_s}$ is an ADHM quotient of type (P,m) for any point $s\in S$.

Two flat families of ADHM quotients $(\mathcal{E}_S, q_S, \Xi_{S,1,2}, \xi_S, \eta_S)$, $(\mathcal{E}_S, q_S, \Xi_{S,1,2}, \xi_S, \eta_S)'$ are isomorphic if there exists an isomorphism $\rho_S : \mathcal{E}_S \to \mathcal{E}_S$ of flat families of stable ADHM sheaves so that the obvious diagrams analogous to those in definition (3.4) are commutative.

Let $\mathfrak{Q}_{ADHM}(X,P,m) : \mathfrak{S}\to Sets$ be the functor which assigns to any S the set of isomorphism classes of flat families of ADHM quotients of type (P,m) parameterized by S. Then we claim

Lemma 3.6. *The functor $\mathfrak{Q}_{ADHM}(X,P,m)$ is represented by a quasi-projective scheme R over $\mathbb{C}$ equipped with a universal flat family $(\mathcal{E}_R, q_R, \Xi_{R,1,2}, \xi_R, \eta_R)$ of ADHM quotients of type (P,m).*

Proof. This is a standard construction for moduli problems of decorated sheaves [39, 38, 65, 29, 66]. R will be constructed as a subscheme of an affine scheme over the *Quot*-scheme Q of coherent quotients $V\otimes\mathcal{O}_X(-m)\twoheadrightarrow E$ on X with Hilbert polynomial P. Note that Q is a projective scheme over $\mathbb{C}$. Let

$$q_Q : V\otimes p_X^*\mathcal{O}_X(-m)\twoheadrightarrow E_Q$$

denote the universal quotient on Q (see for example [40, Sect. 2.2] or [26, Ch. 5] for construction and existence proofs.)

The family of quotients $[q : V\otimes\mathcal{O}_X(-m)\twoheadrightarrow E]$ with fixed Hilbert polynomial is bounded [40, Lemma 1.7.6]. Therefore we can choose the integer m_0 sufficiently large so that condition (A) above holds for all quotients $[V\twoheadrightarrow E]$ parameterized by Q. We will assume this to be the case from this point on.

Given conditions (A) and (B) above, the base change theorem [34, Thm 7.7.5], [8, Thm. III.3.4] implies that all direct images

$$\begin{aligned}\mathcal{H} &= p_{Q*}(E_Q\otimes_{X_Q}\mathcal{O}_X(m)_Q), & \mathcal{H}_i &= p_{Q*}(E_Q\otimes_{X_Q}(M_i^{-1}(m))_Q),\\ \mathcal{H}_\infty &= p_{Q*}(E_\infty(m))_Q, & \mathcal{H}_{\infty,12} &= p_{Q*}(E_\infty(m)\otimes_X M_{12}^{-1})_Q,\end{aligned}$$

with $i=1,2$, are locally free $\mathcal{O}_Q$-modules. Let $\mathcal{V}$ denote the locally free $\mathcal{O}_Q$-module

$$\begin{aligned}\mathcal{V} &= V^\vee\otimes(\mathcal{H}_1\oplus\mathcal{H}_2\oplus\mathcal{H}_{\infty,12})\oplus V_\infty^\vee\otimes\mathcal{H}\\ &= V^\vee\otimes(\mathcal{H}_1\oplus\mathcal{H}_2)\oplus(V^\vee\otimes V_{\infty,12}\otimes\mathcal{O}_Q)\oplus V_\infty^\vee\otimes\mathcal{H}.\end{aligned}\tag{3.2}$$

Let $S(\mathcal{V}^\vee)$ denote the symmetric tensor algebra of $\mathcal{V}^\vee$ and let $T=\mathrm{Spec}S(\mathcal{V}^\vee)$. Note that we have an affine projection $p_{T/Q} : T\to Q$. The fiber of $p_{T/Q}$

over a point $[q : V \otimes \mathcal{O}_X(-m) \twoheadrightarrow E]$ of Q is isomorphic to the linear space

$$\begin{aligned} &\mathrm{Hom}\left(V, H^0(X, E(m) \otimes_X M_1^{-1}) \oplus H^0(X, E(m) \otimes_X M_2^{-1})\right) \\ &\oplus \mathrm{Hom}(V, V_{\infty,12}) \oplus \mathrm{Hom}(V_\infty, H^0(X, E(m))). \end{aligned} \tag{3.3}$$

Let $E_T = (p_{T/Q})_X^* E_Q$ and note that by flat base change we have

$$p_{T/Q}^* p_{Q*}(E_Q \otimes_{X_Q} F_Q) \simeq p_{T*}(E_T \otimes_{X_T} F_T) \tag{3.4}$$

for any coherent $\mathcal{O}_X$-module F since $p_{T/Q} : T \to Q$ is affine, hence flat. Moreover we also have a tautological section τ of $p_{T/Q}^*\mathcal{V}$ over T. Taking into account (3.4), τ yields canonical morphisms of $\mathcal{O}_T$-modules

$$\begin{aligned} \Xi_{T,i} &: V \otimes \mathcal{O}_T \to p_{T*}(E_T \otimes_{X_T} (\mathcal{O}_X(m) \otimes_X M_i^{-1})_T) \\ \xi_T &: V \otimes \mathcal{O}_T \to p_{T*}(E_\infty(m) \otimes_X M_{12}^{-1})_T) \\ \eta_T &: V_\infty \otimes \mathcal{O}_T \to p_{T*}(E_T \otimes_{X_T} \mathcal{O}_X(m)_T) \end{aligned} \tag{3.5}$$

Then we obtain the following tautological diagrams on X_T

$$\begin{tikzcd} V \otimes \mathcal{O}_{X_T} \arrow[r, "p_T^*\Xi_{T,i}"] \arrow[dr, "\Lambda_{T,i}"'] & p_T^*p_{T*}(E_T \otimes_{X_T} (\mathcal{O}_X(m) \otimes_X M_i^{-1})_T) \arrow[d, "\mathrm{ev}"] \\ & E_T \otimes_{X_T} (\mathcal{O}_X(m) \otimes_X M_i^{-1})_T \end{tikzcd}$$

$$\begin{tikzcd} V \otimes \mathcal{O}_{X_T} \arrow[r, "p_T^*\xi_T"] \arrow[dr, "\lambda_T"'] & p_T^*p_{T*}(E_\infty(m) \otimes_X M_{12}^{-1})_T \arrow[d, "\mathrm{ev}"] \\ & (E_\infty(m) \otimes_X M_{12}^{-1})_T \end{tikzcd}$$

$$\begin{tikzcd} V_\infty \otimes \mathcal{O}_{X_T} \arrow[r, "p_T^*\eta_T"] \arrow[dr, "\varrho_T"'] & p_T^*p_{T*}(E_T \otimes_{X_T} \mathcal{O}_X(m)_T) \arrow[d, "\mathrm{ev}"] \\ & E_T \otimes_{X_T} \mathcal{O}_X(m)_T \end{tikzcd}$$

where $i = 1, 2$.

In order to obtain a parameter space for stable ADHM sheaves, we have to establish the existence of a closed subscheme of $Z \subset T$ so that the morphisms $\Lambda_{T,1},\ \Lambda_{T,2}\ \lambda_T,\ \varrho_T$ factor through the projections

$$\begin{aligned} (p_{T/Q})_X^* q_Q(m)|_Z &: V \otimes \mathcal{O}_Z \to E_T \otimes_{X_T} \mathcal{O}_X(m)_T|_Z \\ (p_{T/Q})_X^* q_Q(m)|_Z &: V \otimes \mathcal{O}_Z \to E_T \otimes_{X_T} \mathcal{O}_X(m)_T|_Z \\ (p_{T/Q})_X^* q_Q(m)|_Z &: V \otimes \mathcal{O}_Z \to E_T \otimes_{X_T} \mathcal{O}_X(m)_T|_Z \\ (q_\infty(m))_T|_Z &: V_\infty \otimes \mathcal{O}_Z \to E_\infty(m)_T|_Z \end{aligned}$$

respectively, for $i = 1, 2$.

This follows for example from [29, Lemma 3.1], which is reproduced below for convenience.

Lemma 3.7. *Let Y be a scheme, and let $f : G \to F$ be a homomorphism of coherent $\mathcal{O}_{X\times Y}$-modules so that F is flat over Y. Then there is a unique closed subscheme Z of Y satisfying the following universal property: given a Cartesian diagram*

$$\begin{tikzcd}
X_S \arrow[r, "\overline{h}"] \arrow[d, "p_S"'] & X_Y \arrow[d, "p_Y"] \\
S \arrow[r, "h"] & Y
\end{tikzcd}$$

$\overline{h}^* f = 0$ if and only if h factors through Z.

In lemma (3.7) take $Y = T$,

$$G = \mathrm{Ker}\left((p_{T/Q})_X^*(q_Q(m)^{\oplus 3}) \oplus p_X^* q_\infty(m)\right)$$

$$F = E_T \otimes_{X_T} (M_1^{-1}(m) \oplus M_2^{-1}(m))_T \oplus (E_\infty(m) \otimes_X M_{12}^{-1})_T \\ \oplus E_T \otimes_{X_T} \mathcal{O}_X(m)_T$$

and

$$f = (\Lambda_{T,1} \oplus \Lambda_{T,2} \oplus \lambda_T \oplus \varrho_T)|_G.$$

Then we obtain a closed subscheme $Z \subset T$ with the desired properties. Let $E_Z = E_T|_{X_Z}$. By construction we have morphisms of $\mathcal{O}_{X_Z}$-modules $\Phi_{Z,i} : E_Z \otimes_{X_Z} (M_i)_Z \to E_Z$, $i = 1, 2$, $\phi_Z : E_Z \otimes_{X_Z} (M_{12}) \to (E_\infty)_Z$, $\psi_Z : (E_\infty)_Z \to E_Z$ and data

$$\begin{aligned}
q_Z &: V \otimes \mathcal{O}_X(-m)_Z \twoheadrightarrow E_Z \\
\Xi_{Z,i} &: V \otimes \mathcal{O}_Z \to p_{Z*}(E_Z \otimes_{X_Z} (\mathcal{O}_X(m) \otimes_X M_i^{-1})_Z) \\
\xi_Z &: V \otimes \mathcal{O}_Z \to p_{Z*}(E_Z \otimes_{X_Z} (\mathcal{O}_X(m) \otimes_X M_{12}^{-1})_Z) \\
\eta_Z &: V_\infty \otimes \mathcal{O}_Z \to p_{Z*}(E_S \otimes_{X_Z} \mathcal{O}_X(m)_Z)
\end{aligned}$$

so that the factorizations required by definition (3.5) hold.

Next, let $R \subset Z$ be the subscheme of Z so that the restriction of the data $(\mathcal{E}_Z, q_Z, \Xi_{Z,1,2}, \xi_Z, \eta_Z)$ to X_z is an ADHM quotient of type (P, m) on X_z for any closed point $z \in R$. Since the factorizations required in definition (3.3) hold by construction, this means we just have to require the data $(E_Z, \Phi_{Z,1,2}, \phi_Z, \psi_Z)|_z$ to be a stable ADHM sheaf on X_z for any $z \in R$, and $H^0(q_Z(m)|_{X_z})$ to be an isomorphism. Restricting the data $(\mathcal{E}_Z, q_Z, \Xi_{Z,1,2}, \xi_Z, \eta_Z)$ to R, we obtain a flat family of ADHM quotients of type (P, m) $(\mathcal{E}_R, q_R, \Xi_{R,1,2}, \xi_R, \eta_R)$ on X parameterized by R.

We claim that R is the parameter space for ADHM quotients required in proposition (3.6). In order to prove this, suppose $(\mathcal{E}_S, q_S, \Xi_{S,1,2}, \xi_S, \eta_S)$ is a flat family of ADHM quotients of type (P, m) parameterized by S. In particular we have a flat $\mathcal{O}_{X_S}$-module E_S so that $H^0(X, q_S|_{X_s})$ is an isomorphism for each $s \in S$. Therefore we obtain a classifying map $g : S \to Q$ so that $g_X^* E_Q \simeq E_S$.

Taking into account conditions (A), (B) above (3.3), the base change theorem [34, Thm 7.7.5], [8, Thm. III.3.4] yields isomorphisms

$$\begin{gathered} g^*p_{Q*}(E_Q \otimes_{X_Q} (\mathcal{O}_X(m) \otimes_X M_i^{-1})_Q) \simeq p_{S*}(E_S \otimes_{X_S} (\mathcal{O}_X(m) \otimes_X M_i^{-1})_S) \\ g^*p_{Q*}(E_\infty(m) \otimes_X M_{12}^{-1})_Q \simeq p_{S*}(E_\infty(m) \otimes_X M_{12}^{-1})_S \\ g^*p_{Q*}(E_Q \otimes_{X_Q} \mathcal{O}_X(m)_Q) \simeq p_{S*}(E_S \otimes_{X_S} \mathcal{O}_X(m)_S) \end{gathered}$$

for $i = 1, 2$. Then it follows for example from [26, Thm. 5.8] that there is a lift $h : S \to T$ so that

$$(E_S, \Xi_{S,1,2}, \xi_S, \eta_S) = h^*(E_T, \Xi_{T,1,2}, \xi_T, \eta_T).$$

Since $(\mathcal{E}_S, q_S, \Xi_{S,1,2}, \xi_S, \eta_S)$ is a flat family of ADHM quotients, lemma (3.7) implies that h factors through a morphism $f : S \to R$. Moreover, we have

$$(E_S, \Xi_{S,1,2}, \xi_S, \eta_S) = f^*(E_R, \Xi_{R,1,2}, \xi_R, \eta_R).$$

Finally, since the ADHM data $\mathcal{E}_S = (E_S, \Phi_{S,1,2}, \phi_2, \psi_S)$ is determined by the data $(E_S, \Xi_{S,1,2}, \xi_S, \eta_S)$ for any base scheme S, it follows that

$$(\mathcal{E}_S, \Xi_{S,1,2}, \xi_S, \eta_S) = f^*(\mathcal{E}_R, \Xi_{R,1,2}, \xi_R, \eta_R).$$

□

Note that there is a natural algebraic group action of $G = GL(V)$ on R so that the universal ADHM quotient $(\mathcal{E}_R, q_R, \Xi_{R,1,2}, \xi_R, \eta_R)$ is G-equivariant. We will prove below that the quotient stack $[R/G]$ is isomorphic to the stack of stable ADHM sheaves on X. First we need the following lemma.

Lemma 3.8. *Let X be an arbitrary scheme over $\mathbb{C}$, $f : P \to B$ be a principal G-bundle where G is a smooth algebraic group scheme over $\mathbb{C}$. Let G act trivially on X.*

(i) Let E_P be a coherent $\mathcal{O}_{X_P}$-module. Suppose E_P is equivariant with respect to the natural action of G on X_P. Then E_P descends to a coherent $\mathcal{O}_{X_B}$-module E_B. Moreover, if E_P is flat over P then E_B is flat over B.

(ii) Let $\phi_P : E_P \to F_P$ be a morphism of coherent $\mathcal{O}_{X_P}$-modules. Suppose E_P, F_P, ϕ_P are equivariant with respect to the natural action of G on X_P. Then $\phi_P : E_P \to F_P$ descends to a morphism of coherent $\mathcal{O}_{X_B}$-modules $\phi_B : E_B \to F_B$.

Proof. This lemma is a corollary of [1, Thm. 3.3]. Alternatively, one can give a direct proof by analogy with the proof of [40, Thm. 4.2.14]. First note that for a principal G-bundle $f : P \to B$ as above we have a canonical isomorphism

$$\sigma \times q_P : P \times G \xrightarrow{\sim} P \times_B P$$

where $\sigma : P \times G \to P$ denotes the right action of G on P and $q_P : P \times G \to P$ is the canonical projection. Then the main observation is that an equivariant structure of a coherent $\mathcal{O}_{X_P}$-module E_P is equivalent to descent data for E_P with respect to the morphism $f_X : X_P \to X_B X$. Therefore lemma (3.8) follows from the descent theorem for faithfully flat morphisms [26, Thm. 4.21], [9, Thm. A.2(a)] and [35, Prop. 2.5.1].

□

Proposition 3.9. *There is an isomorphism of groupoids* $[R/G] \simeq \mathfrak{M}_{ADHM}(\mathcal{X}, P)$ *over* $\mathfrak{S}$. *In particular,* $\mathfrak{M}_{ADHM}(\mathcal{X}, P)$ *is an Artin stack.*

Proof. The proof is very similar to the proof of [28, Prop. 3.3]. Let

$$\begin{tikzcd} P \arrow[r, "f"] \arrow[d] & R \\ B & \end{tikzcd}$$

be an object of $[R/G]$ over B. Then $(f \times 1_X)^*(E_R, \Phi_{R,1,2}, \phi_R, \psi_R)$ is a flat family of stable ADHM sheaves on X parameterized by P. Since f is G-equivariant, this family is equivariant with respect to the natural right G-action on X_P. Since $X_P \to X_B$ is a principal G-bundle, the family $f_X^*(E_R, \Phi_{R,1,2}, \phi_R, \psi_R)$ descends to a similar family $\mathcal{E}_B$ parameterized by B according to lemma (3.8).

A morphism between two objects

$$\begin{tikzcd} P \arrow[r, "f"] \arrow[d] & R & P' \arrow[r, "f'"] \arrow[d] & R \\ B & & B' & \end{tikzcd}$$

of $[R]$ is a commutative diagram of the form

(3.6)
$$\begin{tikzcd} & & R \\ P \arrow[r, "\xi"'] \arrow[rru, "f"] \arrow[d] & u^*P' \arrow[ru, "u^*f'"'] \arrow[d] & \\ B \arrow[r, "1_B"] & B & \end{tikzcd}$$

where $u : B \to B'$ is a morphism of schemes of finite type over $\mathbb{C}$, and $\xi : P \to u^*P'$ is an isomorphism. Since $u^*f' \circ \xi = f$, and all morphisms are G-equivariant, it follows that we have an isomorphism $f_X^*\mathcal{E}_R \xrightarrow{\sim} u_X^* f'^*_X \mathcal{E}_R$ which descends to an isomorphism $\mathcal{E}_B \xrightarrow{\sim} u_X^*\mathcal{E}_{B'}$ according to lemma (3.8). Therefore we obtain a functor $\mathfrak{f} : [R/G] \to \mathfrak{M}_{ADHM}(\mathcal{X}, P)$.

Next we construct a functor $\mathfrak{g} : \mathfrak{M}_{ADHM}(\mathcal{X}, P) \to [R/G]$ employing a construction similar to the proof of [40, Lemma 4.3.1]. Suppose $\mathcal{E}_B$ is a flat family of stable ADHM sheaves with Hilbert polynomial P on X parameterized by B. By definition, $\mathcal{E}_B|_{X_b}$ is a stable ADHM sheaf on X with Hilbert polynomial P for any point $b \in B$. Therefore it satisfies condition (A) above definition (3.3). Then the base change theorem [34, Thm 7.7.5], [8, Thm.

III.3.4] implies that the direct images

$$\begin{aligned} \mathcal{H} &= p_{B*}(E_B \otimes_{X_B} \mathcal{O}_X(m)_B), & \mathcal{H}_i &= p_{B*}(E_B \otimes_{X_B} M_i^{-1}(m)_B), \\ \mathcal{H}_\infty &= p_{B*}E_\infty(m)_B, & \mathcal{H}_{\infty,12} &= p_{B*}(E_\infty(m) \otimes_X M_{12}^{-1})_B, \end{aligned}$$

$i = 1, 2$, are locally free and all higher direct images vanish. Let $P = Isom(V \otimes \mathcal{O}_B \to \mathcal{H})$ denote the frame bundle associated to the locally free $\mathcal{O}_B$-module $\mathcal{H}$, which has rank $N = P(m)$ [40, Ex. 4.2.3]. Then we have a canonical surjective morphism of $\mathcal{O}_{X_B}$-modules

$$p_B^*\mathcal{H} \otimes_{X_B} \mathcal{O}_X(-m)_B \twoheadrightarrow E_B. \tag{3.7}$$

Let $\pi : P \to B$ denote the natural projection map. Note that the pullback $\pi^*\mathcal{H}$ has a canonical trivialization

$$V \otimes \mathcal{O}_P \xrightarrow{\sim} \pi^*\mathcal{H} \tag{3.8}$$

on P. Composing (3.8) and (3.7) we obtain a surjective morphism of $\mathcal{O}_{X_P}$-modules

$$q_P : V \otimes \mathcal{O}_X(-m)_P \twoheadrightarrow \pi_X^* E_B.$$

Let $E_P = (\pi \times 1_X)^* E_B$. Given condition (A) above (3.3), the base change theorem [34, Thm 7.7.5], [8, Thm. III.3.4] implies that the have an isomorphism

$$\pi^*\mathcal{H} \simeq p_{P*}(E_P \otimes_{X_P} \mathcal{O}_X(m)_P).$$

Moreover, it also implies that

$$p_{P*}(E_P \otimes_{X_P} \mathcal{O}_X(m)_P)|_{pt} \simeq H^0(X, E_P|_{\{pt\}\times X}(m))$$

for any point $\{pt\} \in P$. Then it follows that

$$(q_P|_{\{pt\}\times X})(m) : V \otimes \mathcal{O}_{\{pt\}\times X} \to E_P|_{\{pt\}\times X}(m)$$

induces an isomorphism in degree zero cohomology for any point $pt \in P$. This allows us to construct a flat family of ADHM quotients of type m parameterized by P as follows. Let $\mathcal{E}_P = \pi_X^* \mathcal{E}_B$. According to the base change theorem we have isomorphisms

$$\begin{aligned} \pi^*\mathcal{H} &\simeq p_{P*}(E_P \otimes_{X_P} \mathcal{O}_X(m)_P) \\ \pi^*\mathcal{H}_i &\simeq p_{P*}(E_P \otimes_{X_P} M_i^{-1}(m)_P) \\ \pi^*\mathcal{H}_{\infty,12} &\simeq p_{P*}(E_\infty(m) \otimes_X M_{12}^{-1})_P \end{aligned}$$

Then the data $(\Xi_{P,i}, \xi_P, \eta_P)$ is determined by applying p_{P*} to the following morphisms

$$\begin{tikzcd} V \otimes \mathcal{O}_{X_P} \arrow[r, "q_P(m)"] & E_P \otimes_X \mathcal{O}_X(m)_P \arrow[r, "\Phi_{P,i}(m)"] & E_P \otimes_X M_i^{-1}(m)_P \\ V \otimes \mathcal{O}_{X_P} \arrow[r, "q_P(m)"] & E_P \otimes_X \mathcal{O}_X(m)_P \arrow[r, "\phi_P(m)"] & (E_\infty(m) \otimes_X M_{12}^{-1})_P \\ V_\infty \otimes \mathcal{O}_{X_P} \arrow[r, "p_X^* q_\infty(m)"] & E_\infty(m)_P \arrow[r, "\psi_P(m)"] & E_P \otimes_X \mathcal{O}_X(m)_P \ . \end{tikzcd}$$

Therefore we obtain a flat family of ADHM quotients of type (P, m) parameterized by P. This family is naturally G-equivariant by construction, therefore it determines a G-equivariant classifying morphism $f : P \to R$.

Let (u, ϱ) be a morphism between two flat families of stable ADHM sheaves $\mathcal{E}_B, \mathcal{E}_{B'}$ where $u : B \to B'$ and $\varrho : \mathcal{E}_B \xrightarrow{\sim} u^*\mathcal{E}_{B'}$. Then repeating the above construction we obtain two flat families of ADHM quotients $(\mathcal{E}_P, \Xi_{P,1,2}, \xi_P, \eta_P)$, $(\mathcal{E}_P, \Xi_{P,1,2}, \xi_P, \eta_P)'$ over B and B' respectively. Since we have the isomorphism $\varrho : \mathcal{E}_B \xrightarrow{\sim} u^*\mathcal{E}_{B'}$, we will obtain an isomorphism $\xi : P \xrightarrow{\sim} u^*P'$ over B using the base change theorem.

Let $v : u^*P' \to P'$ be the morphism defined by the commutative diagram

$$\begin{tikzcd}
u^*P' \arrow[r, "v"] \arrow[d] & P' \arrow[d] \\
B \arrow[r, "u"] & B'
\end{tikzcd}$$

Employing again the base change theorem, it is straightforward to check that the families of ADHM quotients $(\mathcal{E}_P, \Xi_{P,1,2}, \xi_P, \eta_P)$, $\xi^*v^*(\mathcal{E}_P, \Xi_{P,1,2}, \xi_P, \eta_P)'$ are canonically isomorphic. Therefore the G-equivariant classifying maps $f : P \to R$, $f' : u^*P' \to R$ fit in a diagram of the form (3.6). In conclusion, we obtain a functor $\mathfrak{g} : \mathfrak{M}_{ADHM}(\mathcal{X}, P) \to [R/G]$ over S.

Finally, it is straightforward to check that the $\mathfrak{f} \circ \mathfrak{g}$, $\mathfrak{g} \circ \mathfrak{f}$ are naturally isomorphic to the identity functors, hence we obtain indeed an isomorphism of groupoids $[R/G] \simeq \mathfrak{M}_{ADHM}(\mathcal{X}, P)$.

□

3.2. **The Moduli Space.** In this section we prove theorem (1.1) and corollary (1.2). We will need the following preliminary lemma.

Lemma 3.10. *Let $\rho : A \to B$ be a local morphism of Noetherian local rings, k the residual field of A and M, N two finitely generated B-modules, N flat over A. Let $u : M \to N$ a morphism of B-modules. Then the following conditions are equivalent*

- (i) *u is injective and* Coker(u) *is flat over A*
- (ii) *$u \otimes 1 : M \otimes_A k \to N \otimes_A k$ is injective.*

Proof. [33, 10.2.4]

□

Proof of theorem (1.1). Lemma (2.3.*ii.*) and [44, Cor. 8.1.1] imply that $\mathfrak{M}_{ADHM}(\mathcal{X}, P)$ is an algebraic space over $\mathbb{C}$.

We will prove that $\mathfrak{M}_{ADHM}(\mathcal{X}, P)$ is separated using the valuative criterion of separatedness [44, Prop. 7.8]. Note that $\mathfrak{M}_{ADHM}(\mathcal{X}, P)$ is locally of finite type over $\mathbb{C}$ since the smooth atlas R provided by proposition (3.9) is quasi-projective. Then we can restrict ourselves to discrete valuation rings with residual field isomorphic to $\mathbb{C}$.

Therefore let R be a discrete valuation ring with residual field $k = \mathbb{C}$, and let K be its fraction field. We will denote by $\pi \in R$ a uniformizing element.

Let $\mathcal{E}_R^{(1)}, \mathcal{E}_R^{(2)}$ be flat families of stable ADHM sheaves on X parameterized by $\mathrm{Spec}(R)$ so that the restrictions $\mathcal{E}_K^{(1)}$, $\mathcal{E}_K^{(2)}$ to $\mathrm{Spec}(K)$ are isomorphic. Let $\xi_K : \mathcal{E}_K^{(1)} \to \mathcal{E}_K^{(2)}$ be an isomorphism of flat families of ADHM sheaves. We will denote $X \times \mathrm{Spec}(R)$ by X_R, and $X \times \mathrm{Spec}(K)$ by X_K. Therefore we are given an isomorphism of $\mathcal{O}_{X_K}$-modules $\xi_K : E_K^{(1)} \to E_K^{(2)}$ so that the diagrams (2.1) for $\mathcal{E}_K^{(1)}$, $\mathcal{E}_K^{(2)}$ and ξ_K are commutative.

First we claim that there exists an integer $n \in \mathbb{Z}$ so that $\pi^n \xi_K$ extends to a morphism of $\mathcal{O}_{X_R}$-modules $\xi_R : E_S^{(1)} \to E_S^{(2)}$ so that the restriction of ξ_R to any fiber of $X_R \to \mathrm{Spec}(R)$ is nontrivial. This follows from an argument analogous to the one used in the proof of [2, Lemma 7.8]. We will reproduce it below for completeness.

Let S be an arbitrary base scheme of finite type over $\mathbb{C}$, and F, G be coherent $\mathcal{O}_{X_S}$-modules F, G, with G flat over S. According to [2, Sect. 1.1], the functor

$$I \to \mathrm{Hom}_{X_S}(F, G \otimes_{X_S} p_S^* I)$$

from quasi-coherent $\mathcal{O}_S$-modules to sets is represented by the coherent $\mathcal{O}_S$-module $p_{S*}\mathcal{H}om_{X_S}(F,G)$ and the tautological morphism

$$h(F,G) \in \mathrm{H}om_{X_S}(F, G \otimes_{X_S} p_S^* p_{S*}\mathcal{H}om_{X_S}(F,G)).$$

Then for any scheme $f : T \to S$ over S there is a functorial isomorphism

$$\mathrm{Hom}_T(f^* p_{S*}\mathcal{H}om_{X_S}(F,G), \mathcal{O}_T) \simeq \mathrm{Hom}_{X_T}(f_X^* F, f_X^* G)$$

as in equation (1.1.1) of [2]. Taking $S = \mathrm{Spec}(R)$, $T = \mathrm{Spec}(K)$ and $f : T \to S$ the canonical open immersion, it follows that ξ_K corresponds to a nonzero element in

$$\eta_K \in \mathrm{Hom}_T(f^* p_{S*}\mathcal{H}om_{X_S}(E_R^{(1)}, E_R^{(2)}), \mathcal{O}_T).$$

However, since R is a discrete valuation ring, we have a direct sum decomposition of $\mathcal{O}_S$-modules

$$p_{S*}\mathcal{H}om_{X_S}(F,G) \simeq \mathcal{H}_0 \oplus \mathcal{H}_1$$

where $\mathcal{H}_1$ is free and $\mathcal{H}_0$ is a torsion module. Since η_K is non-zero, it follows that $\mathcal{H}_1$ is nontrivial. Therefore there exists a surjective morphism

$$\eta_R \in \mathrm{Hom}_S(p_{S*}\mathcal{H}om_{X_S}(F,G), \mathcal{O}_S)$$

so that $\eta_R|_T = \pi^n \eta_K$ for some $n \in \mathbb{Z}$. Let $\xi_R \in \mathrm{Hom}_{X_S}(F,G)$ be the morphism of $\mathcal{O}_{X_S}$ modules which corresponds to η_R. Then we have $\xi_R|_T = \pi^n \xi_K$ and the restriction of ξ_R to any fiber of $X_R \to S$ is nontrivial.

Next we prove that the stability condition for ADHM sheaves implies $n = 0$. Since ξ_K is an isomorphism of ADHM sheaves we have

$$\xi_K \circ \psi_K^{(1)} = \psi_K^{(2)},$$

hence also

$$\pi^n \xi_K \circ \psi_K^{(1)} = \pi^n \psi_K^{(2)}. \tag{3.9}$$

By assumption, $\psi_K^{(1)}, \psi_K^{(2)}$ extend to morphisms of $\mathcal{O}_{X_R}$-modules $\psi_R^{(1)} : (E_\infty)_R \to E_R^{(1)}$, $\psi_R^{(2)} : (E_\infty)_R \to E_R^{(2)}$. Moreover, since $\mathcal{E}_R^{(1)}$ is a flat family of stable ADHM sheaves, the restriction of $\psi_R^{(1)}$ to the closed fiber must be nontrivial. Therefore the extension $\xi_R \circ \psi_R^{(1)}$ of $\pi^n \xi_K \circ \psi_K^{(1)}$ to S has nontrivial restriction to the closed fiber. The extension $\psi_R^{(2)}$ of $\psi_K^{(2)}$ also has nontrivial restriction to the closed fiber since $\mathcal{E}_R^{(2)}$ is a flat family of stable ADHM sheaves. Then (3.9) implies $n = 0$.

In conclusion the isomorphism $\xi_K : E_K^{(1)} \to E_K^{(2)}$ extends to a morphism of $\mathcal{O}_{X_R}$-modules $\xi_R : E_R^{(1)} \to E_R^{(2)}$ with nontrivial restriction to the central fiber.

Recall that ξ_K is an morphism of ADHM sheaves on X_K, i.e. the diagrams (2.1) for $\mathcal{E}_K^{(1)}$, $\mathcal{E}_K^{(2)}$ and ξ_K are commutative. Moreover, $E_R^{(1)}, E_R^{(2)}$ are torsion free $\mathcal{O}_{X_R}$-modules since they are flat over S, and torsion free along each fiber. Then it follows that ξ_R is a morphism of ADHM sheaves on X_R since a section of a torsion free sheaf which vanishes over an open subset must vanish everywhere.

In particular the restriction $\xi_R|_{X_k} : \mathcal{E}_R^{(1)}|_{X_k} \to \mathcal{E}_R^{(2)}|_{X_k}$ of ξ_R to the closed fiber is an automorphism of stable ADHM sheaves. Since we have already established that $\xi_R|_{X_k}$ is nontrivial, it must be an isomorphism according to lemma (2.3.*i*.) Then lemma (3.10) implies that ξ_R is an isomorphism.

Proposition (3.9) implies that $\mathfrak{M}_{ADHM}(\mathcal{X}, P)$ is of finite type over $\mathbb{C}$ as an Artin stack. Since the category of algebraic spaces is a full subcategory of the category of Artin stacks, this further implies that $\mathfrak{M}_{ADHM}(\mathcal{X}, P)$ is of finite type as an algebraic space. Alternatively, this follows from the proof of lemma (3.11) below forgetting the equivariant structure.

□

3.3. **Torus Actions and Fixed Loci.** Since the moduli space of ADHM sheaves is not proper, an algebraic torus action with proper fixed loci will be required in order in order to obtain a well defined residual theory in the sense of [18]. Let us start with a general result.

Lemma 3.11. *Suppose we have an algebraic torus action* $\mathbf{T}\times\mathfrak{M}_{ADHM}(\mathcal{X}, P) \to \mathfrak{M}_{ADHM}(\mathcal{X}, P)$ *so that the natural smooth atlas* $R \to \mathfrak{M}_{ADHM}$ *provided by proposition (3.9) is* $\mathbf{T}$*-equivariant. Then the algebraic space* $\mathfrak{M}_{ADHM}(\mathcal{X}, P)$ *admits a* $\mathbf{T}$*-equivariant étale atlas* $\iota : S \to \mathfrak{M}_{ADHM}(\mathcal{X}, P)$ *where* S *is scheme of finite type over* $\mathbb{C}$ *equipped with an algebraic* $\mathbf{T}$*-action. In particular there exists a finite collection of* $\mathbf{T}$*-equivariant étale morphisms* $\{\iota_\alpha : S_\alpha \to \mathfrak{M}_{ADHM}(X, P)\}$, $\alpha \in \Lambda$, *with* S_α *affine of finite type over* $\mathbb{C}$ *which cover* $\mathfrak{M}_{ADHM}(\mathcal{X}, P)$.

Proof. Let V be any affine étale atlas of $\mathfrak{M}_{ADHM}(\mathcal{X}, P)$, where V is a scheme over $\mathbb{C}$. Then $S = R \times_{\mathfrak{M}_{ADHM}(\mathcal{X},P)} V$ is again an étale atlas of

$\mathfrak{M}_{ADHM}(\mathcal{X},P)$. Moreover S is of finite type over $\mathbb{C}$ since R is quasi-projective, and $S\to\mathfrak{M}_{ADHM}(\mathcal{X},P)$ is $\mathbf{T}$-equivariant since $R\to\mathfrak{M}_{ADHM}(\mathcal{X},P)$ is $\mathbf{T}$-equivariant.

□

Next let us provide two explicit examples of algebraic torus actions.

Example 3.12. *Let* $\mathbf{T}=\mathbb{C}^\times\times\mathbb{C}^\times$ *and define an action* $\mathbf{T}\times\mathfrak{M}_{ADHM}(\mathcal{X},P)\to\mathfrak{M}_{ADHM}(\mathcal{X},P)$ *by*

$$(E_S,\Phi_{S,1},\Phi_{S,2},\phi_S,\psi_S)\to(E_S,t_1\Phi_{S,1},t_2\Phi_{S,2},t_1t_2\phi_S,\psi_S) \tag{3.10}$$

for any $(t_1,t_2)\in\mathbf{T}(S)$ *and any flat family* $\mathcal{E}_S=(E_S,\Phi_{S,1,2},\phi_S,\psi_S)$ *of stable ADHM sheaves. Note that in this case the hypothesis of lemma (3.12) is satisfied i.e. there is a natural* $\mathbf{T}$*-action on* R *so that the universal family* $\mathcal{E}_R$ *is* $\mathbf{T}$*-equivariant.*

Example 3.13. *Suppose* $E_\infty=\oplus_{a=1}^{r_\infty}L_a$*, where* $r_\infty\in\mathbb{Z}$*,* $r_\infty\geq 1$*, and* L_a*,* $a=1,\ldots,r_\infty$*, are invertible* $\mathcal{O}_X$*-modules. Let* $\mathbf{T}=\mathbb{C}^\times\times\mathbb{C}^\times\times(\mathbb{C}^\times)^{r_\infty-1}$*. There is a natural torus action*

$$(\mathbb{C}^\times)^{r_\infty-1}\times E_\infty\to E_\infty$$

given by the identification of $(\mathbb{C}^\times)^{r_\infty-1}$ *with the hypersurface*

$$\prod_{a=1}^{r_\infty}z_a=1$$

in $(\mathbb{C}^\times)^{r_\infty}$*. This provides a* $(\mathbb{C}^\times)^{r_\infty-1}$ *action* $\xi_\infty:(\mathbb{C}^\times)^{r_\infty-1}\to\mathrm{Aut}(E_\infty)$ *given by*

$$\xi_\infty(\underline{z})|_{L_a}=z_a1_{L_a}$$

for any $a=1,\ldots,r_\infty$*, where* $\underline{z}=(z_a)_{a=1,\ldots,r_\infty}\in(\mathbb{C}^\times)^{r_\infty-1}$*. Using the above presentation of the torus* $(\mathbb{C}^\times)^{r_\infty-1}$*, its characters will be denoted in the following by* $\underline{m}=(m_1,\ldots,m_{r_\infty})\in\mathbb{Z}^{r_\infty}$*.*

If S *is a scheme of finite type over* $\mathbb{C}$*, we will denote by* $(\xi_\infty)_S:(\mathbb{C}^\times)^{r_\infty-1}(S)\to\mathrm{Aut}(E_S)$ *the relative version of this action. Then we have a natural torus action on* $\mathfrak{M}_{ADHM}(\mathcal{X},P)$ *given by*

$$(E_S,\Phi_{S,1},\Phi_{S,2},\phi_S,\psi_S)\to(E_S,t_1\Phi_{S,1},t_2\Phi_{S,2},t_1t_2(\xi_\infty)_S(\underline{z})\circ\phi_S,\psi_S\circ\xi_\infty(\underline{z})^{-1}) \tag{3.11}$$

for any $(t_1,t_2,\underline{z})\in\mathbf{T}(S)$ *and any flat family* $\mathcal{E}_S=(E_S,\Phi_{S,1,2},\phi_S,\psi_S)$ *of stable ADHM sheaves.*

Remark 3.14. *Note that the actions defined in examples (3.12), (3.13) coincide if* E_∞ *is a line bundle on* X*.*

The following proposition shows that the fixed loci of the action defined in (3.13) are proper if X is a curve. In this case the Hilbert polynomial P is determined by the rank $r\geq 1$ and the degree $e\in\mathbb{Z}$, and the moduli spaces will be denoted by $\mathfrak{M}_{ADHM}(\mathcal{X},r,e)$.

Proposition 3.15. *Let X be a smooth projective curve over $\mathbb{C}$, and suppose $E_\infty = \oplus_{a=1}^{r_\infty} L_a$ as in example (3.13). Then the fixed locus $\mathfrak{M}(\mathcal{X}, r, e)^{\mathbf{T}}$ of the torus action given in example (3.13) is a closed algebraic subspace of $\mathfrak{M}(\mathcal{X}, r, e)$ isomorphic to a projective scheme over $\mathbb{C}$.*

Proof. A standard argument [23, Sect 1.1] shows that $\mathfrak{M}(\mathcal{X}, r, e)^{\mathbf{T}}$ is a closed algebraic subspace of $\mathfrak{M}(\mathcal{X}, r, e)$ if $\mathfrak{M}(\mathcal{X}, r, e)$ is separated. Therefore it is separated and of finite type over $\mathbb{C}$.

We will rely on [63] for a systematic treatment of group actions on stacks. Let S be a scheme of finite type over $\mathbb{C}$. A fixed point in $\mathfrak{M}(\mathcal{X}, r, e)(S)$ is an isomorphism class of a flat family of stable ADHM sheaves $\mathcal{E}_S$ on X_S equipped with an isomorphism of ADHM sheaves

$$\xi_S(t_1, t_2, \underline{z}) : (E_S, \Phi_{S,1}, \Phi_{S,2}, \phi_S, \psi_S) \to (E_S, t_1\Phi_{S,1}, t_2\Phi_{S,2}, t_1t_2\phi_S, \psi_S)$$

for each $(t_1, t_2, \underline{z}) \in \mathbf{T}(S)$ satisfying a natural cocycle identity.

In particular, we have a group homomorphism $\xi_S : \mathbf{T}(S) \to \mathrm{Aut}(E_S)$ i.e. a $\mathbf{T}(S)$-action on E_S. Therefore, according to [21], E_S decomposes in a direct sum of $\mathcal{O}_{X_S}$-modules

$$E_S = \bigoplus_{(n_1,n_2)\in\mathbb{Z}^2} \bigoplus_{\underline{m}\in\mathbb{Z}^{r_\infty}} E_S(n_1, n_2, \underline{m}) \tag{3.12}$$

where only finitely many terms are nontrivial. An element $(t_1, t_2, \underline{z}) \in \mathbf{T}(S)$ acts by multiplication by $t_1^{n_1} t_2^{n_2} z_1^{m_1} \ldots z_{r_\infty}^{m_{r_\infty}}$ on $E_S(n_1, n_2, \underline{m})$. Moreover each direct summand $E_S(n_1, n_2, \underline{m})$ must be flat over S since E_S is flat over S.

By definition the restriction of $\mathcal{E}_S$ to any fiber X_s, $s \in S$, is a stable ADHM sheaf $\mathcal{E}_s$ on the smooth projective curve X_s over the residual field $k(s)$. Since E_S is flat over S, the restriction $\xi_S(t_1, t_2, \underline{z})|_{X_s}$ is an isomorphism of stable ADHM sheaves

$$\xi_s(t_1, t_2, \underline{z}) : (E_s, \Phi_{s,1}, \Phi_{s,2}, \phi_s, \psi_s) \to (E_s, t_1\Phi_{s,1}, t_2\Phi_{s,2}, t_1t_2\phi_s, \psi_s).$$

In particular we have the following commutative diagrams of $\mathcal{O}_{X_s}$-modules.
(3.13)

$$\begin{tikzcd}
E_s \otimes_{X_s} (M_i)_s \arrow[r, "\Phi_{s,i}"] \arrow[d, "\xi_s(t_1,t_2,\underline{z})\otimes 1_{(M_i)_s}"'] & E_s \arrow[d, "\xi_s(t_1,t_2,\underline{z})"] \\
E_s \otimes_{X_s} (M_i)_s \arrow[r, "t_i\Phi_{s,i}"] & E_s
\end{tikzcd}
\qquad
\begin{tikzcd}
(E_\infty)_s \arrow[r, "\psi_s"] \arrow[d, "(\xi_\infty)_s(\underline{z})^{-1}"'] & E_s \arrow[d, "\xi_s(t_1,t_2,\underline{z})"] \\
(E_\infty)_s \arrow[r, "\psi_s"] & E_s
\end{tikzcd}$$

$$\begin{tikzcd}
E_s \otimes_{X_s} (M_{12})_s \arrow[r, "\phi_s"] \arrow[d, "\xi_s(t_1,t_2,\underline{z})\otimes 1_{(M_{12})_s}"'] & (E_\infty)_s \arrow[d, "(\xi_\infty)_s(\underline{z})^{-1}"] \\
E_s \otimes_{X_s} (M_{12})_s \arrow[r, "t_1t_2\phi_s"] & (E_\infty)_s
\end{tikzcd}$$

for any $t = (t_1, t_2, \underline{z}) \in \mathbf{T}(S)$ and $i = 1, 2$.

The commutative diagrams (3.13) imply that the only nontrivial components of $\Phi_{s,1,2}, \phi_s, \psi_s$ with respect to the character decomposition (3.12) are

$$\begin{aligned} \Phi_{s,1}(n_1,n_2,\underline{m}) &: E_s(n_1,n_2,\underline{m})\otimes_{X_s}(M_1)_s \to E_s(n_1+1,n_2,\underline{m}) \\ \Phi_{s,2}(n_1,n_2,\underline{m}) &: E_s(n_1,n_2,\underline{m})\otimes_{X_s}(M_2)_s \to E_s(n_1,n_2+1,\underline{m}) \\ \phi_s(-1,-1,\underline{\delta}^a) &: E_s(-1,-1,\underline{\delta}^a)\otimes_{X_s}(M_{12})_s \to (L_a)_s \\ \psi_s(a) &: (L_a)_s \to E_s(0,0,\underline{\delta}^a) \end{aligned} \tag{3.14}$$

where

$$\underline{\delta}^a = (\underbrace{0,0,\ldots,0}_{a-1},-1,\underbrace{0,\ldots,0}_{r_\infty-a})$$

for $a=1,\ldots,r_\infty$. In particular note that

$$\mathrm{Im}(\psi_s)\subseteq\bigoplus_{a=1}^{r_\infty}E_s(0,0,\underline{\delta}^a).$$

Moreover, the following relations must also be satisfied

$$\begin{aligned} &\Phi_{s,2}(n_1+1,n_2,\underline{m})\circ(\Phi_{s,1}(n_1,n_2,\underline{m})\otimes 1_{(M_2)_s}) = \\ &\Phi_{s,1}(n_1,n_2+1,\underline{m})\circ(\Phi_{s,2}(n_1,n_2,\underline{m})\otimes 1_{(M_1)_s}) \end{aligned} \tag{3.15}$$

for any $(n_1,n_2)\in\mathbb{Z}^2_{\geq 0}$ and any $\underline{m}\in\mathbb{Z}^{r_\infty}$.

Then it follows from the first two equations in (3.14) that the canonical destabilizing subsheaf $(E_s)_0$ constructed in lemma (2.4) is a subsheaf of

$$\bigoplus_{n_1,n_2\geq 0}\bigoplus_{a=1}^{r_\infty}E_s(n_1,n_2,\underline{\delta}^a).$$

Given the direct sum decomposition (3.12), where the terms are torsion free sheaves, the stability condition implies that

$$E_s(n_1,n_2,\underline{m})=0$$

if $n_1<0$ or $n_2<0$ or $\underline{m}\neq\underline{\delta}^a$. Otherwise, the saturation $\overline{(E_s)_0}$ would be a proper subsheaf of E_s. Therefore we must have $\phi_s=0$ for any fixed ADHM sheaf.

Taking into account relations (3.15), same argument implies that the saturations

$$\overline{\mathrm{Im}(\Phi_{s,1}(n_1,n_2,\underline{\delta}^a))}\subseteq E_s(n_1+1,n_2,\underline{\delta}^a)$$
$$\overline{\mathrm{Im}(\Phi_{s,2}(n_1,n_2,\underline{\delta}^a))}\subseteq E_s(n_1,n_2+1,\underline{\delta}^a)$$

must be equal to $E_s(n_1+1,n_2,\underline{\delta}^a)$, respectively $E_s(n_1,n_2+1,\underline{\delta}^a)$ for any $(n_1,n_2)\in\mathbb{Z}^2_{\geq 0}$, and any $a=1,\ldots,r_\infty$. Moreover the saturation

$$\overline{\mathrm{Im}(\psi_s(a))}\subseteq E_s(0,0,\underline{\delta}^a)$$

must also be equal to $E_s(0,0,\underline{\delta}^a)$ for any $a=1,\ldots,r_\infty$. Since $(L_a)_s$, $a=1,\ldots,r_\infty$ are line bundles on X_s, and X_s is a smooth projective curve over s, this implies that the data $(E_s(n_1,n_2,\underline{\delta}^a), \Phi_{s,i}(n_1,n_2,a), \psi_s(a))$ satisfies the following conditions

- (a) All nontrivial direct summands $(E_s(n_1,n_2,\underline{\delta}^a)$ in (3.12) are invertible $\mathcal{O}_{X_s}$-modules.
- (b) The morphism of $\mathcal{O}_{X_s}$-modules $\psi_s(a)$ must be injective for all $a = 1,\ldots,r_\infty$.
- (c) The morphisms of $\mathcal{O}_{X_s}$-modules $\Phi_s(n_1,n_2,\underline{\delta}^a)$ must be injective for any $a = 1,\ldots,r_\infty$ and any $(n_1,n_2) \in \mathbb{Z}^2_{\geq 0}$ so that the target is nontrivial.
- (d) Relations (3.15) must be satisfied.

Note that the degree of each line bundle $E_s(n_1,n_2,\underline{\delta}^a)$ is constant along each connected component of S since $E_S(n_1,n_2,\underline{\delta}^a)$ are flat over S. For simplicity we will assume in the following that S is connected. The arguments below generalize obviously to the case when S has several connected components. Let $e(n_1,n_2,a) = \deg E_S(n_1,n_2,\underline{\delta}^a)$. For each $a = 1,\ldots,r_\infty$ let Δ_a the set of pairs $(n_1,n_2) \in \mathbb{Z}^\times$ so that the direct summand $E_S(n_1,n_2,\underline{\delta}^a)$ in (3.12) is nontrivial.

Let Z be the moduli space of isomorphism classes of data

$$(L(n_1,n_2,a), \Phi_{1,2}(n_1,n_2,a), \psi(a))$$

with $a = 1,\ldots,r_\infty$ and $(n_1,n_2) \in \Delta_a$ so that

- $L(n_1,n_2,a)$ is a line bundle on X of degree $e(n_1,n_2,a)$.
- $\Phi_1(n_1,n_2,a) : L(n_1,n_2,a) \otimes_X M_1 \to L(n_1+1,n_2,a)$, $\Phi_2(n_1,n_2,a) : L(n_1,n_2,a) \otimes_X M_2 \to L(n_1,n_2+1,a)$, $\psi(a) : L_a \to L(0,0,a)$ are injective morphisms of $\mathcal{O}_X$-modules for any $a = 1,\ldots,r_\infty$ and any $(n_1,n_2) \in \Delta_a$ whenever the target is nontrivial.
- $\Phi_2(n_1+1,n_2,a) \circ (\Phi_1(n_1,n_2,a) \otimes 1_{M_2}) = \Phi_1(n_1,n_2+1,a) \circ (\Phi_2(n_1,n_2,a) \otimes 1_{M_1})$ for any $a = 1,\ldots,r_\infty$ and any $(n_1,n_2) \in \Delta_a$.

Note that Z is a proper scheme over $\mathbb{C}$ since by construction it is a closed subscheme of a certain direct product of symmetric powers of X. Moreover Z is a fine moduli space i.e. there is a universal object on $Z \times X$. This follows from the fact that the symmetric product $S^d(X)$, $d \geq 1$ is a fine moduli space for degree d effective divisors on X.

Then we claim that $\mathfrak{M}_{ADHM}(\mathcal{X},r,e)^{\mathbf{T}}$ is isomorphic to the projective scheme Z. Clearly, the structure results for $\mathbf{T}$-fixed S-valued points of $\mathfrak{M}_{ADHM}(X,r,e)$ proven above imply that any such point $[\mathcal{E}_S]$ determines a morphism $S \to Z$ so that the data $(E_S(n_1,n_2,a), \Phi_{S,1,2}(n_1,n_2,a), \psi(a))$ is isomorphic to the pull back of the universal data on Z. Conversely, any morphism $S \to Z$ determines a set of data $L_S(n_1,n_2,a), \Phi_S(n_1,n_2,a), \psi_S(a)$, $a = 1,\ldots,r_\infty$, $(n_1,n_2) \in \Delta_a$ satisfying conditions analogous to $(a-c)$ above and relations (3.15), up to isomorphism.

Then the data

$$\begin{aligned} E_S &= \oplus_{a=1}^{r_\infty} \oplus_{(n_1,n_2)\in\Delta_a} L_S(n_1,n_2,a) \\ \Phi_{S,1,2} &= \oplus_{a=1}^{r_\infty} \oplus_{(n_1,n_2)\in\Delta_a} \Phi_{S,1,2} \\ \psi_S &= \oplus_{a=1}^{r_\infty} \psi_S(a). \end{aligned}$$

determine an ADHM sheaf $\mathcal{E}_S$ on X_S with $\phi_S = 0$. Note that the restriction $\mathcal{E}_S|_{X_s}$ to each fiber X_s is a stable ADHM sheaf since the saturation of the canonical destabilizing subsheaf $(E_s)_0$ is equal to E_s by construction. Moreover, one can easily check that $[\mathcal{E}_S]$ is a fixed point with respect to the **T** action defined in example (3.13).

□

4. Deformation Theory of ADHM Sheaves on Curves

In this section we provide some basics on deformation theory of ADHM sheaves on curves. Therefore in the following X will be a fixed smooth projective curve over $\mathbb{C}$ of genus g. Our main references for deformation theory are [6], [7], [48].

The following observation will be very useful throughout this section.

Lemma 4.1. *Suppose X is a smooth complex projective curve. Let $\mathcal{E}_S$ be a flat family of stable ADHM sheaves on X parameterized by a scheme S of finite type over $\mathbb{C}$. Then E_S is a locally free $\mathcal{O}_{X_S}$-module.*

Proof. Since $\mathcal{E}_S$ is a flat family of stable ADHM sheaves, it follows that $E_S|_{X_s}$ is a torsion free $\mathcal{O}_X$-module, for any point $s \in S$. Since X is a smooth projective curve, this implies that $E_S|_{X_s}$ is locally free. Then lemma (4.1) follows from [40, Lemma 2.1.7] since E_S is flat over S.

□

4.1. Deformation Complex.

Let S be a scheme of finite type over $\mathbb{C}$. In the following we will call a nilpotent thickening of S a scheme S' of finite type over $\mathbb{C}$ so that

- S' is a scheme over S
- there is closed embedding $S \subset S'$ of schemes over S so that the defining ideal sheaf I of S in S' is nilpotent.

Note that if these conditions are satisfied, the projection $\pi_{S'/S} : S' \to S$ is a finite morphism. Therefore it follows from the base change theorem that higher direct images vanish and $\pi_{S'/S*}$ is an exact functor from $\mathcal{O}_{S'}$-modules to $\mathcal{O}_S$-modules. Moreover, the induced projection $(\pi_{S'/S})_X : X_{S'} \to X_S$ satisfies the same properties. For simplicity we will not explicitely write $(\pi_{S'/S})_{X*}$ in the following. Given a $\mathcal{O}_{X_{S'}}$-module F we will denote $(\pi_{S'/S})_{X*}F$ also by F regarded as a $\mathcal{O}_{X_S}$-module. The same conventions will apply to morphisms of $\mathcal{O}_{X_{S'}}$-modules. In particular we have a canonical exact sequence of $\mathcal{O}_S$-modules

$$0 \to I \to \mathcal{O}_{S'} \to \mathcal{O}_S \to 0.$$

Since we have a closed embedding $S \to S'$ of schemes over S, it follows that there is a canonical splitting of $\mathcal{O}_S$-modules $\mathcal{O}_{S'} \simeq \mathcal{O}_S \oplus I$. In the following we will identify $\mathcal{O}_{S'} = \mathcal{O}_S \oplus I$ as $\mathcal{O}_S$-modules.

Given a coherent $\mathcal{O}_S$-module I, the trivial nilpotent thickening of S determined by I is the nilpotent thickening S' so that $\mathcal{O}_{S'} = \mathcal{O}_S \oplus I$ as $\mathcal{O}_S$-modules, the ring structure of $\mathcal{O}_{S'}$ being determined by

$$(s,x)\cdot(s',x') = (ss', sx + s'x') \tag{4.1}$$

for any local sections s, s' of $\mathcal{O}_S$ and x, x' of I. In particular $I^2 = 0$ with respect to the multiplicative structure of $\mathcal{O}_{S'}$.

Definition 4.2. (*i*) *Let $\mathcal{E}_S$ be a flat family of stable ADHM sheaves on X parameterized by a scheme S of finite type over $\mathbb{C}$ and let S' be a nilpotent thickening of S. An extension of $\mathcal{E}_S$ to S' is a flat family $\mathcal{E}_{S'}$ of stable ADHM sheaves on X so that $\mathcal{E}_{S'} \otimes_{S'} \mathcal{O}_S = \mathcal{E}_S$.*

(*ii*) *Two extensions $\mathcal{E}^{(1)}_{S'}$, $\mathcal{E}^{(2)}_{S'}$ of $\mathcal{E}_S$ to S' are equivalent if there exists an isomorphism $\xi_{S'} : \mathcal{E}^{(1)}_{S'} \to \mathcal{E}^{(2)}_{S'}$ of ADHM sheaves on $X_{S'}$ so that $\xi_{S'}|_{X_S} = 1_{\mathcal{E}_S}$.*

Definition 4.3. *Let $\mathcal{E}^{(1)}_S, \mathcal{E}^{(2)}_S$ be two flat families of ADHM sheaves on X parameterized by a scheme S of finite type over $\mathbb{C}$ and let I be a coherent $\mathcal{O}_S$-module. We define the complex $\mathcal{C}(\mathcal{E}^{(1)}_S, \mathcal{E}^{(2)}_S, I)$ to be the following complex of $\mathcal{O}_{X_S}$-modules*

$$0 \to \mathcal{H}om_{X_S}(E^{(1)}_S, E^{(2)}_S \otimes_{X_S} I) \xrightarrow{d_1} \begin{array}{c} \mathcal{H}om_{X_S}(E^{(1)}_S \otimes_{X_S} (M_1)_S, E^{(2)}_S \otimes_{X_S} I) \\ \oplus \\ \mathcal{H}om_{X_S}(E^{(1)}_S \otimes_{X_S} (M_2)_S, E^{(2)}_S \otimes_{X_S} I) \\ \oplus \\ \mathcal{H}om_{X_S}(E^{(1)}_S \otimes_{X_S} (M_{12})_S, (E_\infty)_S \otimes_{X_S} I) \\ \oplus \\ \mathcal{H}om_{X_S}((E_\infty)_S, E^{(2)}_S \otimes_{X_S} I) \end{array} \tag{4.2}$$

$$\xrightarrow{d_2} \mathcal{H}om_{X_S}(E^{(1)}_S \otimes_{X_S} (M_{12})_S, E^{(2)}_S \otimes_{X_S} I) \to 0$$

where

$$\begin{aligned} d_1(\alpha) = {}^t\big(& -\alpha\circ\Phi^{(1)}_{S,1} + (\Phi^{(2)}_{S,1}\otimes 1_I)\circ(\alpha\otimes 1_{(M_1)_S}), \\ & -\alpha\circ\Phi^{(1)}_{S,2} + (\Phi^{(2)}_{S,2}\otimes 1_I)\circ(\alpha\otimes 1_{(M_2)_S}), \\ & (\phi^{(2)}_S\otimes 1_I)\circ(\alpha\otimes 1_{(M_{12})_S}), -\alpha\circ\psi^{(1)}_S\big) \end{aligned}$$

for any local section α of $\mathcal{H}om_{X_S}(E^{(1)}_S, E^{(2)}_S \otimes_{X_S} I)$, and

$$\begin{aligned} d_2(\beta_1,\beta_2,\gamma,\delta) = & \beta_1\circ(\Phi^{(1)}_{S,2}\otimes 1_{(M_1)_S}) - (\Phi^{(2)}_{S,2}\otimes 1_I)\circ(\beta_1\otimes 1_{(M_2)_S}) \\ & - \beta_2\circ(\Phi^{(1)}_{S,1}\otimes 1_{(M_2)_S}) + (\Phi^{(2)}_{S,1}\otimes 1_I)\circ(\beta_2\otimes 1_{(M_1)_S}) \\ & + (\psi^{(2)}_S\otimes 1_I)\circ\gamma + \delta\circ\phi^{(1)}_S \end{aligned}$$

for any local sections $(\beta_1, \beta_2, \gamma, \delta)$ *of*

$$\mathcal{H}om_{X_S}(E_S^{(1)} \otimes_{X_S} (M_1)_S, E_S^{(2)} \otimes_{X_S} I) \oplus \mathcal{H}om_{X_S}(E_S^{(1)} \otimes_{X_S} (M_2)_S, E_S^{(2)} \otimes_{X_S} I)$$
$$\oplus\, \mathcal{H}om_{X_S}(E_S^{(1)} \otimes_{X_S} (M_{12})_S, (E_\infty)_S \otimes_{X_S} I) \oplus \mathcal{H}om_{X_S}((E_\infty)_S, E_S^{(2)} \otimes_{X_S} I),$$

The degrees of the terms of (4.2) *are* $0, 1, 2$ *respectively.*

In the following we will use the following notation conventions.

- *If* $I = \mathcal{O}_S$, *we will denote* $\mathcal{C}(\mathcal{E}_S^{(1)}, \mathcal{E}_S^{(2)}, I)$ *by* $\mathcal{C}(\mathcal{E}_S^{(1)}, \mathcal{E}_S^{(2)})$.
- *If* $\mathcal{E}_S^{(1)} = \mathcal{E}_S^{(2)} = \mathcal{E}_S$, *we will denote* $\mathcal{C}(\mathcal{E}_S^{(1)}, \mathcal{E}_S^{(2)}, I)$ *by* $\mathcal{C}(\mathcal{E}_S, I)$.
- *If* $\mathcal{E}_S^{(1)} = \mathcal{E}_S^{(2)} = \mathcal{E}_S$ *and* $I = \mathcal{O}_S$, *we will denote* $\mathcal{C}(\mathcal{E}_S^{(1)}, \mathcal{E}_S^{(2)}, I)$ *by* $\mathcal{C}(\mathcal{E}_S)$.

Remark 4.4. *Under the conditions of definition (4.3), suppose* S' *is a nilpotent thickening of* S *and* I *is the ideal sheaf of* S *in* S'. *Let* $\mathcal{E}_S$ *be a flat family of stable ADHM sheaves on* X *parameterized by* S, *and let* $\mathcal{E}_{S'}$ *be an extension of* $\mathcal{E}_S$ *to* S. *Then the ADHM sheaf* $\mathcal{E}_{S'}$ *has a natural structure of ADHM sheaf on* X_S *as explained above definition (4.2). Therefore we can take* $\mathcal{E}_S^{(1)} = \mathcal{E}_S$, $\mathcal{E}_S^{(2)} = \mathcal{E}_{S'}$ *in definition (4.3) obtaining a complex* $\mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S'}, I)$ *of* $\mathcal{O}_{X_S}$*-modules.*

Proposition 4.5. (*i*) *Let* $\mathcal{E}_S$ *be a flat family of stable ADHM sheaves on* X *parameterized by a separated scheme* S *of finite type over* $\mathbb{C}$, *let* S' *be a nilpotent thickening of* S *and let* $\mathcal{E}_{S'}$ *be an extension of* $\mathcal{E}_S$ *to* S'. *Then* $\mathcal{E}_{S'}$ *determines a hypercohomology class* $\mathfrak{e}(\mathcal{E}_S, \mathcal{E}_{S'}, I) \in \mathbb{H}^1(X_S, \mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S'}, I))$.

(*ii*) *Let* $\mathcal{E}_{S'}^{(1)}, \mathcal{E}_{S'}^{(2)}$ *be two extensions of* $\mathcal{E}_S$ *to* S'. *Let* $\xi_{S'} : \mathcal{E}_{S'}^{(1)} \to \mathcal{E}_{S'}^{(2)}$ *be a morphism of ADHM sheaves on* $X_{S'}$ *so that* $\xi_{S'}|_{X_S} = 1_{\mathcal{E}_S}$. *Then there are natural induced morphisms of hypercohomology groups*

$$\xi^k_{S'*} : \mathbb{H}^k(X_S, \mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S'}^{(1)}, I)) \to \mathbb{H}^k(X_S, \mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S'}^{(2)}, I)) \tag{4.3}$$

with $k \in \mathbb{Z}_{\geq 0}$, *so that*

$$\xi^1_{S'*}(\mathfrak{e}(\mathcal{E}_S, \mathcal{E}_{S'}^{(1)}, I)) = \mathfrak{e}(\mathcal{E}_S, \mathcal{E}_{S'}^{(2)}, I)$$

Proof. Suppose $\mathcal{E}_{S'} = (E_{S'}, \Phi_{S',1,2}, \phi_{S'}, \psi_{S'})$ is such an extension. Since $E_{S'}$ is a locally free $\mathcal{O}_{X_{S'}}$-module according to lemma (4.1), we have an exact sequence of $\mathcal{O}_{X_{S'}}$-modules

$$0 \to E_{S'} \otimes_{X_{S'}} I \to E_{S'} \to E_S \to 0$$

which yields an exact sequence of $\mathcal{O}_{X_S}$-modules

$$0 \to E_{S'} \otimes_{X_S} I \xrightarrow{\iota_S} E_{S'} \to E_S \to 0. \tag{4.4}$$

Since $\mathcal{O}_{S'} = \mathcal{O}_S \oplus I$, as $\mathcal{O}_S$-modules, we also have a canonical identification

$$(E_\infty)_{S'} = (E_\infty)_S \oplus (E_\infty)_{S'} \otimes_{X_S} I$$

as $\mathcal{O}_{X_S}$-modules. By construction the following morphisms of $\mathcal{O}_{X_S}$-modules are obviously trivial
(4.5)

$$\begin{tikzcd}
E_{S'} \otimes_{X_S} (M_i)_S \otimes_{X_S} I \arrow[r, hook] & E_{S'} \otimes_{X_S} (M_i)_S \arrow[r, "\Phi_{S',i}"] & E_{S'} \arrow[r, two heads] & E_S
\end{tikzcd}$$

$$\begin{tikzcd}
E_{S'} \otimes_{X_S} (M_{12})_S \otimes_{X_S} I \arrow[r, hook] & E_{S'} \otimes_{X_S} (M_{12})_S \arrow[r, "\phi_{S'}"] & (E_\infty)_{S'} \arrow[r, two heads] & (E_\infty)_S
\end{tikzcd}$$

$$\begin{tikzcd}
(E_\infty)_{S'} \otimes_{X_S} I \arrow[r, hook] & (E_\infty)_{S'} \arrow[r, "\psi_{S'}"] & E_{S'} \arrow[r, two heads] & E_S
\end{tikzcd}$$

where $i = 1, 2$.

The extension of $\mathcal{O}_{X_S}$-modules (4.4) determines an extension class in $\mathrm{Ext}^1_{X_S}(E_S, E_{S'} \otimes_{X_S} I)$. Since E_S is locally free according to lemma (4.1), we have

$$\mathrm{Ext}^1_{X_S}(E_S, E_{S'} \otimes_{X_S} I) \simeq H^1(X_S, \mathcal{H}om_{X_S}(E_S, E_{S'} \otimes_{X_S} I)).$$

Since S is separated and of finite type over $\mathbb{C}$, there exists an affine Čech cover $\mathcal{U}_S = \{U_{S,\alpha}\}$ of X_S. Given an open embedding, $U_{S,\alpha_1\ldots\alpha_q} \subset U_{S,\alpha_1\ldots\alpha_p}$ with $p < q$, we will denote the restriction of any $\mathcal{O}_{U_{S,\alpha_1\ldots\alpha_p}}$-module or morphism of $\mathcal{O}_{U_{S,\alpha_1\ldots\alpha_p}}$-modules to $U_{S,\alpha_1\ldots\alpha_q}$ by a subscript $(\alpha_1 \ldots \alpha_q)$.

By restriction to each affine open subset $U_{S,\alpha} \subset X_S$, the morphisms of $\mathcal{O}_{X_S}$-modules $\Phi_{S,i}, \phi_S, \psi_S$ yield Čech 0-cocycles $\{(\Phi_{S,i})_\alpha\}, \{(\phi_S)_\alpha\}, \{(\psi_S)_\alpha\}$.

Next note that the restriction of the extension (4.4) to any affine open subset $U_{S,\alpha}$ is trivial since E_S is a locally free $\mathcal{O}_{X_S}$-module. Therefore we can choose local splittings

$$\begin{tikzcd}
0 \arrow[r] & (E_{S'} \otimes_{X_S} I)_\alpha \arrow[r] & (E_{S'})_\alpha \arrow[r, bend left] & (E_S)_\alpha \arrow[l, bend left, "\eta_\alpha"] \arrow[r] & 0
\end{tikzcd} \tag{4.6}$$

where $\eta = \{\eta_\alpha\} \in C^0(\mathcal{U}_S, \mathcal{H}om_{X_S}(E_S, E_{S'}))$. Then

$$e = \{e_{\alpha\beta}\} = \{(\delta\eta)_{\alpha\beta}\} \in C^1(\mathcal{U}_S, \mathcal{H}om_{X_S}(E_S, E_{S'} \otimes_{X_S} I))$$

is a 1-cocycle representing the extension class determined by (4.4).

The local splittings $\{\eta_\alpha\}$ determine local isomorphisms

$$\eta_\alpha \oplus (\iota_S)_\alpha : E_S \oplus (E_{S'} \otimes_{X_S} I)_\alpha \xrightarrow{\sim} (E_{S'})_\alpha$$

where $\iota_S : E_{S'} \otimes_{X_S} I \to E_{S'}$ is the canonical injection of $\mathcal{O}_{X_S}$ modules in (4.4). Moreover, we have commutative diagrams of $\mathcal{O}_{U_{S,\alpha}}$ modules of the form

(4.7)

$$\begin{tikzcd}
(E_S \oplus E_{S'} \otimes_{X_S} I)_\alpha \otimes_{U_{S,\alpha}} ((M_i)_S)_\alpha \arrow[r, "(\eta_\alpha \oplus (\iota_S)_\alpha) \otimes 1_{((M_i)_S)_\alpha}"] \arrow[d, "(\Lambda_\alpha)_i"'] & (E_{S'})_\alpha \otimes_{U_{S,\alpha}} ((M_i)_S)_\alpha \arrow[d, "(\Phi_{S',i})_\alpha"] \\
(E_S \oplus E_{S'} \otimes_{X_S} I)_\alpha \arrow[r, "\eta_\alpha \oplus (\iota_S)_\alpha"] & (E_{S'})_\alpha
\end{tikzcd}$$

$$\begin{tikzcd}
(E_S \oplus E_{S'} \otimes_{X_S} I)_\alpha \otimes_{U_{S,\alpha}} ((M_{12})_S)_\alpha \arrow[r, "(\eta_\alpha \oplus (\iota_S)_\alpha) \otimes 1_{((M_{12})_S)_\alpha}"] \arrow[d, "\lambda_\alpha"'] & (E_{S'})_\alpha \otimes_{U_{S,\alpha}} ((M_{12})_S)_\alpha \arrow[d, "(\phi_{S'})_\alpha"] \\
((E_\infty)_S)_\alpha \oplus ((E_\infty)_{S'} \otimes_{X_S} I)_\alpha \arrow[r, "1_{((E_\infty)_S)_\alpha} \oplus 1_{((E_\infty)_{S'} \otimes_{X_S} I)_\alpha}"] & ((E_\infty)_S)_\alpha \oplus ((E_\infty)_{S'} \otimes_{X_S} I)_\alpha
\end{tikzcd}$$

$$\begin{tikzcd}
((E_\infty)_S)_\alpha \oplus ((E_\infty)_{S'} \otimes_{X_S} I)_\alpha \arrow[r, "1_{((E_\infty)_S)_\alpha} \oplus 1_{((E_\infty)_{S'} \otimes_{X_S} I)_\alpha}"] \arrow[d, "\rho_\alpha"'] & ((E_\infty)_S)_\alpha \oplus ((E_\infty)_{S'} \otimes_{X_S} I)_\alpha \arrow[d, "(\psi_{S'})_\alpha"] \\
(E_S \oplus E_{S'} \otimes_{X_S} I)_\alpha \arrow[r, "\eta_\alpha \oplus (\iota_S)_\alpha"] & (E_{S'})_\alpha
\end{tikzcd}$$

for $i = 1, 2$. Since the morphisms (4.5) are trivial, $\{\Lambda_\alpha\}, \{\lambda_\alpha\}, \{\rho_\alpha\}$ are 0-cochains of the form

(4.8)

$$\begin{aligned}
(\Lambda_\alpha)_i &= \begin{pmatrix} (\Phi_{S,i})_\alpha & 0 \\ (\widetilde{\Phi}_\alpha)_i & (\Phi_{S',i})_\alpha \otimes 1_{I_\alpha} \end{pmatrix} \in C^0(\mathcal{U}_S, \mathcal{H}om_{X_S}((E_S \oplus E_{S'} \otimes_{X_S} I) \otimes_{X_S} (M_i)_S, \\
&\qquad\qquad\qquad E_S \oplus E_{S'} \otimes_{X_S} I)) \\
\lambda_\alpha &= \begin{pmatrix} (\phi_S)_\alpha & 0 \\ \widetilde{\phi}_\alpha & (\phi_{S'})_\alpha \otimes 1_{I_\alpha} \end{pmatrix} \in C^0(\mathcal{U}_S, \mathcal{H}om_{X_S}((E_S \oplus E_{S'} \otimes_{X_S} I) \otimes_{X_S} (M_{12}))_S, \\
&\qquad\qquad\qquad (E_\infty)_S \oplus (E_\infty)_{S'} \otimes_{X_S} I)) \\
\rho_\alpha &= \begin{pmatrix} (\psi_S)_\alpha & 0 \\ \widetilde{\psi}_\alpha & (\psi_{S'})_\alpha \otimes 1_{I_\alpha} \end{pmatrix} \in C^0(\mathcal{U}_S, \mathcal{H}om_{X_S}((E_\infty)_S \oplus (E_\infty)_{S'} \otimes_{X_S} I, \\
&\qquad\qquad\qquad E_S \oplus E_{S'} \otimes_{X_S} I))
\end{aligned}$$

with $i = 1, 2$, satisfying the conditions listed below. First, we have gluing conditions

$$\begin{aligned}
(g_S)_{\alpha\beta}((\Lambda_\beta)_i)_{\alpha\beta} &= ((\Lambda_\alpha)_i)_{\alpha\beta}(g_{S,i})_{\alpha\beta} \\
((g_\infty)_S)_{\alpha\beta}(\lambda_\beta)_{\alpha\beta} &= (\lambda_\alpha)_{\alpha\beta}(g_{S,12})_{\alpha\beta} \\
(g_S)_{\alpha\beta}(\rho_\beta)_{\alpha\beta} &= (\rho_\alpha)_{\alpha\beta}((g_\infty)_S)_{\alpha\beta}
\end{aligned} \tag{4.9}$$

where

$$
\begin{aligned}
(g_S)_{\alpha\beta} &= \begin{pmatrix} 1_{(E_S)_{\alpha\beta}} & 0 \\ (e_S)_{\alpha\beta} & 1_{(E_{S'}\otimes_{X_S} I)_{\alpha\beta}} \end{pmatrix} \\
(g_{S,i})_{\alpha\beta} &= \begin{pmatrix} 1_{(E_S)_{\alpha\beta}} \otimes 1_{((M_i)_S)_{\alpha\beta}} & 0 \\ (e_S)_{\alpha\beta} \otimes 1_{((M_i)_S)_{\alpha\beta}} & 1_{(E_{S'}\otimes_{X_S} I)_{\alpha\beta}} \otimes 1_{((M_i)_S)_{\alpha\beta}} \end{pmatrix} \\
(g_{S,12})_{\alpha\beta} &= \begin{pmatrix} 1_{(E_S)_{\alpha\beta}} \otimes 1_{((M_{12})_S)_{\alpha\beta}} & 0 \\ (e_S)_{\alpha\beta} \otimes 1_{((M_{12})_S)_{\alpha\beta}} & 1_{(E_{S'}\otimes_{X_S} I)_{\alpha\beta}} \otimes 1_{((M_{12})_S)_{\alpha\beta}} \end{pmatrix} \\
((g_\infty)_S)_{\alpha\beta} &= \begin{pmatrix} 1_{((E_\infty)_S)_{\alpha\beta}} & 0 \\ 0 & 1_{((E_\infty)_{S'}\otimes_{X_S} I)_{\alpha\beta}} \end{pmatrix}
\end{aligned}
$$

for $i = 1, 2$. In addition, the ADHM relation

$$\Phi_{S',1} \circ (\Phi_{S',2} \otimes 1_{(M_1)_S}) - \Phi_{S',2} \circ (\Phi_{S',1} \otimes 1_{(M_2)_S}) + \psi_{S'} \circ \phi_{S'} = 0$$

is equivalent to the following condition

$$
\begin{aligned}
&\widetilde{\Phi}_1(\Phi_{S,2} \otimes 1_{(M_1)_S}) - (\Phi_{S',2} \otimes 1_I)(\widetilde{\Phi}_1 \otimes 1_{(M_2)_S}) \\
&- \widetilde{\Phi}_2(\Phi_{S,1} \otimes 1_{(M_2)_S}) + (\Phi_{S',1} \otimes 1_I)(\widetilde{\Phi}_2 \otimes 1_{(M_1)_S}) \\
&+ (\psi_{S'} \otimes 1_I)\widetilde{\phi} + \widetilde{\psi}\phi_S = 0
\end{aligned} \tag{4.10}
$$

A simple computation shows that the conditions (4.9) are equivalent to

$$
\begin{aligned}
\delta(\widetilde{\Phi}_{S,i}) &= -e\Phi_{S,i} + (\Phi_{S',i} \otimes 1_I)(e \otimes 1_{(M_i)_S}) \\
\delta(\widetilde{\phi}_S) &= (\phi_{S'} \otimes 1_I)(e \otimes 1_{(M_{12})_S}) \\
\delta(\widetilde{\psi}_S) &= -e\psi_S
\end{aligned} \tag{4.11}
$$

where $i = 1, 2$.

Note that equations (4.10), (4.10) imply that the collection of cochains $\left(e, \widetilde{\Phi}_1, \widetilde{\Phi}_2, \widetilde{\phi}, \widetilde{\psi}\right)$ defines a 1-cocycle in the total hypercohomology complex associated to the complex of $\mathcal{O}_S$-modules (4.2), i.e. we have

$$\delta(e) = 0 \qquad d_1(e) - \delta(\widetilde{\Phi}_{1,2}, \widetilde{\phi}, \widetilde{\psi}) = 0 \qquad d_2(\widetilde{\Phi}_{1,2}, \widetilde{\phi}, \widetilde{\psi}) = 0.$$

Next suppose we make a different choice of local splittings $\{\eta'_\alpha\}$ in (4.6). Then note that the difference $\{\gamma_\alpha = \eta'_\alpha - \eta_\alpha\}$ is a 0-cochain

$$\{\gamma_\alpha\} \in C^0(\mathcal{U}_S, \mathcal{H}om_{X_S}(E_S, E_{S'} \otimes_{X_S} I)).$$

An elementary computation shows that the cocycle e and the cochains $(\widetilde{\Phi}_i, \widetilde{\phi}, \widetilde{\psi})$ change as follows

$$
\begin{aligned}
e' - e &= \delta(\gamma) \\
\widetilde{\Phi}'_i - \widetilde{\Phi}_i, &= -\gamma\Phi_{S,i} + (\Phi_{S',i} \otimes 1_I)(\gamma \otimes 1_{(M_i)_S}) \\
\widetilde{\phi}' - \widetilde{\phi} &= (\phi_{S'} \otimes 1_I)(\gamma \otimes 1_{(M_{12})_S}) \\
\widetilde{\psi}' - \widetilde{\psi} &= -\gamma\psi_S
\end{aligned}
$$

for $i = 1, 2$. Therefore it follows that the hypercohomology 1-cocycle $\left(e, \widetilde{\Phi}_1, \widetilde{\Phi}_2, \widetilde{\phi}, \widetilde{\psi}\right)$ changes by a coboundary in the total hypercohomology complex associated to (4.2), that is

$$\begin{gathered} e' - e = \delta(\gamma) \\ \left(\widetilde{\Phi}_{1,2}, \widetilde{\phi}, \widetilde{\psi}\right)' - \left(\widetilde{\Phi}_{1,2}, \widetilde{\phi}, \widetilde{\psi}\right) = d_1(\gamma). \end{gathered}$$

This proves proposition $(4.5.i)$. The required hypercohomology class is

$$\mathfrak{e}(\mathcal{E}_S, \mathcal{E}_{S'}, I) = \left[\left(e, \widetilde{\Phi}_1, \widetilde{\Phi}_2, \widetilde{\phi}, \widetilde{\psi}\right)\right].$$

In order to prove $(4.5.ii)$ let $\mathcal{E}^{(1)}_{S'}$, $\mathcal{E}^{(2)}_{S'}$ be extensions of $\mathcal{E}_S$ to S', and let $\xi_{S'} : \mathcal{E}^{(1)} \to \mathcal{E}^{(2)}$ be a morphism of ADHM sheaves on X so that $\xi_{S'}|_S = 1_{\mathcal{E}_S}$. Since $\xi_{S'}$ is a morphism of ADHM sheaves, it induces a natural morphism of complexes of $\mathcal{O}_{X_S}$-modules

$$\xi_{S'*} : \mathcal{C}(\mathcal{E}_S, \mathcal{E}^{(1)}_{S'}, I) \to \mathcal{C}(\mathcal{E}_S, \mathcal{E}^{(2)}_{S'}, I)$$

which induces in turn the morphisms (4.3).

It remains to check that $\xi^1_{S'*}$ maps the class $e(\mathcal{E}_S, \mathcal{E}^{(1)}_{S'}, I)$ to $e(\mathcal{E}_S, \mathcal{E}^{(2)}_{S'}, I)$. Let $\{\eta^{(1)}_\alpha\} \in C^0(\mathcal{U}_S, \mathcal{H}om_{X_S}(E_S, E^{(1)}_{S'}))$ be local splittings as in (4.6) for the extension (4.4) with $E_{S'} = E^{(1)}_{S'}$. Since $\xi_{S'}|_S = 1_{\mathcal{E}_S}$, the 0-cochain $\{\eta^{(2)}_\alpha\} \in C^0(\mathcal{U}_S, \mathcal{H}om_{X_S}(E_S, E^{(2)}_{S'}))$,

$$\eta^{(2)}_\alpha = (\xi_{S'})_\alpha \circ \eta^{(1)}_\alpha$$

provides analogous local splittings for the extension (4.4) with $E_{S'} = E^{(2)}_{S'}$. In particular this implies that

$$e^{(2)} = \xi_{S'} \circ e^{(1)}$$

Moreover we have commutative diagrams of $\mathcal{O}_{U_{S,\alpha}}$ modules

(4.12)
$$\begin{tikzcd}
(E_S \oplus E^{(1)}_{S'} \otimes_{X_S} I)_\alpha \arrow[r, "\eta^{(1)}_\alpha \oplus \iota^{(1)}_\alpha"] \arrow[d, "1_{(E_S)_\alpha} \oplus ((\xi_{S'})_\alpha \otimes 1_{I_\alpha})"'] & E^{(1)}_{S'} \arrow[d, "(\xi_{S'})_\alpha"] \\
(E_S \oplus E^{(2)}_{S'} \otimes_{X_S} I)_\alpha \arrow[r, "\eta^{(2)}_\alpha \oplus \iota^{(1)}_\alpha"] & E^{(2)}_{S'}
\end{tikzcd}$$

in which the horizontal arrows are isomorphisms. Then, using the commutative diagrams (4.7), (4.12) and the fact that $\xi_{S'}$ is a morphism of ADHM sheaves, a routine computation yields the following relations

$$\begin{aligned} ((\xi_{S'})_\alpha \otimes 1_{I_\alpha}) \circ (\widetilde{\Phi}^{(1)}_\alpha)_i &= (\widetilde{\Phi}^{(2)}_\alpha)_i \\ \widetilde{\phi}^{(1)}_\alpha &= \widetilde{\phi}^{(2)}_\alpha \\ ((\xi_{S'})_\alpha \otimes 1_{I_\alpha}) \circ \widetilde{\psi}^{(1)}_\alpha &= \widetilde{\psi}^{(2)}_\alpha. \end{aligned}$$

Obviously, these relations hold for any choice of local splittings $\{\eta_\alpha^{(1)}\}$. Since $\xi_{S'}$ is a global morphism of ADHM sheaves, this implies that indeed $\xi^1_{S'*}(\mathfrak{e}(\mathcal{E}_S, \mathcal{E}_{S'}^{(1)}, I)) = \mathfrak{e}(\mathcal{E}_S, \mathcal{E}_S^{(2)}, I)$.

□

Remark 4.6. *Under the hypotheses of proposition (4.5.i), suppose S' is the trivial nilpotent thickening of S determined by a coherent $\mathcal{O}_S$-module I. Then we have a canonical isomorphism $\mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S'}, I) \simeq \mathcal{C}(\mathcal{E}_S, I)$ since $I^2 = 0$ in the ring structure of $\mathcal{O}_{S'}$, therefore $\mathcal{E}_{S'} \otimes_{X_S} I \simeq \mathcal{E}_S \otimes_{X_S} I$. In particular, in this case the complex $\mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S'}, I)$ is independent of $\mathcal{E}_{S'}$.*

Corollary 4.7. *Let $\mathcal{E}_S$ be a flat family of stable ADHM sheaves on X parameterized by an affine scheme S of finite type over $\mathbb{C}$, and let S' be the trivial nilpotent thickening of S determined by a coherent $\mathcal{O}_S$-module I. Then there is a one-to-one correspondence between equivalence classes of extensions $\mathcal{E}_{S'}$ of $\mathcal{E}_S$ to S' and hypercohomology classes in $\mathbb{H}^1(X_S, \mathcal{C}(\mathcal{E}_S, I))$.*

Proof. Proposition (4.5.i) implies that any extension $\mathcal{E}_{S'}$ determines a hypercohomology class $\mathfrak{e}(\mathcal{E}_S, \mathcal{E}_{S'}, I) \in \mathbb{H}^1(X_S, \mathcal{C}(\mathcal{E}_S, I))$. Moreover, (4.5.$ii$) implies that equivalent extensions determine identical hypercohomology classes.

Conversely, let $\mathfrak{e} \in \mathbb{H}^1(X_S, \mathcal{C}(\mathcal{E}_S, I))$ and let

$$(e, \widetilde{\Phi}_1, \widetilde{\Phi}_2, \widetilde{\phi}, \widetilde{\psi}) \in \begin{array}{c} C^1(\mathcal{U}_S, \mathcal{H}om_{X_S}(E_S, E_S \otimes_{X_S} I)) \\ \oplus \\ C^0(\mathcal{U}_S, \mathcal{H}om_{X_S}((M_1)_S \otimes_{X_S} E_S, E_S \otimes_{X_S} I)) \\ \oplus \\ C^0(\mathcal{U}_S, \mathcal{H}om_{X_S}((M_2)_S \otimes_{X_S} E_S, E_S \otimes_{X_S} I)) \\ \oplus \\ C^0(\mathcal{U}_S, \mathcal{H}om_{X_S}((M_{12})_S \otimes_{X_S} E_S, (E_\infty)_S \otimes_{X_S} I)) \\ \oplus \\ C^0(\mathcal{U}_S, \mathcal{H}om_{X_S}((E_\infty)_S, E_S \otimes_{X_S} I)) \end{array}$$

be a hypercohomology 1-cocycle representing $\mathfrak{e}$. In particular $\delta(e) = 0$, therefore e determines an extension class

$$[e] \in H^1(X_S, \mathcal{H}om_{X_S}(E_S, E_S \otimes_{X_S} I)) \simeq \mathrm{Ext}^1_{X_S}(E_S, E_S \otimes_{X_S} I). \tag{4.13}$$

This determines in turn an equivalence class of extensions of $\mathcal{O}_{X_S}$-modules of the form

$$0 \to E_S \otimes_{X_S} I \xrightarrow{\iota_S} E'_S \xrightarrow{j_S} E_S \to 0.$$

Let us consider such an extension with the property that there exist local splittings $\{\eta_\alpha\} \in C^0(\mathcal{U}_S, \mathcal{H}om_{X_S}(E_S, E_{S'}))$ as in (4.6) so that

$$\delta(\eta) = e.$$

The data $\Big(\{(\widetilde{\Phi}_\alpha)_i\}, \{\widetilde{\phi}_\alpha\}, \{\widetilde{\psi}_\alpha\}\Big)$ satisfying the relations (4.11), (4.10) determines 0-cochains $(\{\Lambda_\alpha\}_i, \{\lambda_\alpha\}, \{\rho_\alpha\})$ as in (4.8) satisfying relations (4.9).

Then using the commutative diagrams of the form (4.7), in which the horizontal arrows are isomorphisms, we obtain local morphisms

$$\begin{aligned}\Phi'_{S,i,\alpha} &: (E'_S)_\alpha \otimes_{U_{S,\alpha}} ((M_i)_S)_\alpha \to (E_{S'})_\alpha\\ \phi'_{S,\alpha} &: (E'_S)_\alpha \otimes_{U_{S,\alpha}} ((M_{12})_S)_\alpha \to ((E_\infty)_{S'})_\alpha\\ \psi'_{S,\alpha} &: ((E_\infty)_{S'})_\alpha \to (E'_S)_\alpha\end{aligned}$$

for $i = 1, 2$. Moreover, relations (4.9) imply that $\Big(\{\Phi'_{S,i,\alpha}\}, \{\phi'_{S,\alpha}\}, \{\psi'_{S,\alpha}\}\Big)$, $i = 1, 2$ are in fact 0-cocycles. Therefore they define global morphisms of $\mathcal{O}_{X_S}$-modules

$$\begin{aligned}\Phi'_{S,i} &: E'_S \otimes_{X_S} (M_i)_S \to E'_S\\ \phi'_S &: E'_S \otimes_{X_S} (M_{12})_S \to (E_\infty)_{S'}\\ \psi'_S &: (E_\infty)_{S'} \to E'_S.\end{aligned} \tag{4.14}$$

which satisfy the ADHM relation, according to relations (4.10). Moreover, by construction the morphisms of $\mathcal{O}_{X_S}$-modules ι_S, j_S in (4.13) yield naturally morphisms of ADHM sheaves on X_S.

Next let us prove that E'_S has a $\mathcal{O}_{X_{S'}}$-module structure, and the morphisms (4.14) are morphisms of $\mathcal{O}_{X_{S'}}$-modules. Since E'_S is an $\mathcal{O}_{X_S}$ by construction, and we have a canonical splitting $\mathcal{O}_{S'} \simeq \mathcal{O}_S \oplus I$, it suffices to define multiplication by local sections of I. According to [69, Lemma 3.4], E'_S this is given by

$$x \cdot u = \iota_S(j_S(u) \otimes x)$$

for any local sections x, u of I, E'_S respectively. Therefore E'_S has indeed a structure of $\mathcal{O}_{X_{S'}}$-module.

We also have to prove that the morphisms $\Phi'_{S,i}, \phi'_S, \psi'_S$ are morphisms of $\mathcal{O}_{X_{S'}}$-modules. Let us check for example that $\Phi'_{S,i}$ are morphisms of $\mathcal{O}_{X_{S'}}$ modules, for $i = 1, 2$. The proof for ϕ'_S, ψ'_S is the identical. Since ι_S, j_S are morphisms of ADHM sheaves on X_S, we have

$$\begin{aligned}\Phi'_{S,i}(x \cdot u) &= \Phi'_{S,i}(\iota_S(j_S(u) \otimes x)) = \iota_S((\Phi_{S,i} \otimes 1_I)(j_S(u) \otimes x))\\ &= \iota_S(\Phi_{S,i}(j_S(u)) \otimes x) = \iota_S(j_S(\Phi'_{S,i}(u)) \otimes x)\\ &= x \cdot \Phi'_{S,i}(u).\end{aligned} \tag{4.15}$$

Therefore $\Phi'_{S,i}$, $i = 1, 2$ are indeed morphisms of $\mathcal{O}_{X_{S'}}$-modules.

In order to finish the proof note that E'_S is flat over S' according to [69, Lemma 3.7]. Moreover, a routine computation shows that different hypercohomology cocycles representing $\mathfrak{e}$ yield equivalent extensions.

□

Next let us determine the obstructions in the deformation theory of stable ADHM sheaves. Recall [6], [48] that a standard deformation situation consists of a sequence of closed embeddings of schemes over S

$$S \subset S' \subset S'' \tag{4.16}$$

where S is a scheme of finite type over $\mathbb{C}$, and S', S'' are nilpotent thickenings of S. We will assume in the following that S is also separated over $\mathbb{C}$.

Let $I_{S\subset S'}, I_{S\subset S''}, I_{S'\subset S''}$ be the ideal sheaves corresponding to the closed embeddings $S \subset S'$, $S \subset S''$ and $S' \subset S''$ respectively. We will assume that that $I_{S\subset S''} \cdot I_{S'\subset S''} = 0$ in $\mathcal{O}_{S''}$, hence in particular $I^2_{S'\subset S''} = 0$. Therefore $I_{S'\subset S''}$ has a $\mathcal{O}_S$-module structure, and we have an exact sequence of $\mathcal{O}_S$-modules

$$0 \to I_{S'\subset S''} \to I_{S\subset S''} \to I_{S\subset S'} \to 0. \tag{4.17}$$

Suppose we have a flat family $\mathcal{E}_{S'}$ of stable ADHM sheaves on X parameterized by S', which restricts to a given family $\mathcal{E}_S$ over S. Recall that E_S, $E_{S'}$ are locally free $\mathcal{O}_{X_S}$, respectively $\mathcal{O}_{X_{S'}}$-modules. Moreover, since the projection morphism $S' \to S$ is finite, the base change theorem implies that $E_{S'}$ is also a locally free $\mathcal{O}_{X_S}$-module. Then the exact sequence of $\mathcal{O}_S$-modules (4.17) yields an exact sequence of complexes of $\mathcal{O}_{X_S}$-modules

$$0 \to \mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S'\subset S''}) \to \mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S''}) \to \mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'}) \to 0. \tag{4.18}$$

Note that we have a canonical isomorphism

$$\mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S'\subset S''}) \simeq \mathcal{C}(\mathcal{E}_S, \mathcal{E}_S, I_{S'\subset S''})$$

of $\mathcal{O}_S$-modules since $I^2_{S'\subset S''} = 0$. Therefore (4.18) yields a long exact sequence of hypercohomology groups which reads in part

$$\begin{aligned} \cdots \to \mathbb{H}^1(X_S, \mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S''})) &\to \mathbb{H}^1(X_S, \mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'})) \\ \xrightarrow{\partial} \mathbb{H}^2(X_S, \mathcal{C}(\mathcal{E}_S, \mathcal{E}_S, I_{S'\subset S''})) &\to \cdots \end{aligned} \tag{4.19}$$

According to proposition (4.5.i), the data $(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'})$ determines a hypercohomology class $\mathfrak{e}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'}) \in \mathbb{H}^1(X_S, \mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'})$.

Definition 4.8. *Given a deformation situation* (4.16)*, a flat family* $\mathcal{E}_S$ *and an extension* $\mathcal{E}_{S'}$ *to* S'*, we define the obstruction class*

$$\mathfrak{o}b(\mathcal{E}_{S'}, S', S'') \in \mathbb{H}^2(X_S, \mathcal{C}(\mathcal{E}_S, \mathcal{E}_S, I_{S'\subset S''}))$$

to be

$$\mathfrak{o}b(\mathcal{E}_{S'}, S', S'') = \partial(\mathfrak{e}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'})).$$

Proposition 4.9. $\mathcal{E}_{S'}$ *can be extended to a flat family of stable ADHM sheaves on* X *parameterized by* S'' *if and only if* $\mathfrak{o}b(\mathcal{E}_{S'}, S', S'') = 0$.

Proof. For the direct implication, suppose that the family $\mathcal{E}_{S'}$ extends to a flat family $\mathcal{E}_{S''}$ of stable ADHM sheaves on X parameterized by S'' so that $\mathcal{E}_{S''} \otimes_{X_{S''}} \mathcal{O}_{X_{S'}} = \mathcal{E}_{S'}$. Then, according to proposition (4.5.i), we have hypercohomology classes

$$\mathfrak{e}(\mathcal{E}_S, \mathcal{E}_{S''}, I_{S\subset S''}) \subset \mathbb{H}^1(X_S, \mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S''}, I_{S\subset S''}))$$
$$\mathfrak{e}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'}) \subset \mathbb{H}^1(X_S, \mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'}))$$

and it is straightforward to check that $\mathfrak{e}(\mathcal{E}_S, \mathcal{E}_{S''}, I_{S\subset S''})$ is a lift of $\mathfrak{e}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'})$ in the exact sequence (4.19). Therefore $\mathfrak{o}b(\mathcal{E}_S, S', S'') = 0$.

Conversely, suppose $\mathcal{E}_{S'}$ is an extension of $\mathcal{E}_S$ to S' so that

$$\partial(\mathfrak{e}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'})) = 0.$$

Then there exists a lift $\mathfrak{e}''$ of $\mathfrak{e}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'})$ in the exact sequence (4.19).

Let $\{\eta'_\alpha\} \in C^0(\mathcal{U}_S, \mathcal{H}om X_S(E_S, E_{S'}))$ be local splittings for the extension of $\mathcal{O}_S$-modules

$$0 \to E_{S'} \otimes_{X_S} I_{S\subset S'} \xrightarrow{\iota'_S} E_{S'} \to E_S \to 0$$

and let $(e', \widetilde{\Phi}'_i, \widetilde{\phi}', \widetilde{\psi}')$ be the corresponding hypercohomology representative of $\mathfrak{e}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'})$ constructed in the proof of proposition (4.5.i). Let $\{(e'', \widetilde{\Phi}''_i, \widetilde{\phi}'', \widetilde{\psi}'')\}$ be a hypercohomology representative of $\mathfrak{e}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'})$ which maps to $(e', \widetilde{\Phi}'_i, \widetilde{\phi}', \widetilde{\psi}')$ under the natural morphism induced by the projection $I_{S\subset S"} \to I_{S\subset S'}$.

The 1-cocycle $e'' \in C^1(\mathcal{U}_S, \mathcal{H}om X_S(E_S, E_{S'} \otimes_{X_S} I_{S\subset S''}))$ determines an equivalence class of extensions of $\mathcal{O}_{X_S}$-modules of the form

$$0 \to E_{S'} \otimes_{X_S} I_{S\subset S''} \xrightarrow{\iota''_S} E''_S \to E_S \to 0.$$

Let E''_S be such an extension equipped with local splittings $\{\eta''_\alpha\} \in C^0(\mathcal{U}_S, \mathcal{H}om_{X_S}(E_S, E''_S))$ so that $\delta(\eta'') = e''$. Then, proceeding by analogy with the proof of corollary (4.7), one constructs morphisms of $\mathcal{O}_{X_S}$-modules

$$\begin{aligned} \Phi''_{S,i} &: E''_S \otimes_{X_S} (M_i)_S \to E''_S \\ \phi''_S &: E''_S \otimes_{X_S} (M_{12})_S \to (E_\infty)_{S''} \\ \psi''_S &: (E_\infty)_{S''} \to E''_S. \end{aligned}$$

satisfying the ADHM relation.

Next let us prove that the data $(E''_S, \Phi''_{S,i}, \phi''_S, \psi''_S)$ is an ADHM sheaf on $X_{S''}$ which extends $\mathcal{E}_{S'}$. Note that on each affine open subset $U_{S,\alpha}$ we have a diagram of $\mathcal{O}_{U_{S,\alpha}}$-modules

$$\begin{tikzcd} (E_S)_\alpha \oplus (E_{S'})_\alpha \otimes_{U_{S,\alpha}} (I_{S\subset S''})_\alpha \arrow[r, "\eta''_\alpha \oplus (\iota''_S)_\alpha"] \arrow[d, two heads] & (E''_S)_\alpha \\ (E_S)_\alpha \oplus (E_{S'})_\alpha \otimes_{U_{S,\alpha}} (I_{S\subset S'})_\alpha \arrow[r, "\eta'_\alpha \oplus (\iota''_S)_\alpha"] & (E_{S'})_\alpha \end{tikzcd} \tag{4.20}$$

in which the horizontal rows are isomorphisms. Therefore there exist local surjective morphisms $j_\alpha : (E''_S)_\alpha \to (E_{S'})_\alpha$ which complete (4.20) to a commutative diagram of $\mathcal{O}_{U_{S,\alpha}}$-modules. Furthermore, using the fact that $e_{S''}$ is a lift of $e_{S'}$ under the natural morphisms induced by the projection $I_{S\subset S''} \twoheadrightarrow I_{S\subset S'}$, a simple computation shows that the morphisms $\{j_\alpha\}$

agree on overlaps. Therefore we obtain global surjective morphism of $\mathcal{O}_{X_S}$-modules $j_S : E''_S \twoheadrightarrow E_{S'}$ which fits in a commutative diagram of $\mathcal{O}_{X_S}$-modules

$$\begin{tikzcd}
0 \arrow[r] & E'_S \otimes_{X_S} I_{S\subset S''} \arrow[r, "\iota''_S"] \arrow[d, two heads] & E''_S \arrow[r] \arrow[d, two heads, "j_S"] & E_S \arrow[r] \arrow[d, two heads, "1_{E_S}"] & 0 \\
0 \arrow[r] & E_{S'} \otimes_{X_S} I_{S\subset S'} \arrow[r, "\iota'_S"] & E_{S'} \arrow[r] & E_S \arrow[r] & 0
\end{tikzcd}$$

in which the horizontal rows are exact and the vertical arrows are surjective.

Since E''_S is a $\mathcal{O}_{X_S}$-module by construction, and we have a canonical splitting $\mathcal{O}_{S''} = \mathcal{O}_S \oplus I_{S\subset S''}$, it suffices to define multiplication by local sections of $I_{S\subset S''}$. According to [69, Prop. 3.13] this is given by

$$x'' \cdot u = \iota''_S((j_S \otimes 1_{I_{S\subset S''}})(u \otimes x''))$$

for any local sections x'', u of $I_{S\subset S''}$, E''_S respectively. Then a computation identical to (4.15) shows that $\Phi''_{S,i}, \phi''_S, \psi''_S$ are morphisms of $\mathcal{O}_{S''}$-modules.

In order to finish the proof, note that $E_{S''}$ is flat over S'' according to lemma [69, Lemma 3.7].

□

Next we prove a vanishing result needed later in the proof of virtual smoothness.

Lemma 4.10. *Suppose X is a smooth projective curve over $\mathbb{C}$, and let $\mathcal{E}$ be a stable ADHM sheaf on X. Then $\mathbb{H}^i(X, \mathcal{C}(\mathcal{E})) = 0$, for all $i \geq 3$ and for all $i \leq 0$.*

Proof. Let us first prove vanishing for $i \geq 3$. $\mathcal{C}(\mathcal{E})$ is obtained by setting $S = \mathrm{S}pec(\mathbb{C})$ and $I = \mathcal{O}_S$ in definition (4.3). Since the degrees of the three terms in (4.2) are $0, 1, 2$ respectively, it follows that all terms $E_1^{p,q}$, $p+q \geq 4$ in the standard hypercohomology spectral sequence are trivially zero. Therefore $\mathbb{H}^i(X, \mathcal{C}(\mathcal{E})) = 0$ for $i \geq 4$. Moreover, the only nonzero term on the diagonal $p+q = 3$ is

$$E_1^{2,1} = H^1(X, \mathcal{H}om_X(E \otimes_X M_{12}, E)).$$

The differential

$$d_1^{1,1} : E_1^{1,1} \to E_1^{2,1}$$

is the map

$$\begin{array}{c}
H^1(X, \mathcal{H}om_X(E \otimes_X M_1, E)) \\
\oplus \\
H^1(X, \mathcal{H}om_X(E \otimes_X M_2, E)) \\
\oplus \\
H^1(X, \mathcal{H}om_X(E \otimes_X M_{12}, E_\infty)) \\
\oplus \\
H^1(X, \mathcal{H}om_X(E_\infty, E))
\end{array}
\xrightarrow{d_1^{1,1}} H^1(X, \mathcal{H}om_X(E \otimes_X M_{12}, E))$$

induced by d_2.

We claim that this map is surjective if $\mathcal{E}$ is stable. In order to prove this claim we will prove that the dual map $(d_1^{1,1})^\vee : (E_1^{2,1})^\vee \to (E_1^{1,1})^\vee$ is injective if $\mathcal{E}$ is stable. Using Serre duality, the dual differential is a linear map

$$\mathrm{Hom}_X(E, E\otimes M_{12}\otimes_X K_X)) \xrightarrow{(d_1^{1,1})^\vee} \begin{array}{c} \mathrm{Hom}_X(E, E\otimes_X M_1\otimes_X K_X)) \\ \oplus \\ \mathrm{Hom}_X(E, E\otimes_X M_2\otimes_X K_X)) \\ \oplus \\ \mathrm{Hom}_X(E_\infty, E\otimes_X M_{12}\otimes_X K_X) \\ \oplus \\ \mathrm{Hom}_X(E, E_\infty\otimes_X K_X)) \end{array}$$

which maps a global homomorphism $\alpha \in \mathrm{Hom}_X(E, E\otimes_X M_{12}\otimes_X K_X))$ to

$$\begin{array}{c} -(\alpha\otimes 1_{M_2^{-1}})\circ\Phi_2 + (\Phi_2\otimes 1_{M_1}\otimes 1_{K_X})\circ\alpha \\ (\alpha\otimes 1_{M_1^{-1}})\circ\Phi_1 - (\Phi_1\otimes 1_{M_2}\otimes 1_{K_X})\circ\alpha \\ \alpha\circ\psi \\ (\phi\otimes 1_{K_X})\circ\alpha \end{array}$$

If $\alpha \in \mathrm{Ker}((d_1^{1,1})^\vee)$, it follows that $\mathrm{Ker}(\alpha)$ is Φ-invariant and $\mathrm{Im}(\psi) \subseteq \mathrm{Ker}(\alpha)$. Moreover, since α is a morphism of locally free sheaves, $\mathrm{Ker}(\alpha)$ must be saturated subsheaf of E. Then the stability of $\mathcal{E}$ implies that $\mathrm{Ker}(\alpha) = E$, hence $\alpha = 0$. This proves the claim.

The second assertion is trivial for $i < 0$. The case $i = 0$ follows again from the stability of $\mathcal{E}$ by an identical argument. □

4.2. **ADHM Sheaves with Trivial Framing.**

We conclude this section with some specific results for ADHM sheaves with $E_\infty = \mathcal{O}_X$. Under the conditions of definition (4.3) note that we have a morphism of $\mathcal{O}_{X_S}$-modules

$$\mathcal{H}om_{X_S}((E_\infty)_S, (E_\infty)_S\otimes_{X_S} I) \xrightarrow{\kappa} \begin{array}{c} \mathcal{H}om_{X_S}(E_S^{(1)}\otimes_{X_S}(M_{12})_S, (E_\infty)_S\otimes_{X_S} I) \\ \oplus \\ \mathcal{H}om_{X_S}((E_\infty)_S, E_S^{(2)}\otimes_{X_S} I) \end{array}$$

given by

$$\alpha_\infty \to {}^t(-\alpha_\infty\circ\phi_S^{(1)}, (\psi_S^{(2)}\otimes 1_I)\circ\alpha_\infty)$$

for any local section α_∞ of $\mathcal{H}om_{X_S}((E_\infty)_S, (E_\infty)_S\otimes_{X_S} I)$. Moreover, it is straightforward to check that $d_2\circ\kappa = 0$, therefore κ yields a morphism of complexes

$$\mathcal{H}om_{X_S}((E_\infty)_S, (E_\infty)_S\otimes X_S I)[-1] \xrightarrow{\kappa[-1]} \mathcal{C}(\mathcal{E}_S^{(1)}, \mathcal{E}_S^{(2)}, I). \tag{4.21}$$

Let

$$\widetilde{\mathcal{C}}(\mathcal{E}_S^{(1)}, \mathcal{E}_S^{(2)}, I) = \mathsf{Cone}(\kappa[-1]). \tag{4.22}$$

Using notation conventions analogous to definition (4.3), we will denote $\widetilde{\mathcal{C}}(\mathcal{E}_S, \mathcal{E}_S, I)$ by $\widetilde{\mathcal{C}}(\mathcal{E}_S, I)$ and $\widetilde{\mathcal{C}}(\mathcal{E}_S^{(1)}, \mathcal{E}_S^{(2)}, \mathcal{O}_S)$ by $\widetilde{\mathcal{C}}(\mathcal{E}_S^{(1)}, \mathcal{E}_S^{(2)})$. If $I = \mathcal{O}_S$, $\widetilde{\mathcal{C}}(\mathcal{E}_S, I)$ will be denoted by $\widetilde{\mathcal{C}}(\mathcal{E}_S)$.

Note that there is a canonical 0-th hypercohomology class $1_{\mathcal{E}_S} \in \mathbb{H}^0(X_S, \widetilde{\mathcal{C}}(\mathcal{E}_S))$ determined by the pair

$$\left(1_{E_S}, 1_{(E_\infty)_S}\right) \in H^0(X_S, \mathcal{H}om_{X_S}(E_S, E_S)) \oplus H^0(X_S, \mathcal{H}om_{X_S}((E_\infty)_S, (E_\infty)_S))$$

It is straightforward to check that

$$\widetilde{d}_1 \left(1_{E_S}, 1_{(E_\infty)_S}\right) = 0$$

therefore $1_{\mathcal{E}_S}$ is indeed a well defined 0-th hypercohomology class.

By construction we also have an exact sequence of complexes of $\mathcal{O}_{X_S}$-modules

(4.23)
$$0 \to \mathcal{C}(\mathcal{E}_S^{(1)}, \mathcal{E}_S^{(2)}, I) \to \widetilde{\mathcal{C}}(\mathcal{E}_S^{(1)}, \mathcal{E}_S^{(2)}, I) \to \mathcal{H}om_{X_S}((E_\infty)_S, (E_\infty)_S \otimes_{X_S} I) \to 0.$$

Lemma 4.11. *Under the conditions of definition (4.3), suppose* $\mathcal{E}_S^{(1)} = \mathcal{E}_S^{(2)} = \mathcal{E}_S$ *and* $E_\infty = \mathcal{O}_X$*. Then the exact sequence* (4.23) *has a canonical splitting, and we have isomorphisms of hypercohomology groups*

$$\mathbb{H}^k(X_S, \widetilde{\mathcal{C}}(\mathcal{E}_S, \mathcal{E}_S, I)) \simeq \mathbb{H}^k(X_S, \mathcal{C}(\mathcal{E}_S, \mathcal{E}_S, I)) \oplus H^k(X_S, p_S^* I) \tag{4.24}$$

for all $k \in \mathbb{Z}$*.*

Proof. Since $(E_\infty)_S = \mathcal{O}_{X_S}$, $\mathcal{H}om_{X_S}((E_\infty)_S, (E_\infty)_S \otimes_{X_S} I) \simeq p_S^* I$, and we have a morphism

$$\begin{aligned} p_S^* I \to \ & \mathcal{H}om_{X_S}(E_S, E_S \otimes_{X_S} I) \oplus p_S^* I \\ \alpha_\infty \to \ & (1_{E_S} \otimes \alpha_\infty, \alpha_\infty) \end{aligned}$$

It is straightforward to check that this is a splitting of (4.23).

□

Under the conditions of proposition (4.5), we have an exact sequence of complexes of $\mathcal{O}_{X_S}$-modules

$$0 \to \widetilde{\mathcal{C}}(\mathcal{E}_S, \mathcal{E}_{S'}, I) \to \widetilde{\mathcal{C}}(\mathcal{E}_S, \mathcal{E}_{S'}, \mathcal{O}_{X_S}) \to \widetilde{\mathcal{C}}(\mathcal{E}_S) \to 0. \tag{4.25}$$

Let

$$\widetilde{\mathfrak{e}}(\mathcal{E}_S, \mathcal{E}_{S'}, I) = \widetilde{\partial}^1(1_{\mathcal{E}_S}) \tag{4.26}$$

where

$$\widetilde{\partial}^1 : \mathbb{H}^0(X_S, \widetilde{\mathcal{C}}(\mathcal{E}_S)) \to \mathbb{H}^1(X_S, \widetilde{\mathcal{C}}(\mathcal{E}_S, \mathcal{E}_{S'}, I))$$

is the connecting isomorphism determined by (4.25). We will also denote by

$$\widetilde{i}^1 : \mathbb{H}^1(X_S, \mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S'}, I)) \to \mathbb{H}^1(X_S, \widetilde{\mathcal{C}}(\mathcal{E}_S, \mathcal{E}_{S'}, I))$$

the natural morphism of hypercohomology groups determined by the canonical injective morphism of complexes $\mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S'}, I) \hookrightarrow \widetilde{\mathcal{C}}(\mathcal{E}_S, \mathcal{E}_{S'}, I)$ and by

$$\widetilde{i}^2 : \mathbb{H}^2(X_S, \mathcal{C}(\mathcal{E}_S, \mathcal{E}_S, I)) \hookrightarrow \mathbb{H}^2(X_S, \widetilde{\mathcal{C}}(\mathcal{E}_S, \mathcal{E}_S, I))$$

the injection determined by the splitting (4.24).

Now consider a deformation situation of the form (4.16), and let $\mathcal{E}_{S'}$ be a flat family of stable ADHM sheaves parameterized by S' which restricts to a given family $\mathcal{E}_S$ on X_S. Then we have an exact sequence of complexes of $\mathcal{O}_{X_S}$-modules

$$0 \to \widetilde{\mathcal{C}}(\mathcal{E}_S, I_{S'\subset S''}) \to \widetilde{\mathcal{C}}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S''}) \to \widetilde{\mathcal{C}}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'}) \to 0. \tag{4.27}$$

Lemma 4.12. *Still assuming* $E_\infty = \mathcal{O}_X$, *we have*

$$\widetilde{i}^2(\mathfrak{ob}(\mathcal{E}_{S'}, S', S'')) = (\widetilde{i}^1 \circ \widetilde{\partial}^2)(\widetilde{\mathfrak{e}}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'})) \tag{4.28}$$

where

$$\widetilde{\partial}^2 : \mathbb{H}^1(X_S, \widetilde{\mathcal{C}}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'})) \to \mathbb{H}^2(X_S, \widetilde{\mathcal{C}}(\mathcal{E}_S, \mathcal{E}_S, I_{S'\subset S''}))$$

is the connecting morphism determined by (4.27).

Proof. By construction we have a morphism of complexes of $\mathcal{O}_{X_S}$-modules

$$\begin{tikzcd}
0 \arrow[r] & \mathcal{C}(\mathcal{E}_S, I_{S'\subset S''}) \arrow[r] \arrow[d] & \mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S''}) \arrow[r] \arrow[d] & \mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'}) \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & \widetilde{\mathcal{C}}(\mathcal{E}_S, I_{S'\subset S''}) \arrow[r] & \widetilde{\mathcal{C}}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S''}) \arrow[r] & \widetilde{\mathcal{C}}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'}) \arrow[r] & 0
\end{tikzcd}$$

in which the horizontal arrows are exact. This yields a commutative diagram of morphisms of hypercohomology groups

$$\begin{tikzcd}
\mathbb{H}^1(X_S, \mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'})) \arrow[r, "\partial^2"] \arrow[d, "\widetilde{i}^1"'] & \mathbb{H}^2(X_S, \mathcal{C}(\mathcal{E}_S, \mathcal{E}_S, I_{S'\subset S''})) \arrow[d, "\widetilde{i}^2"] \\
\mathbb{H}^1(X_S, \widetilde{\mathcal{C}}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'})) \arrow[r, "\widetilde{\partial}^2"] & \mathbb{H}^2(X_S, \widetilde{\mathcal{C}}(\mathcal{E}_S, \mathcal{E}_S, I_{S'\subset S''}))
\end{tikzcd}$$

Therefore lemma (4.12) follows if we prove that

$$\widetilde{i}^1(\mathfrak{e}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'})) = \widetilde{\mathfrak{e}}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'})$$

However, this follows easily form the detailed construction of the class $\mathfrak{e}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'})$ given in the proof of proposition (4.5). Taking into account the fact that

$$(E_\infty)_{S'} = (E_\infty)_S \oplus (E_\infty)_{S'} \otimes_{X_S} I_{S\subset S'}$$

as $\mathcal{O}_{X_S}$-modules. The construction of the class

$$\widetilde{\mathfrak{e}}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'})) = \widetilde{\partial}^1(1_{\mathcal{E}_S})$$

is formally identical to the construction of the class $\mathfrak{e}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'})$ by definition of the coboundary morphism.

□

5. Virtual Smoothness for ADHM Sheaves on Curves

From this point on we take X to be a smooth projective curve of genus g over $\mathbb{C}$. Then the Hilbert polynomial of any torsion-free $\mathcal{O}_X$-module is determined by a pair of integers (r, e) with $r \geq 1$, namely the rank and degree respectively. In the following we will denote the moduli stack $\mathfrak{M}_{ADHM}(\mathcal{X}, P)$ by $\mathfrak{M}_{ADHM}(\mathcal{X}, r, e)$.

In this section our goal is to prove that the moduli stacks $\mathfrak{M}_{ADHM}(\mathcal{X}, r, e)$ are virtually smooth, i.e. they carry natural perfect obstruction theories [48, 10]. Similar results have been previously obtained for moduli spaces of (decorated) sheaves in [69, 57, 52, 61]. Our treatment is closer to [69], relying on the construction of [48], or, more precisely, the generalization presented in [47, Sect 2.2]. Moreover we will carry out our construction in the equivariant setting as in [31] with respect to an algebraic torus action on $\mathfrak{M}_{ADHM}(\mathcal{X}, r, e)$ satisfying the hypothesis of lemma (3.11.) The results of this section are valid for any such action, in particular for those presented in examples (3.12), (3.13).

5.1. **The Tangent-Obstruction Complex.**

The following lemmas establish the existence of a tangent-obstruction theory for any flat family of stable ADHM sheaves on X parameterized by an affine scheme S of finite type over $\mathbb{C}$. We will proceed by a step-by-step verification of the conditions formulated in [48, Sect 1] for such families.

Lemma 5.1. *Let $\mathcal{E}_S$ be a flat family of stable ADHM sheaves on X parameterized by an affine scheme S of finite type over $\mathbb{C}$. Let I be a coherent $\mathcal{O}_S$-module. Then the following hold.*

(i) In the notation of remark (4.6), we have

$$\mathbb{H}^1(X_S, \mathcal{C}(\mathcal{E}_S, I)) \simeq \Gamma_S(\mathbf{R}^1 p_{S*}\mathcal{C}(\mathcal{E}_S, I)). \tag{5.1}$$

(ii) Given a base change morphism $f : T \to S$ with T affine of finite type over $\mathbb{C}$, a coherent $\mathcal{O}_T$-module J and a morphism of $\mathcal{O}_T$-modules $\xi : f^*I \to J$, we have a canonical morphism

$$c(f, \xi) : f^*\mathbf{R}^1 p_{S*}\mathcal{C}(\mathcal{E}_S, I) \to \mathbf{R}^1 p_{T*}\mathcal{C}(\mathcal{E}_T, J) \tag{5.2}$$

of $\mathcal{O}_T$-modules, where $\mathcal{E}_T = f_X^*\mathcal{E}_S$.

(iii) Given a sequence of morphisms $U \xrightarrow{g} T \xrightarrow{f} S$, with U, T affine of finite type over $\mathbb{C}$, a coherent $\mathcal{O}_U$-module K, a coherent $\mathcal{O}_T$-module J, a coherent $\mathcal{O}_S$-module I, and morphisms $\eta : g^*J \to K$, $\xi : f^*I \to J$, there is a commutative diagram of canonical morphisms of the form

$$\begin{tikzcd} g^* f^* \mathbf{R} p_{S*}\mathcal{C}(\mathcal{E}_S, I) \arrow[r, "\simeq"] \arrow[d, "g^*c(f,\xi)"'] & (f \circ g)^* \mathbf{R}^1 p_{S*}\mathcal{C}(\mathcal{E}_S, I) \arrow[d, "c(f\circ g, \zeta)"] \\ g^*\mathbf{R}^1 p_{T*}\mathcal{C}(\mathcal{E}_T, J) \arrow[r, "c(g,\eta)"] & \mathbf{R}^1 p_{U*}\mathcal{C}(\mathcal{E}_U, K) \end{tikzcd} \tag{5.3}$$

where $\mathcal{E}_T = f_X^*\mathcal{E}_S$, $\mathcal{E}_U = g_X^*\mathcal{E}_T$ *and* $\zeta : (f\circ g)^*I \to K$ *is defined by the composition*

$$(f\circ g)^*I \simeq g^*f^*I \xrightarrow{g^*\xi} g^*J \xrightarrow{\eta} K.$$

(iv) Suppose there is a $\mathbf{T}$*-action on* S *so that the family* $\mathcal{E}_S$ *is* $\mathbf{T}$*-equivariant. Then* $\mathbf{R}^1 p_{S*}(\mathcal{E}_S, \mathcal{E}_S\otimes_{X_S} I)$ *has a natural* $\mathbf{T}$*-equivariant structure for any* $\mathbf{T}$*-equivariant* $\mathcal{O}_S$*-module* I*. Moreover, if all base change morphisms as well as the sheaves* I, J, K *are* $\mathbf{T}$*-equivariant the canonical morphism* $c(f,\xi)$ *is* $\mathbf{T}$*-equivariant, and the diagram* (5.3) *is* $\mathbf{T}$*-equivariant.*

Proof. Since S is affine, we have

$$\mathbb{H}^k(X_S, \mathcal{C}(\mathcal{E}, I)) \simeq \Gamma_S(\mathbf{R}^k p_S * \mathcal{C}(\mathcal{E}, I)) \tag{5.4}$$

for all $k\in\mathbb{Z}$, hence (5.1) follows.

Given a morphism $f : T\to S$ as in (5.1.*ii*), we have a canonical isomorphism

$$f_X^*\mathcal{C}(\mathcal{E}_S, I) \simeq \mathcal{C}(\mathcal{E}_T, f^*I)$$

where $\mathcal{E}_T = f_X^*\mathcal{E}_S$. From definition (4.3) it follows that a morphism $\xi : f^*I \to J$ induces a canonical morphism

$$\mathcal{C}(\mathcal{E}_T, f^*I) \to \mathcal{C}(\mathcal{E}_T, J).$$

Therefore we obtain by composition a canonical morphism

$$f_X^*\mathcal{C}(\mathcal{E}_S, I) \to \mathcal{C}(\mathcal{E}_T, J).$$

Applying the functorial properties of direct images (A.1), we obtain a canonical morphisms of $\mathcal{O}_T$-modules

$$f^*\mathbf{R}^1 p_{S*}\mathcal{C}(\mathcal{E}_S, I) \to \mathbf{R}^1 p_{T*}\mathcal{C}(\mathcal{E}_T, f^*I) \to \mathbf{R}^1 p_{T*}\mathcal{C}(\mathcal{E}_T, J)$$

whose composition is the required morphism (5.2).

Lemma (5.1.*iii*) follows from a routine diagram chasing argument using lemma (A.1.*ii*) and (A.1.*iii*). Although straightforward, the details are tedious and will be omitted.

According for example to [12], since the torus $\mathbf{T}$ has trivial group cohomology, all above manipulations are valid in the equivariant setting, employing equivariant derived functors of equivariant complexes of sheaves. Therefore the proof of (5.1.*iv*) is formally identical to the above.

□

Lemma 5.2. *Let* $\mathcal{E}_S$ *be a flat family of stable ADHM sheaves on* X *parameterized by an affine scheme* S *of finite type over* $\mathbb{C}$*. Let* $S\subset S'\subset S''$ *be a deformation situation as in* (4.16)*. Then the following hold.*

(i) In the notation of remark (4.6), we have

$$\mathbb{H}^2(X_S, \mathcal{C}(\mathcal{E}_S, I_{S'\subset S''})) \simeq \Gamma_S(\mathbf{R}^2 p_{S*}\mathcal{C}(\mathcal{E}_S)\otimes_{\mathcal{O}_S} I_{S'\subset S''}). \tag{5.5}$$

(ii) Given a base change morphism $f : T\to S$ *with* T *affine of finite type over* $\mathbb{C}$ *we have a canonical isomorphism*

$$c(f) : f^*\mathbf{R}^2 p_{S*}\mathcal{C}(\mathcal{E}_S) \xrightarrow{\sim} \mathbf{R}^2 p_{T*}\mathcal{C}(\mathcal{E}_T) \tag{5.6}$$

compatible with base change, where $\mathcal{E}_T = f_X^*\mathcal{E}_S$.

(iii) *Suppose* $S \subset S' \subset S''$ *is a deformation situation of the form* (4.16) *with* S, S', S'' *affine of finite type over* $\mathbb{C}$. *Suppose moreover we have a base change diagram*

$$\begin{tikzcd} T'' \arrow[r, "f''"] \arrow[d] & S'' \arrow[d] \\ T \arrow[r, "f"] & S \end{tikzcd}$$

where T *is an affine scheme of finite type over* $\mathbb{C}$ *and* $T'' \to T$ *has a section* $\sigma_T : T \to T''$ *so that the diagram*

$$\begin{tikzcd} T'' \arrow[r, "f''"] & S'' \\ T \arrow[r, "f"] \arrow[u, "\sigma_T"] & S \arrow[u, "\sigma_S"'] \end{tikzcd}$$

is commutative. Let $T' = S' \times_{S''} T''$, $f' : T' \to S'$ *be the natural projection, and* $\mathcal{E}_T = f_X^*\mathcal{E}_S$. *Suppose furthermore we have an extension* $\mathcal{E}_{S'}$ *of* $\mathcal{E}_S$ *to* S', *and let* $\mathcal{E}_{T'} = (f'_X)^*\mathcal{E}_{S'}$. *Then there is a canonical morphism of* $\mathcal{O}_T$*-modules*

$$(5.7)\quad c(f,S',S'') : f^*\mathbf{R}^2 p_{S*}(\mathcal{C}(\mathcal{E}_S)\otimes_S I_{S'\subset S''}) \to \mathbf{R}^2 p_{T*}(\mathcal{C}(\mathcal{E}_T)\otimes_T I_{T'\subset T''})$$

so that

$$(5.8)\qquad ob(\mathcal{E}_{T'}, T', T'') = (\Gamma_T(c(f,S,S''))\circ f^*)ob(\mathcal{E}_{S'}, S', S'')$$

(iv) *Suppose there is a* $\mathbf{T}$*-action on* S *so that the family* $\mathcal{E}_S$ *is* $\mathbf{T}$*-equivariant. Then* $\mathbf{R}^2 p_{S*}(\mathcal{C}(\mathcal{E}_S, \mathcal{E}_S\otimes_{X_S} I))$ *has a natural* $\mathbf{T}$*-equivariant structure for any* $\mathbf{T}$*-equivariant* $\mathcal{O}_S$*-module* I. *Moreover, statements* (i), (ii) *hold for* $\mathbf{T}$*-equivariant base change, and the canonical morphism* (5.7) *is* $\mathbf{T}$*-equivariant if the deformation situations* $S \subset S' \subset S''$, $T \subset T' \subset T''$ *are* $\mathbf{T}$*-equivariant.*

Proof. Setting $k = 2$ in (5.4) we obtain

$$\mathbb{H}^2(X_S, \mathcal{C}(\mathcal{E}_S, I_{S'\subset S''})) = \Gamma_S(\mathbf{R}^2 p_{S*}\mathcal{C}(\mathcal{E}_S, I_{S'\subset S''})).$$

Recall that by construction, $\mathcal{C}(\mathcal{E}_S)$ is a finite complex of locally free $\mathcal{O}_{X_S}$ modules according to lemma (4.1). Moreover, definition (4.3) implies that there is an isomorphism of complexes of $\mathcal{O}_{X_S}$-modules

$$(5.9)\qquad \mathcal{C}(\mathcal{E}_S, I_{S'\subset S''}) \simeq \mathcal{C}(\mathcal{E}_S)\otimes_{X_S} I_{S'\subset S''}.$$

The vanishing lemma (4.10) and the base change theorem [34, Thm 7.7.5], [37, Thm III.12.11] imply that

$$\mathbf{R}^k p_{S*}\mathcal{C}(\mathcal{E}, I_{S'\subset S''}) = 0$$

for all $k \geq 3$ and

$$\mathbf{R}^2 p_{S*}(\mathcal{C}(\mathcal{E}, I_{S'\subset S''})) \simeq \mathbf{R}^2 p_{S*}(\mathcal{C}(\mathcal{E})\otimes_S I_{S'\subset S''}).$$

This proves (5.2.i). The morphism (5.6) is the canonical morphism provided by lemma (A.1.i).

In order to prove (5.2.*iii*) recall that

$$ob(\mathcal{E}_{S'}, S', S'') = \partial(\mathfrak{e}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'}))$$

where

$$\mathfrak{e}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'}) \in \mathbb{H}^1(X_S, \mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'}))$$

is the hypercohomology class constructed in proposition (4.5.*i*) and

$$\partial : \mathbb{H}^1(X_S, \mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'})) \to \mathbb{H}^2(X_S, \mathcal{C}(\mathcal{E}_S, I_{S'\subset S''})$$

is the connecting homomorphism determined by the short exact sequence of complexes of $\mathcal{O}_{X_S}$-modules (4.18). The later is reproduced below for convenience

$$0 \to \mathcal{C}(\mathcal{E}_S, I_{S'\subset S''}) \to \mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S''}) \to \mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'}) \to 0.$$

In the situation given in (5.2.*iii*) we have an analogous exact sequence complexes of $\mathcal{O}_{X_T}$-modules

$$0 \to \mathcal{C}(\mathcal{E}_T, I_{T'\subset T''}) \to \mathcal{C}(\mathcal{E}_T, \mathcal{E}_{T'}, I_{T\subset T''}) \to \mathcal{C}(\mathcal{E}_T, \mathcal{E}_{T'}, I_{T\subset T'}) \to 0.$$

which yields a similar connecting homomorphism. Since the projections $S' \to S$, respectively $T' \to T$ are finite morphisms, and $\mathcal{E}_{T'} = {f'_X}^* \mathcal{E}_{S'}$, the base change theorem implies that we have a canonical isomorphism of $\mathcal{O}_{X_T}$ modules

$$f_X^* \mathcal{E}_{S'} \simeq \mathcal{E}_{T'}.$$

This induces an isomorphism of complexes of $\mathcal{O}_T$-modules

$$f_X^* \mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'}) \simeq \mathcal{C}(\mathcal{E}_T, \mathcal{E}_{T'}, f^* I_{S\subset S'}). \tag{5.10}$$

However we also have a natural morphisms of $f'^* I_{S\subset S'} \to I_{T\subset T'}$, $f''^* I_{S\subset S''} \to I_{T\subset T''}$, $f''^* I_{S'\subset S''} \to I_{T'\subset T''}$ of $\mathcal{O}_{T'}$, $\mathcal{O}_{T''}$ and respectively $\mathcal{O}_{T''}$-modules (see for example [37, II.7.12.2]) which are also morphisms of $\mathcal{O}_T$-modules. Therefore we obtain a commutative diagram of complexes of $\mathcal{O}_{X_T}$-modules of the form

(5.11)

$$\begin{tikzcd}
f_X^* \mathcal{C}(\mathcal{E}_S, I_{S'\subset S''}) \arrow[r] \arrow[d] & \mathcal{C}(\mathcal{E}_T, I_{T'\subset T''}) \arrow[d] \\
f_X^* \mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S''}) \arrow[r] \arrow[d] & \mathcal{C}(\mathcal{E}_T, \mathcal{E}_{T'}, I_{T\subset T''}) \arrow[d] \\
f_X^* \mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'}) \arrow[r] & \mathcal{C}(\mathcal{E}_T, \mathcal{E}_{T'}, I_{T\subset T'}).
\end{tikzcd}$$

Note that the right column is exact while the left column is not since f_X^* is not an exact functor. Then we further obtain a natural commutative diagram of hypercohomology groups

(5.12)

$$\begin{tikzcd}
\mathbb{H}^1(X_S, \mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'})) \arrow[r] \arrow[d, "\partial"'] & \mathbb{H}^1(X_S, \mathcal{C}(\mathcal{E}_T, \mathcal{E}_{T'}, I_{T\subset T'})) \arrow[d, "\partial"'] \\
\mathbb{H}^2(X_S, \mathcal{C}(\mathcal{E}_S, I_{S'\subset S''})) \arrow[r] & \mathbb{H}^2(X_T, \mathcal{C}(\mathcal{E}_T, I_{T'\subset T''})).
\end{tikzcd}$$

Let the morphism (5.7) be the tensor product of the canonical morphism (5.6) and the canonical morphism of $\mathcal{O}_T$-modules $f'^*I_{S'\subset S} \to I_{T'\subset T}$. The bottom horizontal arrow of (5.12) is then identified by construction with $\Gamma_T(c(f,S',S''))\circ f^*$.

Moreover, we claim that the top horizontal arrow of (5.12) maps the extension class $\mathfrak{e}'_S$ to $\mathfrak{e}'_T$. This follows by observing that the isomorphism (5.10) yields a natural morphism of hypercohomology groups

$$f_X^* : \mathbb{H}^1(X_S, \mathcal{C}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'})) \to \mathbb{H}^1(X_T, \mathcal{C}(\mathcal{E}_T, \mathcal{E}_{T'}, f^*I_{S\subset S'}))$$

Since S is affine we can choose the affine Čech cover $\mathcal{U}_S$ in the proof of proposition (4.5) to be of the form $\mathcal{U}_S = \{(U_\alpha)_S\}$ where $\{U_\alpha\}$ is an affine Čech cover of X. Since T is also affine, by base change we obtain an affine Čech cover $\mathcal{U}_T = \{(U_\alpha)_T\}$ of X_T. Then it is obvious that the construction of the hypercohomology class $\mathfrak{e}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'})$ is compatible with base change, i.e. we have

$$\mathfrak{e}(\mathcal{E}_T, \mathcal{E}_{T'}, f^*I_{S\subset S'}) = f_X^*\mathfrak{e}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'}).$$

Moreover, it is also obvious by construction that the morphism

$$\mathbb{H}^1(X_T, \mathcal{C}(\mathcal{E}_T, \mathcal{E}_{T'}, f^*I_{S\subset S'})) \to \mathbb{H}^1(X_T, \mathcal{C}(\mathcal{E}_T, \mathcal{E}_{T'}, I_{T\subset T'}))$$

induced by the canonical morphism $f^*I_{S\subset S'} \to I_{T\subset T'}$ maps $\mathfrak{e}(\mathcal{E}_T, \mathcal{E}_{T'}, f^*I_{S\subset S'})$ to $\mathfrak{e}(\mathcal{E}_T, \mathcal{E}_{T'}, I_{T\subset T'})$.

□

Next we prove that the tangent-obstruction theory of any affine chart $S \to \mathfrak{M}_{ADHM}(\mathcal{X}, r, e)$ of finite type over $\mathbb{C}$ is perfect, verifying the conditions of [48, Def. 1.3].

Lemma 5.3. *Let $\mathcal{E}_S$ be a flat family of stable ADHM sheaves parameterized by an affine scheme S of finite type over $\mathbb{C}$. Then there exists a two-term complex $\mathbb{E}^\bullet = \left(\mathbb{E}^1_S \longrightarrow \mathbb{E}^2_S\right)$ of coherent locally free $\mathcal{O}_S$-modules with degrees $(1,2)$ so that*

$$\mathcal{H}^1(\mathbb{E}^\bullet_S \otimes_S I) \simeq \mathbf{R}^1 p_{S*}\mathcal{C}(\mathcal{E}_S, I)$$

$$\mathcal{H}^2(\mathbb{E}^\bullet_S \otimes_S I) \simeq \mathbf{R}^2 p_{S*}\mathcal{C}(\mathcal{E}_S) \otimes_S I$$

for any coherent $\mathcal{O}_S$-module I.

Moreover, if $S \to \mathfrak{M}_{ADHM}(\mathcal{X}, r, e)$ is a $\mathbf{T}$-equivariant chart, and I is a $\mathbf{T}$-equivariant coherent $\mathcal{O}_S$-module, the complex $\mathbb{E}^\bullet_S$ can be chosen $\mathbf{T}$-equivariant as well.

Proof. As observed in the proof of lemma (5.2.i), $\mathcal{C}(\mathcal{E}_S, I) = \mathcal{C}(\mathcal{E}_S) \otimes_{X_S} I$ for any coherent $\mathcal{O}_S$-module I. According to [34, Thm. 6.10.5] or [37, Prop. III.12.2] there exists a complex $\mathbb{F}^\bullet_S$ of finitely generated locally free $\mathcal{O}_S$-modules bounded above so that

$$\mathcal{H}^k(\mathbb{F}^\bullet_S \otimes_S I) = \mathbf{R}^k p_{S*}(\mathcal{C}(\mathcal{E}_S) \otimes_{X_S} I)$$

for any $\mathcal{O}_S$-module I and for all $k \in \mathbb{Z}$. Lemma (4.10) and the base change theorem [34, Thm 7.7.5] [37, Thm III.12.11] imply that

$$\mathbf{R}^k p_{S*}(\mathcal{C}(\mathcal{E}_S, \mathcal{E}_S)) = 0$$

for all $k \geq 3$ and

$$\mathbf{R}^2 p_{S*}(\mathcal{C}(\mathcal{E}_S) \otimes_{X_S} I) \simeq \mathbf{R}^2 p_{S*}\mathcal{C}(\mathcal{E}_S) \otimes_S I$$

for any coherent $\mathcal{O}_S$-module I.

Since $\mathbb{F}_S^\bullet$ is bounded above and has trivial cohomology in degrees $k \geq 3$, an easy induction argument based on [37, Prop. III.9.1A.f] shows that $\mathrm{Ker}(\mathbb{F}_S^2 \to \mathbb{F}_S^3)$ is a flat finitely generated $\mathcal{O}_S$-module. Therefore $\mathrm{Ker}(\mathbb{F}_S^2 \to \mathbb{F}_S^3)$ is a locally free $\mathcal{O}_S$-module, and we can truncate $\mathbb{F}_S^\bullet$ so that $\mathbb{F}_S^k = 0$ for $k \geq 3$.

Moreover, theorem [34, Thm. 6.10.5] also implies that for any closed point $s \in S$, the cohomology of the complex of vector spaces $\mathbb{F}_S^\bullet \otimes k(s)$ is isomorphic to the hypercohomology of the complex $\mathcal{C}(\mathcal{E}_S)|_{X_s}$. Since the later vanishes in degrees $k \leq 0$ according to lemma (4.10), it follows that the cohomology of $\mathbb{F}_S^\bullet \otimes k(s)$ is trivial in degrees $k \leq 0$ for any closed point $s \in S$. Then lemma (3.10) implies that we can truncate $\mathbb{F}_S^\bullet$ to a locally free complex $\mathbb{E}_S^\bullet$ satisfying the required properties.

Now suppose $S \to \mathfrak{M}_{ADHM}(\mathcal{X}, r, e)$ is $\mathbf{T}$-equivariant, hence in particular there is a $\mathbf{T}$-action on S so that $\mathcal{E}_S$ has a $\mathbf{T}$-equivariant structure. Then all above arguments hold in the equivariant setting, as observed for example in [70, Sect 2.2].

□

5.2. **Virtual Smoothness.** In this section we prove theorem (1.5) and corollary (1.6).

Proof of Theorem (1.5). We will construct a perfect tangent-obstruction theory on $\mathfrak{M}_{ADHM}(\mathcal{X}, r, e)$ as defined in [47, Def. 2.1]. According to lemma (3.11) we can choose a finite cover of $\mathfrak{M}_{ADHM}(\mathcal{X}, r, e)$ $\{\iota_\alpha : S_\alpha \to \mathfrak{M}_{ADHM}(\mathcal{X}, r, e)\}$, $\alpha \in \Lambda$ consisting of étale $\mathbf{T}$-equivariant morphisms, with S_α affine schemes of finite type over $\mathbb{C}$. According to lemmas (5.1), (5.2), (5.3), each chart (S_α, ι_α) carries a $\mathbf{T}$-equivariant perfect tangent-obstruction theory. Therefore we just have to check the compatibility conditions formulated in [47, Def. 2.1]. In the following we will denote by $\mathcal{E}_\alpha$ the $\mathbf{T}$-equivariant flat family determined by $\iota_\alpha : S_\alpha \to \mathfrak{M}_{ADHM}(\mathcal{X}, r, e)$, $\alpha \in \Lambda$. We will also denote by $p_{\alpha\beta\ldots} : X_{S_{\alpha\beta\ldots}} \to S_{\alpha\beta\ldots}$ the canonical projections for any overlap $S_{\alpha\beta\ldots}$ with $\alpha, \beta, \ldots \in \Lambda$, and by $\mathcal{E}_{\alpha\beta\ldots}$ the corresponding $\mathbf{T}$-equivariant flat families.

First we check that the collection of $\mathcal{O}_{S_\alpha}$-modules

$$\mathbf{R}^2 p_{S_\alpha *}\mathcal{C}(\mathcal{E}_\alpha)$$

with $\alpha \in \Lambda$ defines a $\mathbf{T}$-equivariant coherent sheaf on $\mathfrak{M}_{ADHM}(\mathcal{X}, r, e)$. Let

$$\begin{tikzcd}
S_{\alpha\beta} \arrow[r, "q_\alpha"] \arrow[d, "q_\beta"'] & S_\alpha \arrow[d, "\iota_\alpha"] \\
S_\beta \arrow[r, "\iota_\beta"] & \mathfrak{M}_{ADHM}(X, r, e)
\end{tikzcd}$$

be an arbitrary overlap. Since ι_α, ι_β are equivariant morphisms, there is a natural $\mathbf{T}$ action on $S_{\alpha\beta}$ so that q_α, q_β are also $\mathbf{T}$-equivariant. Moreover by construction we have a universal isomorphism

$$\xi_{\alpha\beta} : (q_\alpha)_X^*\mathcal{E}_\alpha \xrightarrow{\sim} (q_\beta)_X^*\mathcal{E}_\beta$$

of $\mathbf{T}$-equivariant $\mathcal{O}_{X_{S_{\alpha\beta}}}$-modules which yields a canonical isomorphism of $\mathbf{T}$-equivariant $\mathcal{O}_{X_{S_{\alpha\beta}}}$-modules

$$\eta_{\alpha\beta} : (q_\alpha)_X^*\mathcal{C}(\mathcal{E}_\alpha) \xrightarrow{\sim} (q_\beta)_X^*\mathcal{C}(\mathcal{E}_\beta).$$

Proceeding as in the proof of lemma (5.2.*ii*), the vanishing lemma (4.10) and the base change theorem [34, Thm 7.7.5], [8, Thm. III.3.4] yield canonical isomorphisms of $\mathbf{T}$-equivariant $\mathcal{O}_{S_{\alpha\beta}}$ modules.

$$q_\alpha^*\mathbf{R}^2 p_{\alpha *}\mathcal{C}(\mathcal{E}_\alpha) \simeq \mathbf{R}^2 p_{\alpha\beta *}(q_\alpha)_X^*\mathcal{C}(\mathcal{E}_\alpha)$$
$$q_\beta^*\mathbf{R}^2 p_{\beta *}\mathcal{C}(\mathcal{E}_\beta) \simeq \mathbf{R}^2 p_{\alpha\beta *}(q_\beta)_X^*\mathcal{C}(\mathcal{E}_\beta)$$

Then the isomorphism $\xi_{\alpha\beta}$ yields a canonical isomorphism

$$\phi_{\alpha\beta} : q_\alpha^*\mathbf{R}^2 p_{\alpha *}\mathcal{C}(\mathcal{E}_\alpha) \to q_\beta^*\mathbf{R}^2 p_{\beta *}\mathcal{C}(\mathcal{E}_\beta).$$

Next we have to check that the isomorphisms $\{\phi_{\alpha\beta}\}$ satisfy the cocycle condition. Given a triple overlap $S_{\alpha\beta\gamma}$ let $q_{\alpha\beta} : S_{\alpha\beta\gamma} \to S_{\alpha\beta}$, $q_{\alpha\gamma} : S_{\alpha\beta\gamma} \to S_{\alpha\gamma}$, $q_{\beta\gamma} : S_{\alpha\beta\gamma} \to S_{\beta\gamma}$ denote the canonical projections. Note that there exists a natural $\mathbf{T}$-action on $S_{\alpha\beta\gamma}$ so that these projections are $\mathbf{T}$-equivariant. Then we have to prove that

$$q_{\beta\gamma}^*\phi_{\beta\gamma} \circ q_{\alpha\beta}^*\phi_{\alpha\beta} = q_{\alpha\gamma}^*\phi_{\alpha\gamma} \tag{5.13}$$

This follows from the fact that there is a universal stable ADHM sheaf $\mathfrak{E}$ on $\mathfrak{M}_{ADHM}(\mathcal{X}, r, e) \times X$. This is standard for moduli stacks, and follows for example from the isomorphism (3.9) since R is equipped with a universal ADHM sheaf (see also [28, Ex. 2.49].) In particular the isomorphisms $\{\xi_{\alpha\beta}\}$ satisfy the cocycle condition, which implies that the isomorphisms $\{\eta_{\alpha\beta}\}$ also satisfy the cocycle condition. Then (5.13) follows again from vanishing results and the base change theorem. In fact the resulting obstruction sheaf $\mathfrak{O}b$ on $\mathfrak{M}_{ADHM}(X, r, e)$ is the direct image

$$\mathfrak{O}b = \mathbf{R}^2\mathfrak{p}_*\mathcal{C}(\mathfrak{E})$$

where $\mathcal{C}(\mathfrak{E})$ is the deformation complex of the universal object defined in (4.3) and $\mathfrak{p} : \mathfrak{M}_{ADHM}(X, r, e) \times X \to \mathfrak{M}_{ADHM}(X, r, e)$ is the canonical projection.

We also have to check that the obstruction assignments agree on overlaps. More precisely, suppose we are given two $\mathbf{T}$-equivariant deformation situations

$$S_\alpha \subset S'_\alpha \subset S''_\alpha$$
$$S_\beta \subset S'_\beta \subset S''_\beta$$

so that the induced deformation situations

$$S_{\alpha\beta} \subset S_{\alpha\beta} \times_{S_\alpha} S'_\alpha \subset S_{\alpha\beta} \times_{S_\alpha} S''_\alpha$$
$$S_{\alpha\beta} \subset S_{\alpha\beta} \times_{S_\beta} S'_\beta \subset S_{\alpha\beta} \times_{S_\beta} S''_\beta$$

are identical. We will denote the induced deformation situation on the overlap by

$$S_{\alpha\beta} \subset S'_{\alpha\beta} \subset S''_{\alpha\beta}.$$

and by $q'_\alpha : S'_{\alpha\beta} \to S'_\alpha$, $q'_\beta : S'_{\alpha\beta} \to S'_\beta$ the canonical equivariant projections. Suppose moreover that we are given $\mathbf{T}$-equivariant extensions $\mathcal{E}'_\alpha$ $\mathcal{E}'_\beta$ to S'_α, $S'_{\alpha\beta}$ respectively, and an isomorphism

$$\xi'_{\alpha\beta} : ({q'}_\alpha)^*_X \mathcal{E}'_\alpha \xrightarrow{\sim} ({q'}_\beta)^*_X \mathcal{E}'_\beta \tag{5.14}$$

which reduces to $\xi_{\alpha\beta}$ on $S_{\alpha\beta}$. Then according to lemma (5.2.*iii*) we have canonical $\mathbf{T}$-equivariant morphisms

$$c_\alpha : q_\alpha^* \mathbf{R}^2 p_{\alpha *}(\mathcal{C}(\mathcal{E}_\alpha) \otimes_{S_\alpha} I_{S'_\alpha \subset S''_\alpha}) \to \mathbf{R}^2 p_{\alpha\beta *}((q_\alpha)^*_X \mathcal{C}(\mathcal{E}_\alpha) \otimes_{S_{\alpha\beta}} I_{S'_{\alpha\beta} \subset S''_{\alpha\beta}})$$
$$c_\beta : q_\beta^* \mathbf{R}^2 p_{\alpha *}(\mathcal{C}(\mathcal{E}_\beta) \otimes_{S_\beta} I_{S'_\beta \subset S''_\beta}) \to \mathbf{R}^2 p_{\alpha\beta *}((q_\beta)^*_X \mathcal{C}(\mathcal{E}_\beta) \otimes_{S_{\alpha\beta}} I_{S'_{\alpha\beta} \subset S''_{\alpha\beta}})$$

Moreover we have

$$\mathfrak{o}b((q'_\alpha)^*_X \mathcal{E}'_\alpha, S'_{\alpha\beta}, S''_{\alpha\beta}) = (\Gamma_{S_{\alpha\beta}}(c_\alpha) \circ q_\alpha^*)\mathfrak{o}b(\mathcal{E}'_\alpha, S'_\alpha, S''_\alpha)$$
$$\mathfrak{o}b((q'_\beta)^*_X \mathcal{E}'_\beta, S'_{\alpha\beta}, S''_{\alpha\beta}) = (\Gamma_{S_{\alpha\beta}}(c_\beta) \circ q_\beta^*)\mathfrak{o}b(\mathcal{E}'_\beta, S'_\beta, S''_\beta)$$

and the isomorphism (5.14) yields an identification

$$\mathfrak{o}b((q'_\beta)^*_X \mathcal{E}'_\beta, S'_{\alpha\beta}, S''_{\alpha\beta}) = \Gamma_{S_{\alpha\beta}}(\phi_{\alpha\beta} \otimes 1_{I_{S'_{\alpha\beta} \subset S''_{\alpha\beta}}})\mathfrak{o}b((q'_\alpha)^*_X \mathcal{E}'_\alpha, S'_{\alpha\beta}, S''_{\alpha\beta}).$$

Therefore we obtain the required compatibility relation

$$(\Gamma_{S_{\alpha\beta}}(c_\beta) \circ q_\beta^*)\mathfrak{o}b(\mathcal{E}'_\beta, S'_\beta, S''_\beta) = \Gamma_{S_{\alpha\beta}}(\phi_{\alpha\beta} \otimes 1_{I_{S'_{\alpha\beta} \subset S''_{\alpha\beta}}}) \circ (\Gamma_{S_{\alpha\beta}}(c_\alpha) \circ q_\alpha^*)\mathfrak{o}b(\mathcal{E}'_\alpha, S'_\alpha, S''_\alpha)$$

for obstruction classes.

□

Proof of Corollary (1.6). According to proposition (3.15) the fixed locus $\mathfrak{M}_{ADHM}(X, r, e)^{\mathbf{T}}$ of the torus action defined in example (3.13) is a projective scheme over $\mathbb{C}$. By analogy with [31], taking the $\mathbf{T}$-fixed part of the perfect obstruction-theory of $\mathfrak{M}_{ADHM}(X, r, e)$ determines a perfect tangent-obstruction theory on the fixed locus, hence a virtual cycle

$$[\mathfrak{M}_{ADHM}(X, r, e)^{\mathbf{T}}] \in A_*(\mathfrak{M}_{ADHM}(X, r, e)^{\mathbf{T}}).$$

Alternatively, a direct argument proceeds as follows. Let $\{S_\alpha\}$, $\alpha \in \Lambda$ be an affine open cover of $\mathfrak{M}_{ADHM}(X, r, e)^{\mathbf{T}}$, $\alpha \in \Lambda$. Each S_α, $\alpha \in \Lambda$ parameterizes a flat family $\mathcal{E}_\alpha$ of stable ADHM sheaves on X satisfying the fixed point conditions formulated in the proof of proposition (3.15). Then it is straightforward to check that there is a natural $\mathbf{T}$-action on the hypercohomology groups $\mathbb{H}^k(X_{S_\alpha}, \mathcal{C}(\mathcal{E}_\alpha, I_\alpha))$, $k = 1, 2$ for any $\mathcal{O}_{S_\alpha}$-module I_α. Moreover, repeating proof of proposition (4.5), it follows that for any nilpotent thickening S'_α, there is a one-to-one correspondence between extension of $\mathcal{E}_\alpha$ to S'_α satisfying the $\mathbf{T}$-fixed condition and hypercohomology classes in the fixed part $\mathbb{H}^1(X, \mathcal{C}(\mathcal{E}_\alpha, I_\alpha))^f$, where I_α is the defining ideal of S_α in S'_α. Similarly, repeating the proof of proposition (4.9) in the $\mathbf{T}$-fixed context, it follows that the obstruction class $\mathfrak{ob}(\mathcal{E}_{S'_\alpha}, S_{\alpha'}, S''_\alpha)$ associated to a deformation situation of the form (4.16) lies in the fixed part $\mathbb{H}^2(X, \mathcal{C}(\mathcal{E}_\alpha, I_{S'_\alpha \subset S''_\alpha}))^f$.

Then lemmas (5.1), (5.2) imply that the $\mathbf{T}$-fixed part of the tangent-obstruction theory for $\mathfrak{M}_{ADHM}(X,r,e)$ determines a tangent-obstruction theory for the fixed locus $\mathfrak{M}_{ADHM}(X,r,e)^{\mathbf{T}}$. Moreover, it follows from lemma (5.3) and the proof of theorem (1.5) that this tangent-obstruction theory is perfect. In particular the required two-term complex of $\mathcal{O}_{S_\alpha}$-modules is the $\mathbf{T}$-fixed fixed of the equivariant complex $\mathbb{E}^\bullet_{S_\alpha}$ provided by lemma (5.3).

In order to construct the virtual normal bundle to the fixed locus recall that there exists a universal stable ADHM sheaf $\mathfrak{E}$ on $\mathfrak{M}_{ADHM}(X,r,e) \times X$. Let $\mathfrak{E}^{\mathbf{T}}$ denote its restriction to $\mathfrak{M}_{ADHM}(X,r,e)^{\mathbf{T}} \times X$ and let $\mathfrak{p}^{\mathbf{T}} : \mathfrak{M}_{ADHM}(X,r,e)^{\mathbf{T}} \times X \to \mathfrak{M}_{ADHM}(X,r,e)^{\mathbf{T}}$ denote the canonical projection. Let $\mathcal{C}(\mathfrak{E}^{\mathbf{T}})$ be deformation complex of $\mathfrak{E}^{\mathbf{T}}$ as defined in (4.3). Then an argument identical to lemma [61, Lemma 2.10] shows that there is a two term $\mathbf{T}$-equivariant complex $\mathbb{E}^\bullet$ of coherent locally free sheaves on $\mathfrak{M}_{ADHM}(X,r,e)^{\mathbf{T}}$ so that

$$\mathcal{H}^k(\mathbb{E}^\bullet) \simeq \mathbf{R}^k\mathfrak{p}^{\mathbf{T}}_* \mathcal{C}(\mathfrak{E}^{\mathbf{T}})$$

for $k = 1, 2$. We define the virtual normal bundle to be the $\mathbf{T}$-equivariant K-theory class

$$\begin{aligned} N^{vir}_{\mathfrak{M}_{ADHM}(X,r,e)^{\mathbf{T}}/\mathfrak{M}_{ADHM}(X,r,e)} &= [(\mathbb{E}^1)^m] - [(\mathbb{E}^2)^m] \\ &= [\mathbf{R}^1\mathfrak{p}^{\mathbf{T}}_*\mathcal{C}(\mathfrak{E}^{\mathbf{T}})^m] - [\mathbf{R}^2\mathfrak{p}^{\mathbf{T}}_*\mathcal{C}(\mathfrak{E}^{\mathbf{T}})^m]. \end{aligned} \tag{5.15}$$

where the superscripts m denote moving parts.

□

6. Admissible Pairs

Let X be a smooth projective curve over an infinite field k of characteristic 0, and M_1, M_2 fixed line bundles on X. Let Y be the total space of the projective bundle $\mathrm{Proj}(\mathcal{O}_X \oplus M_1 \oplus M_2)$. In this section we introduce new objects – called admissible pairs on Y – and prove that they are equivalent to stable pairs on Y as defined in [61]. In the next section we will show that admissible pairs are naturally related to stable ADHM sheaves on X by the relative Beilinson spectral sequence.

Let us first fix some notation. Let $\pi : Y \to X$ denote the canonical projection and $k_i = \deg(M_i)$, $i = 1, 2$. Recall that [37, Prop. II.7.1]

$$\pi_*(\mathcal{O}_Y(1)) = \mathcal{O}_X \oplus M_1 \oplus M_2. \tag{6.1}$$

Let $z_0 \in H^0(Y, \mathcal{O}_Y(1))$ denote the section corresponding to $1 \in H^0(X, \mathcal{O}_X)$ under the canonical isomorphism

$$H^0(Y, \mathcal{O}_Y(1)) \simeq H^0(X, \mathcal{O}_X) \oplus H^0(X, M_1) \oplus H^0(X, M_2).$$

We have analogous canonical sections

$$z_i \in H^0(Y, \pi^* M_i^{-1}(1)),$$

with $i = 1, 2$.

Let D_∞ be the zero locus of z_0 on Y, which will be referred to as the divisor at infinity. Let $H \in A_2(Y)$ denote the cycle class of D_∞ and let $F \in A_2(Y)$ denote the fiber class. Let $\sigma : X \to Y$ be the section determined by $z_1 = z_2 = 0$, and let $\beta = \sigma_*[X] \in A_1(Y)$. For future reference let us record the following easy lemma

Lemma 6.1. (*i*) $\mathrm{Ker}(A_1(Y) \xrightarrow{\cap H} A_0(Y)) = \langle \beta \rangle$

(*ii*) $\beta = H^2 - (k_1 + k_2)HF$ *in the intersection ring of* Y*.*

(*iii*) $c_1(Y) = 3H + (2 - 2g - k_1 - k_2)F$*.*

(*iv*) $\mathrm{Pic}(Y) \simeq \mathrm{Pic}(X) \times \mathbb{Z}$ *where the second direct summand is generated by the divisor class* $[D_\infty]$*.*

Proof. According to [27, Thm 3.3], $A_*(Y)$ is a free module over $A_*(X)$ generated by $\{1, H, H^2\}$. In particular $A_2(Y)$ is generated by H and the fiber class F. Let $\gamma = HF$ be a vertical curve class. Then an elementary computation yields the following intersection numbers

$$\begin{aligned} \beta H &= 0 & \beta F &= 1 \\ \gamma H &= 1 & \gamma F &= 0 \end{aligned}$$

Since the intersection pairing is nondegenerate, $A_1(Y)$ is generated by $\{\beta, \gamma\}$, and statement (*i*) above is obvious.

By construction the image of the section $s : X \to Y$ is cut by the equations $z_1 = 0$, $z_2 = 0$ on Y where $z_i \in H^0(Y, \pi^* M_i^{-1}(1))$, $i = 1, 2$ are the canonical sections defined above. Therefore we have

$$\beta = (H - k_1 F)(H - k_2 F) = H^2 - (k_1 + k_2)F$$

in the intersection ring of Y. This proves (*ii*).

The third statement follows by a straightforward computation using the relative Euler sequence. The fourth statement is standard, see for example [37, Ex. II.7.9].

□

Definition 6.2. (*i*) *Let* $d \in \mathbb{Z}_{\geq 1}$, $n \in \mathbb{Z}$*. An admissible pair of type* (d, n) *on* Y *is a pair* (Q, ρ) *consisting of a coherent* $\mathcal{O}_Y$*-module* Q *and a section* $\rho \in H^0(Y, Q)$ *satisfying the following conditions*

(*i*) ρ *is not identically zero.*

(*ii*) Q *is flat over* X*.*

(*iii*) $\mathrm{ch}_0(Q) = 0$, $\mathrm{ch}_1(Q) = 0$, $\mathrm{ch}_2(Q) = d\beta$ *and* $\chi(Q) = n$*.*

(iv) The cokernel $\mathrm{Coker}(\mathcal{O}_Y \xrightarrow{\rho} Q)$ *of the canonical morphism determined by* ρ *is an* $\mathcal{O}_Y$*-module of pure dimension 0.*

(ii) Two admissible pairs of type (d,n) (Q,ρ), (Q',ρ') *are isomorphic if there exists an isomorphism of* $\mathcal{O}_Y$*-modules* $u : Q \to Q'$ *so that the following diagram is commutative*

$$\begin{tikzcd}
\mathcal{O}_Y \arrow[r, "\rho"] \arrow[d, "1_{\mathcal{O}_Y}"'] & Q \arrow[d, "u"] \\
\mathcal{O}_Y \arrow[r, "\rho'"] & Q'
\end{tikzcd}$$

Remark 6.3. *(i) Note that conditions (ii), (iii) imply that the support* $\mathrm{supp}(Q)$ *is disjoint from* D_∞ *and finite over* X*.*

(ii) The admissible pairs (Q,ρ) *are closely related to the stable pairs considered in* [61]*. In the present context, a stable pair of type* (d,n) *on* Y *in the sense of* [61] *is a pair* (Q,ρ) *where* Q *is a coherent* $\mathcal{O}_Y$*-module of pure dimension one with* $\mathrm{ch}_2(Q) = d\beta$ *and* $\chi(Q) = n$ *and* $\rho \in H^0(Y,Q)$ *is a nonzero section so that* $\mathrm{Coker}(\mathcal{O}_Y \xrightarrow{\rho} Q)$ *is a* $\mathcal{O}_Y$*-module of pure dimension zero. The following lemma shows that we can replace condition (6.2.ii) by*

(ii′) Q *is of pure dimension one and* $\mathrm{supp}(Q)$ *is disjoint from* D_∞*.*

Therefore there is a one-to-one correspondence between admissible and respectively stable pairs of type (d,n) *on* Y*.*

Lemma 6.4. *Let* (Q,ρ) *be a pair consisting of a coherent* $\mathcal{O}_Y$*-module and a section* $\rho \in H^0(Y,Q)$ *satisfying conditions (i), (iii) and (iv) of definition (6.2). Then* Q *is flat over* X *if and only if it is of pure dimension one and* $\mathrm{supp}(Q)$ *is disjoint from* D_∞*.*

Proof. We first prove that Q flat over X implies that Q is of pure dimension one and $\mathrm{supp}(Q) \cap D_\infty = \emptyset$.

Since $\beta H = 0$ in the intersection ring of Y according to lemma (6.1.*i*) if $\mathrm{supp}(Q)$ has nontrivial intersection with D_∞, it follows that it must have vertical components. This would violate flatness over X. Therefore $\mathrm{supp}(Q)$ must be disjoint from D_∞.

Note that condition (*iii*) implies that $\mathrm{supp}(Q)$ is codimension 2 in Y. Then the torsion filtration [40, Def. 1.1.4] of Q reduces to

$$0 \subset T_0(Q) \subset T_1(Q) = Q$$

If $T_0(Q)_y \neq 0$ for some closed point $y \in Y$, it follows that $T_0(Q)_y$ is annihilated by any uniformizing parameter $\zeta \in \mathcal{O}_{X,x}$, $x = \pi(y)$. This would violate the local criterion for flatness [24, Thm. 6.8]. Therefore $T_0(Q)$ must be trivial.

Conversely, suppose Q is of pure dimension one and its support has no vertical components. Let $y \in Y$ be a closed point so that $Q_y \neq 0$, $x = \pi(y)$ and $\zeta \in \mathcal{O}_{X,x}$ a uniformizing parameter. Let Z be the support of Q and $\mathcal{I}_Z$ be the ideal sheaf of Z in Y. Since Z is one dimensional, $\mathcal{I}_{Z,y}$ has codimension 2 in $\mathcal{O}_{Y,y}$. Moreover, since Z has no vertical components, it follows that the ideal $\mathcal{J} \subset \mathcal{O}_{Y,y}$ generated by ζ and $\mathcal{I}_{Z,y}$ is codimension 3 in $\mathcal{O}_{Y,y}$. However note that $\mathcal{J}$ annihilates $\mathrm{Ker}(Q_y \xrightarrow{\zeta} Q_y)$. Therefore if $\mathrm{Ker}(Q_y \xrightarrow{\zeta} Q_y)$ is nontrivial, it must be a pure dimension zero submodule of Q. This contradicts the assumption that Q is pure dimension 1.

In conclusion, $\mathrm{Ker}(Q_y \xrightarrow{\zeta} Q_y)$ must be trivial for any $y \in Y$ so that $Q_y \neq 0$. Hence Q is flat over Y by the local criterion of flatness.

□

From now on we take the ground field k to be $\mathbb{C}$. Therefore in the following X is a smooth complex projective curve. Let S be a scheme of finite type over $\mathbb{C}$.

Definition 6.5. (*i*) *A flat family of admissible pairs on Y of type (d,n) parameterized by S is a pair (Q_S, ρ_S) consisting of a $\mathcal{O}_{Y_S}$-module Q_S and a section $\rho_S \in H^0(Y_S, Q_S)$ so that Q_S is flat over S and $(Q_S, \rho_S)|_{Y_s}$ is an admissible pair of type (d,n) on Y_s for each point $s \in S$.*

(*ii*) *An isomorphism of flat families (Q_S, ρ_S), (Q'_S, ρ'_S) of admissible pairs of type (d,n) is an isomorphism $u_S : Q_S \to Q'_S$ of $\mathcal{O}_{Y_S}$-modules so that the diagram*

$$\begin{tikzcd}
\mathcal{O}_{Y_S} \arrow[r, "\rho_S"] \arrow[d, "1_{\mathcal{O}_{Y_S}}"'] & Q_S \arrow[d, "u_S"] \\
\mathcal{O}_{Y_S} \arrow[r, "\rho'_S"] & Q'_S
\end{tikzcd}$$

is commutative.

Proof of Theorem (1.9) Let us define a fibered category $\mathfrak{M}_{Adm}(Y,d,n) \to \mathfrak{S}$ as follows.

- The objects of $\mathfrak{M}_{Adm}(Y,d,n)$ are flat families of admissible pairs of type (d,n) on Y parameterized by schemes S of finite type over $\mathbb{C}$.
- Given two objects (Q_S, ρ_S), (Q_T, ρ_T) a morphism in $\mathrm{Mor}((Q_S,\rho_S),(Q_T,\rho_T))$ is a pair (f, u_S) where $f : S \to T$ is a morphism of schemes of finite type over $\mathbb{C}$, and $u_S : (Q_S, \rho_S) \to (f_Y^* Q_T, f_Y^* \rho_T)$ is an isomorphism of flat families of admissible pairs on Y.

A routine argument shows that $\mathfrak{M}_{Adm}(Y,d,n)$ is a category fibered in groupoids over the category of schemes of finite type over $\mathbb{C}$.

Let $\mathfrak{M}_{St}(Y,d,n)$ be the groupoid of stable pairs (Q,ρ) of type (d,n) on Y. According to [45, 46, 61] there exists a projective moduli space $\mathfrak{M}_{St}(Y,d,n)$ parameterizing isomorphism classes of stable pairs of type (d,n) on Y. Since stable pairs have trivial stabilizers [61, Sect 2.2], this is a fine moduli space. Moreover, according to [61, Thm. 2.15] the moduli space $\mathfrak{M}_{St}(Y,d,n)$ has a natural perfect tangent-obstruction theory, therefore a virtual cycle $[\mathfrak{M}_{St}(Y,d,n)]$.

Lemma (6.4) identifies $\mathfrak{M}_{Adm}(Y,d,n)$ with an open subscheme of $\mathfrak{M}_{St}(Y,d,n)$, therefore $\mathfrak{M}_{Adm}(Y,d,n)$ is a quasi-projective scheme over $\mathbb{C}$ equipped with an induced perfect tangent-obstruction theory. Although the construction presented in [61] is based on the formalism of [10], it is fairly straightforward to provide an alternative construction of a perfect tangent-obstruction theory of $\mathfrak{M}_{St}(Y,d,n)$ relying on [47, Sect. 2.1]. Then [43, Prop. 3] implies that both constructions yield identical virtual cycles.

Let us recall the construction of tangent and respectively obstruction spaces, according to [69, Sect 3], [61, Sect 2]. Given a flat family (Q_S,ρ_S) of stable pairs, let $C(Q_S,\rho_S)$ denote the two term complex of $\mathcal{O}_{Y_S}$-modules

$$\mathcal{O}_{Y_S}\xrightarrow{\rho_S}Q_S \tag{6.2}$$

where the degrees are $0,1$ respectively. Let also $q_S : Y_S \to S$ denote the canonical projection. Suppose that S is affine of finite type over $\mathbb{C}$, and let S' be a trivial nilpotent extension of S determined by a coherent $\mathcal{O}_S$-module I. Then the tangent space to a flat family (Q_S,ρ_S) is the traceless ext group

$$\mathrm{Ext}^1_{Y_S}(C(Q_S,\rho_S),C(Q_S,\rho_S)\otimes_{Y_S}I)_0 \simeq \Gamma_S(\mathcal{E}xt^1_{q_S}(C(Q_S,\rho_S),C(Q_S,\rho_S)\otimes_{Y_S}I)_0)$$

Given a deformation situation of the form (4.16), a flat family (Q_S,ρ_S) and an extension $(Q_{S'},\rho_{S'})$ the obstruction space is the traceless ext group

$$\begin{aligned}\mathrm{Ext}^2_{Y_S}(C(Q_S,\rho_S),&C(Q_S,\rho_S)\otimes_{Y_S} I_{S'\subset S''})_0 \simeq\\ &\Gamma_S(\mathcal{E}xt^2_{q_S}(C(Q_S,\rho_S),C(Q_S,\rho_S))_0\otimes_S I_{S'\subset S''}).\end{aligned}$$

The obstruction class

$$\mathfrak{o}b(C(Q_{S'},\rho_{S'}),S,S'')\in \mathrm{Ext}^2_{Y_S}(C(Q_S,\rho_S),C(Q_S,\rho_S)\otimes_{Y_S} I_{S'\subset S''})_0$$

is given by

$$\mathfrak{o}b(C(Q_{S'},\rho_{S'}),S,S'') = \delta^2(\mathfrak{e}(C(Q_S,\rho_S),C(Q_{S'},\rho_{S'})),I_{S\subset S'})$$

where

$$\delta^2 : \mathrm{Ext}^1_{Y_S}(C(Q_S,\rho_S),C(QS',\rho_{S'})\otimes_{Y_S}I_{S\subset S'}) \to \mathrm{Ext}^2_{Y_S}(C(Q_S,\rho_S),C(Q_S,\rho_S)\otimes_{Y_S}I_{S'\subset S''})$$

is a natural coboundary morphism, and

$$\mathfrak{e}(C(Q_S,\rho_S),C(Q_{S'},\rho_{S'}),I_{S\subset S'})\in \mathrm{Ext}^1_{Y_S}(C(Q_S,\rho_S),C(Q_{S'},\rho_{S'})\otimes_{Y_S} I_{S\subset S'})$$

is the extension class corresponding to $C(Q_{S'},\rho_{S'})$. The later is given in turn by

$$\mathfrak{e}(C(Q_S,\rho_S),C(Q_{S'},\rho_{S'}),I_{S\subset S'}) = \delta^1(1_{C(Q_S,\rho_S)})$$

where

$$\delta^1 : \mathrm{Ext}^0_{Y_S}(C(Q_S,\rho_S),C(Q_S,\rho_S)) \to \mathrm{Ext}^1_{Y_S}(C(Q_S,\rho_S),C(Q_{S'},\rho_{S'})\otimes_{Y_S}I_{S\subset S'})$$

is again a natural coboundary morphism.

Note that $\delta^2(\mathfrak{e}(C(Q_S,\rho_S),C(Q_{S'},\rho_{S'}),I_{S\subset S'})$ belongs to the traceless ext group $\mathrm{Ext}^2_{Y_S}(C(Q_S,\rho_S),C(Q_S,\rho_S)\otimes_{Y_S} I_{S'\subset S''})_0$, which is canonically identified with a subgroup of $\mathrm{Ext}^2_{Y_S}(C(Q_S,\rho_S),C(Q_S,\rho_S)\otimes_{Y_S} I_{S'\subset S''})$ (see [69, Thm. 3.23, Thm 3.28].)

In order to prove that this data defines a perfect tangent-obstruction theory for $\mathfrak{M}_{St}(Y,d,n)$ we have to proceed again with a step-by-step verification of the conditions formulated in [48, Sect. 1], [48, Sect. 2] and [47, Def 2.1]. We will omit the details since they are very similar to [69, Sect. 3.1] as well as section (5) of the present paper.

Finally note that there is a $\mathbf{T} = \mathbb{C}^\times \times \mathbb{C}^\times$ action on $\mathfrak{M}_{St}(Y,d,n)$ induced by the natural fiberwise scaling action on $M_1 \oplus M_2$. Obviously this action preserves the open subspace $\mathfrak{M}_{Adm}(Y,d,n)$. Moreover, by analogy with section (5), the above construction easily generalizes to the equivariant setting, yielding a $\mathbf{T}$-equivariant perfect tangent-obstruction theory on $\mathfrak{M}_{Adm}(Y,d,n)$.

Note also that the fixed locus $\mathfrak{M}_{Adm}(Y,d,n)^{\mathbf{T}}$ is naturally identified with a closed subset of the fixed locus $\mathfrak{M}_{St}(Y,d,n)^{\mathbf{T}}$, which is proper over $\mathbb{C}$. More precisely, it is identified with the closed subset of the fixed locus $\mathfrak{M}_{St}(Y,d,n)^{\mathbf{T}}$ parameterizing $\mathbf{T}$-fixed stable pairs (Q,ρ) on Y so that the support of Q contains no vertical components. Therefore $\mathfrak{M}_{Adm}(Y,d,n)^{\mathbf{T}}$ is proper over $\mathbb{C}$ as well, and has a virtual cycle as well as an equivariant virtual normal bundle, determined by the perfect tangent-obstruction theory of $\mathfrak{M}_{Adm}(Y,d,n)$.

□

7. ADHM Sheaves via Relative Beilinson Spectral Sequence

Our next goal is to prove that there is an isomorphism between the moduli space of stable ADHM sheaves on X with data $\mathcal{X} = (M_1, M_2, \mathcal{O}_X)$ and the moduli space of admissible pairs on Y using a relative Beilinson monad construction. Moreover will prove that this isomorphism is compatible with the natural $\mathbf{T} = \mathbb{C}^\times \times \mathbb{C}^\times$ actions on these spaces and identifies the equivariant perfect obstruction theories.

Recall that under the current notation conventions we set $X_S = S \times X$, $Y_S = S \times Y$ for any scheme S over $\mathbb{C}$, and let $p_X : X_S \to X$, $p_Y : Y_S \to Y$, $\pi_S : Y_S \to X_S$ denote the canonical projections. We will also set $F_S = p_X^* F$, $G_S = p_Y^* G$ for any $\mathcal{O}_X$-module F, respectively $\mathcal{O}_Y$-module G. Note that Y_S is a projective scheme over X_S and we have canonical isomorphisms $\mathcal{O}_{Y_S}(1) \simeq p_Y^*\mathcal{O}_Y(1)$, $\Omega^k_{Y_S/X_S} \simeq p_Y^*\Omega^k_{Y/X}$, $k = 0,1,2$. Moreover, by base change we also have a canonical isomorphism

$$\pi_{S*}\mathcal{O}_{Y_S}(1) \simeq \mathcal{O}_{X_S} \oplus (M_1)_S \oplus (M_2)_S$$

Let

$$z_{S,0} \in H^0(Y_S, \mathcal{O}_{Y_S}(1)), \qquad z_{S,i} \in H^0(Y_S, \pi_S^*(M_i)_S^{-1}(1)),$$

$i = 1,2$, be canonical sections defined by analogy with z_0, z_1, z_2. Note that $z_{S,0}$, $z_{S,i}$, are canonically identified with $p_Y^* z_0$, $p_Y^* z_i$, respectively, for $i = 1,2$.

7.1. Relative Beilinson Monad for Admissible Pairs.

Lemma 7.1. *Let S be a scheme of finite type over $\mathbb{C}$ and let Q_S be an $\mathcal{O}_{Y_S}$-module, flat over S so that $Q_S|_{Y_s}$ satisfies conditions (ii), (iii) of definition*

(6.2). Then there is a spectral sequence of $\mathcal{O}_{Y_S}$-modules with

$$E_1^{i,j} = \mathcal{O}_{Y_S}(j) \otimes_{Y_S} \pi_S^* R^i \pi_{S*}(\Omega^{-j}_{Y_S/X_S}(-j) \otimes_{Y_S} Q_S), \tag{7.1}$$

$i \geq 0$, $j = -2, -1, 0$, converging to Q_S if $i = j = 0$ and 0 otherwise.

Proof. This is the relative Beilinson spectral sequence for Y_S/X_S as described for example in [59]. Let $\mathcal{K}_\Delta$

$$0 \to \mathcal{O}_Y(-2) \boxtimes \Omega^2_{Y/X}(2) \xrightarrow{\kappa_1} \mathcal{O}_Y(-1) \boxtimes \Omega^1_{Y/X}(1) \xrightarrow{\kappa_2} \mathcal{O}_Y \boxtimes \mathcal{O}_Y. \tag{7.2}$$

be the standard Koszul resolution of the diagonal $\Delta \subset Y \times_X Y$. Note that the projections

$$p_{1S}, p_{2S} : Y_S \times_{X_S} Y_S \to Y_S, \qquad p_{Y\times_X Y} : Y_S \times_{X_S} Y_S \simeq (Y \times_X Y)_S \to Y \times_X Y$$

are flat, and we have the obvious relations

$$p_Y \circ p_{1S} = p_1 \circ p_{Y\times_X Y}, \qquad p_Y \circ p_{2S} = p_2 \circ p_{Y\times_X Y}.$$

Then the complex $p^*_{Y\times_X Y}\mathcal{K}_\Delta$, which is canonically isomorphic to

$$0 \to \mathcal{O}_{Y_S} \boxtimes \Omega^2_{Y_S/X_S}(2) \xrightarrow{\kappa_{1S}} \mathcal{O}_{Y_S}(-1) \boxtimes \Omega^1_{Y_S/X_S}(1) \xrightarrow{\kappa_{2S}} \mathcal{O}_{Y_S} \boxtimes \mathcal{O}_{Y_S}, \tag{7.3}$$

is a locally free resolution of the diagonal $\Delta_S = S \times \Delta$. Let us denote by $\mathcal{K}_{\Delta_S}$ the complex (7.3), and by $\mathcal{K}^j_{\Delta_S}$, $-2 \leq j \leq 0$, its terms.

Since the Fourier-Mukai functor $\mathbf{R}p_{1S*}(\mathcal{K}_{\Delta_S} \otimes_{Y_S \times_{X_S} Y_S} p^*_{2S}\,\_\_)$ is isomorphic to the identity, there is a spectral sequence with

$$E_1^{i,j} = R^i p_{1S*}(\mathcal{K}^j_{\Delta_S} \otimes p^*_{2S} Q_S) \tag{7.4}$$

which converges to Q_S if $i + j = 0$ and 0 otherwise. Using the flat base change formula [37, Prop. III.9.3] [36, Prop. II.5.12] for the diagram

$$\begin{tikzcd} Y_S \times_{X_S} Y_S \arrow[r, "p_{2S}"] \arrow[d, "p_{1S}"'] & Y_S \arrow[d, "\pi_S"] \\ Y_S \arrow[r, "\pi_S"] & X_S \end{tikzcd} \tag{7.5}$$

a straightforward computation confirms (7.1).

□

Next we have to compute the terms in (7.1). We will need the following lemma.

Lemma 7.2. *Let $\rho : A \to B$, $\sigma : B \to C$ local morphisms of Noetherian local rings, k the residual field of A and M a finitely generated C-module. Suppose that B is flat over A. Then the following conditions are equivalent.*

(i) M is flat over B
(ii) M is flat over A and $M \otimes_A k$ is flat over $B \otimes_A k$.

Proof. [33, 10.2.5]

□

Given a flat family of admissible pairs (Q_S, ρ_S) parameterized by S, let $C(Q_S, \rho_S)$ denote the two term complex of $\mathcal{O}_{S\times Y}$-modules

$$\mathcal{O}_{Y_S} \xrightarrow{\rho_S} Q_S \tag{7.6}$$

with degrees $(0, 1)$.

Lemma 7.3. *Let (Q_S, ρ_S) be a flat family of admissible pairs parameterized by S and $C(Q_S, \rho_S)$ the complex defined in* (7.6)*. Then $C(Q_S, \rho_S)$ admits a canonical three term locally free resolution supported in degrees $-1, 0, 1$ of the form*

(7.7)

$$0 \to \mathcal{O}_{Y_S}(-2) \otimes_{Y_S} \pi_S^*(E_S \otimes_{X_S} (M_{12})_S)$$

$$\xrightarrow{\sigma_S} \mathcal{O}_{Y_S}(-1) \otimes_{Y_S} \pi_S^*(E_S \otimes_{X_S} (M_1)_S \oplus E_S \otimes_{X_S} (M_2)_S) \oplus \mathcal{O}_{Y_S} \xrightarrow{\tau_S} \pi_S^* E_S \to 0$$

where $E_S = R^0\pi_{S*}Q_S$ and

$$\begin{aligned} \sigma_S &= {}^t(z_{S,0} \otimes \pi_S^*\Phi_{S,2} - z_{S,2},\ -z_{S,0} \otimes \pi_S^*\Phi_{S,1} + z_{S,1},\ 0) \\ \tau_S &= (z_{S,0} \otimes \pi_S^*\Phi_{S,1} - z_{S,1},\ z_{S,0} \otimes \pi_S^*\Phi_{S,2} - z_{S,2},\ \pi_S^*\psi_S) \end{aligned}$$

for certain morphisms of $\mathcal{O}_{X_S}$-modules

$$\Phi_{S,i} : E_S \otimes_{X_S} (M_i)_S \to E_S, \qquad \psi_S : \mathcal{O}_{X_S} \to E_S$$

so that

$$\Phi_{S,1} \circ (\Phi_{S,2} \otimes 1_{(M_1)_S}) - \Phi_{S,2} \circ (\Phi_{S,1} \otimes 1_{(M_2)_S}) = 0 \tag{7.8}$$

and $\psi_S|_{X_s} : \mathcal{O}_X \to E_S|_{X_s}$ is injective for each point $s \in S$.

Proof. Let us first compute the terms in (7.1). Since Q_S is flat over S and $Q_S|_{X_s}$ is flat over X for any point $s \in S$, lemma (7.2) implies that Q_S is flat over X_S. Since it is also finite over X_S according to (6.3.i) the base change theorem [34, Thm 7.7.5], [8, Thm. III.3.4] implies that

$$R^i\pi_{S*}(\Omega^{-j}_{Y_S/X_S}(-j) \otimes_{Y_S} Q_S) = 0 \tag{7.9}$$

for all $i \geq 1$ and all $j = -2, -1, 0$. Moreover $R^0\pi_{S*}(\Omega^{-j}_{Y_S/X_S}(-j) \otimes_{Y_S} Q_S)$ are locally free $\mathcal{O}_{X_S}$-modules for all $j = -2, -1, -0$ [34, Cor. 7.9.10].

Then the first term of the spectral sequence (7.1) reads

(7.10)
$$\begin{tikzcd}
0 \arrow[d, "d_1^{0,-2}"] & \\
\mathcal{O}_{Y_S}(-2)\otimes_{Y_S}\pi_S^*R^0\pi_{S*}(\Omega^2_{Y_S/X_S}(2)\otimes_{Y_S}Q_S) \arrow[d, "d_1^{0,-1}"] & \\
\mathcal{O}_{Y_S}(-1)\otimes_{Y_S}\pi_S^*R^0\pi_{S*}(\Omega^1_{Y_S/X_S}(1)\otimes_{Y_S}Q_S) \arrow[d, "d_1^{0,0}"] & \\
\pi_S^*R^0\pi_{S*}Q_S \arrow[d, "d_1^{0,1}"] & \\
0 & 0
\end{tikzcd}$$

where we have used the notation $d_1^{i,j}: E_1^{i,j}\to E_1^{i,j+1}$, $i=0,1$, $j=-2,-1,0,1$ for differentials.

Note that the relative Euler sequence for $\pi_S: Y_S\to X_S$

$$0\to\Omega^1_{Y_S/X_S}(1)\to\pi_S^*(\mathcal{O}_{X_S}\oplus(M_1\oplus M_2)_S)\xrightarrow{\epsilon_S}\mathcal{O}_{Y_S}(1)\to 0 \tag{7.11}$$

yields an exact sequence of $\mathcal{O}_{X_S}$-modules of the form

$$\begin{aligned}
&0\to R^0\pi_{S*}(\Omega^1_{Y_S/X_S}(1)\otimes_{Y_S}Q_S)\\
&\quad\to R^0\pi_{S*}(\pi_S^*(\mathcal{O}_{X_S}\oplus(M_1\oplus M_2)_S)\otimes_{Y_S}Q_Sx)\xrightarrow{f_S}R^0\pi_{S*}(Q_S(1))\to 0.
\end{aligned}$$

The morphism f_S is of the form

$$f_S(t_0,t_1,t_2)=\Psi_{S,0}(t_1)+\Psi_{S,1}(t_1)+\Psi_{S,2}(t_2)$$

where

$$\begin{aligned}
\Psi_{S,0}&: R^0\pi_{S*}Q_S\to R^0\pi_{S*}(Q_S(1))\\
\Psi_{S,1}&: R^0\pi_{S*}Q_S\otimes_{X_S}(M_1)_S\to R^0\pi_{S*}(Q_S(1))\\
\Psi_{S,2}&: R^0\pi_{S*}Q_S\otimes_{X_S}(M_2)_S\to R^0\pi_{S*}(Q_S(1))
\end{aligned}\tag{7.12}$$

are induced by multiplication by $z_{S,0}, z_{S,1}, z_{S,2}$ respectively. Note that multiplication by $z_{S,0}$ induces an isomorphism $Q_S\simeq Q_S(1)$ since the support of Q_S is disjoint from $S\times D_\infty\subset Y_S$. Therefore $\Psi_{S,0}: R^0\pi_{S*}Q_S\to R^0\pi_{S*}(Q_S(1))$ is an isomorphism.

Then it follows that the morphism

$$\begin{aligned}
(M_1\oplus M_2)_S\otimes_{X_S}R^0\pi_{S*}Q_S&\to(\mathcal{O}_{X_S}\oplus(M_1\oplus M_2)_S)\otimes_{X_S}R^0\pi_{S*}Q_S\\
(t_1,t_2)&\to(-\Psi_{S,0}^{-1}\circ\Psi_{S,1}(t_1)-\Psi_{S,0}^{-1}\circ\Psi_{S,2}(t_2),t_1,t_2)
\end{aligned}$$

maps $(M_1\oplus M_2)_S\otimes_{X_S} R^0\pi_{S*}Q_S$ isomorphically to $\mathrm{Ker}(f_S)$. Therefore we obtain an identification

$$R^0\pi_{S*}(\Omega^1_{Y_S/X_S}(1)\otimes_{Y_S} Q_S)\simeq (M_1\oplus M_2)_S\otimes_{X_S} R^0\pi_{S*}Q_S \tag{7.13}$$

Note also that $\Omega^2_{Y_S/X_S}(2)\simeq \pi_S^*(M_{12})_S(-1)$. Therefore (7.10) reduces to a complex of $\mathcal{O}_{Y_S}$-modules of the form

$$\begin{aligned}&0\to \mathcal{O}_{Y_S}(-2)\otimes_{Y_S}\pi_S^*(R^0\pi_{S*}(Q_S(-1))\otimes_{X_S}(M_{12})_S)\\ &\xrightarrow{d_1^{0,-1}}\mathcal{O}_{Y_S}(-1)\otimes_{Y_S}\pi_S^*(R^0\pi_{S*}Q_S\otimes_{X_S}(M_1\oplus M_2)_S)\\ &\xrightarrow{d_1^{0,0}}\pi_S^*R^0\pi_{S*}Q_S\to 0\end{aligned}\tag{7.14}$$

The differentials $d_1^{0,-1}$, $d_1^{0,0}$ are determined by the morphisms κ_{1S},κ_{2S} in the Koszul complex (7.3). The later are determined in turn by contraction with the section $\Sigma_S\in H^0(Y_S\times_{X_S}Y_S,\mathcal{O}_{Y_S}(1)\boxtimes T_{Y_S/X_S}(-1))$ corresponding to the identity under the canonical identification

$$H^0(Y_S\times_{X_S}Y_S,\mathcal{O}_{Y_S}(1)\boxtimes T_{Y_S/X_S}(-1))\simeq \mathrm{End}_{X_S}(\mathcal{O}_{X_S}\oplus(M_1^{-1}\oplus M_2^{-1})_S).$$

Explicit expressions for κ_{1S},κ_{2S} can be written using the isomorphisms

$$\begin{aligned}&\Omega^2_{Y_S/X_S}(2)\simeq \pi_S^*(M_{12})_S(-1),\\ &\Omega^1_{Y_S/X_S}(1)\simeq \mathrm{Ker}\left(\pi_S^*(\mathcal{O}_{X_S}\oplus(M_1\oplus M_2)_S)\xrightarrow{\epsilon_S}\mathcal{O}_{Y_S}(1)\right).\end{aligned}$$

Then we have

$$\begin{aligned}\kappa_{1S}(u)=(&z_{S,1}\boxtimes z_{S,2}-z_{S,2}\boxtimes z_{S,1},\\ &-z_{S,1}\boxtimes z_{S,0}+z_{S,0}\boxtimes z_{S,1}, z_{S,2}\boxtimes z_{S,0}-z_{S,0}\boxtimes z_{S,2})\end{aligned}\tag{7.15}$$

where $z_{S,i}$, $i=1,2$, are identified with global sections of $\mathcal{O}_{Y_S}(1)\boxtimes\pi_S^*(M_i^{-1})_S$, $i=1,2$ via the obvious isomorphisms

$$(\mathcal{O}_{Y_S}(1)\otimes_{Y_S}\pi_S^*(M_i^{-1})_S\boxtimes\mathcal{O}_{Y_S}\simeq \mathcal{O}_{Y_S}(1)\boxtimes(\pi_S^*(M_i^{-1})_S).$$

A priori, the morphism (7.15) takes values in $\mathcal{O}_{Y_S}(-1)\boxtimes\pi_S^*(\mathcal{O}_{X_S}\oplus(M_1)_S\oplus(M_2)_S)$. However one can easily check that

$$(1\boxtimes\epsilon_S)(\kappa_{1S}(u))=0$$

for any local section u, hence κ_{1S} takes values in $\mathcal{O}_{Y_S}(1)\boxtimes\Omega^1_{Y_S/X_S}(1)$ as expected. The second morphism is locally given by

$$\kappa_{2S}(v_0,v_1,v_2)=p_{1S}^*(z_{S,0})v_0+p_{1S}^*(z_{S,1})v_1+p_{1S}^*(z_{S,2})v_2. \tag{7.16}$$

Using the identification (7.13) and expressions (7.15), (7.16) the complex (7.14) becomes isomorphic to

$$\begin{aligned}&0\to \mathcal{O}_{Y_S}(-2)\otimes_{Y_S}\pi_S^*(R^0\pi_{S*}(Q_S(-1))\otimes_{X_S}(M_{12})_S)\\ &\xrightarrow{\widetilde{\sigma}_S}\mathcal{O}_{Y_S}(-1)\otimes_{Y_S}\pi_S^*(R^0\pi_{S*}Q_S\otimes_{X_S}(M_1\oplus M_2)_S)\\ &\xrightarrow{\widetilde{\tau}_S}\pi_S^*R^0\pi_{S*}Q_S\to 0\end{aligned}\tag{7.17}$$

where

$$\begin{aligned}\widetilde{\sigma}_S &= {}^t(-z_{S,0}\otimes \pi_S^*\varPhi_{S,2} + z_{S,2}\otimes \pi_S^*\varPhi_{S,0},\ z_{S,0}\otimes \pi_S^*\varPhi_{S,1} - z_{S,1}\otimes \pi_S^*\varPhi_{S,0})\\ \widetilde{\tau}_S &= (-z_{S,0}\otimes \pi_S^*(\varPhi_{S,0}^{-1}\circ \varPsi_{S,1}) + z_{S,1},\ -z_{S,0}\otimes \pi_S^*(\varPhi_{S,0}^{-1}\circ \varPhi_{S,2}) + z_{S,2})\end{aligned}$$

and

$$\begin{aligned}\varPhi_{S,0} &: R^0\pi_{S*}(Q_S(-1)) \to R^0\pi_{S*}Q_S\\ \varPhi_{S,1} &: R^0\pi_{S*}(Q_S(-1))\otimes_{X_S}(M_1)_S \to R^0\pi_{S*}Q_S\\ \varPhi_{S,2} &: R^0\pi_{S*}(Q_S(-1))\otimes_{X_S}(M_2)_S \to R^0\pi_{S*}Q_S\end{aligned}\tag{7.18}$$

are induced by multiplication by $z_{S,0}, z_{S,1}, z_{S,2}$ respectively. Note that $\varPhi_{S,0}$ is an isomorphism since the support of Q_S is disjoint from D_∞. Moreover we have obvious commutation relations

$$\varPsi_{S,i}\circ \varPhi_{S,0} = \varPsi_{S,0}\circ \varPhi_{S,i}$$

for $i = 1, 2$.

The complex (7.17) is further isomorphic to

$$\begin{aligned}&0 \to \mathcal{O}_{Y_S}(-2)\otimes_{Y_S}\pi_S^*(E_S\otimes_{X_S}(M_{12})_S)\\ &\xrightarrow{\overline{\sigma}_S}\mathcal{O}_{Y_S}(-1)\otimes_{Y_S}\pi_S^*(E_S\otimes_{X_S}(M_1\oplus M_2)_S)\\ &\xrightarrow{\overline{\tau}_S}\pi_S^*E_S \to 0\end{aligned}\tag{7.19}$$

where $E_S = R^0\pi_*Q_S$,

$$\begin{aligned}\overline{\sigma}_S &= {}^t(-z_{S,0}\otimes \pi_S^*\Phi_{S,2} + z_{S,2},\ z_{S,0}\otimes \pi_S^*\Phi_{S,1} - z_{S,1})\\ \overline{\tau}_S &= (-z_{S,0}\otimes \pi_S^*\Phi_{S,1} + z_{S,1},\ -z_{S,0}\otimes \pi_S^*\Phi_{S,2} + z_{S,2}),\end{aligned}$$

and

$$\Phi_{S,i} = \varPsi_{S,0}^{-1}\circ \varPsi_{S,i} = \varPhi_{S,i}\circ \varPhi_{S,0}^{-1}\tag{7.20}$$

for $i = 1, 2$. Note that $\Phi_{S,1}, \Phi_{S,2}$ satisfy relation (7.8) since the morphisms $\varPhi_{S,1}, \varPhi_{S,2}$ in (7.12) are induced by multiplication by z_1, z_2 respectively, therefore they commute.

Returning to the spectral sequence (7.10) note that all higher differentials vanish for degree reasons. Therefore the cohomology sheaves of the the complex (7.19) must be trivial in degrees $-2, -1$ and isomorphic to Q_S in degree 0. Moreover, we claim that the natural evaluation map

$$\mathrm{ev}_S : \pi_S^*E_S \to Q_S$$

induces an isomorphism between the 0-th cohomology sheaf of the complex (7.19) and Q_S. In order to prove this, note first that the evaluation map is surjective since its restriction to the fiber Y_s is surjective for any $s \in S$. Furthermore, a simple computation shows that

$$\mathrm{Im}(\tau_S') \subseteq \mathrm{Ker}(\mathrm{ev}_S).$$

Therefore the evaluation map induces a surjective morphism of sheaves $\mathcal{H}^0 \to Q_S$ where $\mathcal{H}^0$ is the 0-th cohomology sheaf of (7.19). Since $\mathcal{H}^0 \simeq Q_S$, it follows that this morphism must be an isomorphism. This proves the claim.

Let us denote by $\mathbf{B}(Q_S)$ the $[-1]$ shift of the complex (7.19) and by

$$(0,0,\mathrm{ev}_S) : \mathbf{B}(Q_S) \to Q_S[-1]$$

the quasi-isomorphism induced by the evaluation map.

Next, note that by analogy with lemma (7.1) we have a spectral sequence with first term

$$\mathcal{O}_{Y_S}(j) \otimes_{Y_S} R^i\pi_{S*}\Omega^{-j}_{Y_S/X_S}(-j)$$

converging to $\mathcal{O}_{Y_S}$ for $i = j = 0$ and 0 otherwise. A simple calculation based on the relative Euler sequence and the base change theorem shows that this spectral sequence collapses to $\mathcal{O}_{Y_S}$ for $i = j = 0$ and 0 otherwise.

Now recall that for any coherent $\mathcal{O}_{Y_S}$-module F_S, the derived direct image $\mathbf{R}p_{1S*}(\mathcal{K}_{\Delta_S} \otimes_{Y_S \times_{X_S} Y_S} p_{2S}^*F_S)$ is computed by applying π_{1S*} to an injective resolution of the complex $\mathcal{K}_{\Delta_S} \otimes_{Y_S \times_{X_S} Y_S} p_{2S}^*F_S$. Since injective resolutions are functorial, it follows that the morphism $\rho_S : \mathcal{O}_{Y_S} \to Q_S$ yields a morphism of relative Beilinson monads

$$\mathcal{O}_{Y_S}[-1] \xrightarrow{\pi_S^*\psi_S} \mathbf{B}(Q_S)$$

where $\psi_S : \mathcal{O}_{X_S} \to E_S$ is the natural morphism of $\mathcal{O}_{X_S}$-modules obtained by pushing-forward ρ_S. Moreover, by construction we have diagram of complexes of $\mathcal{O}_{Y_S}$-modules

$$\begin{tikzcd}
\mathcal{O}_{Y_S}[-1] \arrow[r, "1_{\mathcal{O}_{Y_S}}{[-1]}"] \arrow[d, "\pi_S^*\psi_S"] & \mathcal{O}_{Y_S}[-1] \arrow[d, "\rho_S{[-1]}"] \\
\mathbf{B}(Q_S) \arrow[r, "{(0,0,\mathrm{ev}_S)}"] & Q_S[-1]
\end{tikzcd}$$

This yields a quasi-isomorphism of cones

$$\mathsf{Cone}(\ \mathcal{O}_{Y_S}[-1] \xrightarrow{\pi_S^*\psi_S} \mathbf{B}(Q_S)\ ) \to \mathsf{Cone}(\mathcal{O}_{Y_S} \xrightarrow{-\rho_S} Q_S)[-1]$$

given by $(0, 1_{\mathcal{O}_{Y_S}}, \mathrm{ev}_S)$ in degrees $(-1, 0, 1)$ respectively. We will denote this quasi-isomorphism by $(0, 1_{\mathcal{O}_{Y_S}}, \mathrm{ev}_S)$. Note that

$$C(\mathcal{O}_{Y_S} \xrightarrow{-\rho_S} Q_S)[-1] = C(Q_S, \rho_S).$$

Furthermore, note that for each point the restriction $\psi_S|_{X_s} : \mathcal{O}_{X_s} \to E_S|_{X_s}$ is nontrivial by construction since the morphism $\rho_S|_{Y_s} : CO_{Y_s} \to Q_S|_{Y_s}$ is nontrivial by definition. Therefore $\psi_S|_{X_s}$ is injective for any $s \in S$.

In conclusion, the cone $\mathsf{Cone}(\ \mathcal{O}_{Y_S}[-1] \xrightarrow{\pi_S^*\psi_S} \mathbf{B}(Q_S)\ )$ is the required resolution (7.7).

□

Definition 7.4. *(i) Given a flat family of admissible pairs* (Q_S, ρ_S) *parameterized by a scheme* S *of finite type over* $\mathbb{C}$*, we define the relative monad* $\mathbf{B}(Q_S, \rho_S)$ *associated to* (Q_S, ρ_S) *to be the three term complex of locally free* $\mathcal{O}_{Y_S}$*-modules* (7.7)*. In order to keep the notation short, we will denote the terms of this complex by*

$$0 \to \mathcal{B}_{-1}(Q_S, \rho_S) \xrightarrow{\sigma_S} \mathcal{B}_0(Q_S, \rho_S) \xrightarrow{\tau_S} \mathcal{B}_1(Q_S, \rho_S) \to 0. \tag{7.21}$$

(ii) Conversely, given a flat family of stable ADHM sheaves $\mathcal{E}_S$ *on* X *with* $E_\infty = \mathcal{O}_X$*, lemma (2.5) implies that* $\phi_S = 0$*. Then the data* $(E_S, \Phi_{S,1,2}, \psi_S)$ *determines a complex* $\mathbf{B}(\mathcal{E}_S)$ *of the form* (7.7)*. We will denote this complex by*

$$0 \to \mathcal{B}_{-1}(\mathcal{E}_S) \xrightarrow{\sigma_S} \mathcal{B}_0(\mathcal{E}_S) \xrightarrow{\tau_S} \mathcal{B}_1(\mathcal{E}_S) \to 0. \tag{7.22}$$

Note that we have used the same symbols for the differentials in (7.21), (7.22) *since the distinction will be clear from the context.*

For further reference, let

$$0 \to \mathcal{B}'_{-1}(\mathcal{E}_S) \xrightarrow{\sigma'_S} \mathcal{B}'_0(\mathcal{E}_S) \xrightarrow{\tau'_S} \mathcal{B}'_1(\mathcal{E}_S) \to 0 \tag{7.23}$$

be the three term subcomplex of $\mathbf{B}(\mathcal{E}_S)$ *obtained by removing the direct summand* $\mathcal{O}_{Y_S}$ *in the degree* 0 *term of* (7.7) *and truncating* σ_S, τ_S *accordingly. We will denote this complex by* $\mathbf{B}'(\mathcal{E}_S)$.

The following lemma proves that relative monad complexes are compatible with base change. This will be needed later in the proof of theorem (1.11).

Lemma 7.5. *Let* (Q_S, ρ_S) *be a flat family of admissible pairs parameterized by a scheme* S *of finite type over* $\mathbb{C}$*, and let* $C(Q_S, \rho_S)$ *be the complex* (7.6)*. Let* $f : T \to S$ *be a base change morphism, with* T *of finite type over* $\mathbb{C}$*, and let* $(Q_T, \rho_T) = (f_Y)^*(Q_S, \rho_S)$*. Then we have the following commutative diagrams*

(7.24)

$$\begin{tikzcd}
f_Y^*\mathbf{B}(Q_S) \arrow[r, "\simeq"] \arrow[d, "{(0,0,ev_S)}"'] & \mathbf{B}(Q_T) \arrow[d, "{(0,0,ev_T)}"] \\
f_Y^*Q_S \arrow[r, "1_{Q_T}"] & Q_T
\end{tikzcd}$$

$$\begin{tikzcd}
f_Y^*\mathbf{B}(Q_S, \rho_S) \arrow[r, "\simeq"] \arrow[d, "{(0,1_{\mathcal{O}_{Y_S}},ev_S)}"'] & \mathbf{B}(Q_T, \rho_T) \arrow[d, "{(0,1_{\mathcal{O}_{Y_T}},ev_T)}"] \\
f_Y^*C(Q_S, \rho_S) \arrow[r, "{(1_{\mathcal{O}_{Y_T}},1_{Q_T})}"] & Q_T
\end{tikzcd}$$

where the top rows are canonical isomorphisms.

Proof. Note that the Koszul resolution (7.3) is compatible with base change i.e.

$$f_{Y\times_X Y}^*\mathcal{K}_{\Delta_S} \simeq \mathcal{K}_{\Delta_T}.$$

Then lemma (A.1) implies that we have canonical morphisms of complexes

$$f_Y^*\mathbf{B}(Q_S) \to \mathbf{B}(Q_T), \qquad f_Y^*\mathbf{B}(Q_S, \rho_S) \to \mathbf{B}(Q_T, \rho_T). \tag{7.25}$$

Using the flat base change formula [37, Prop. III.9.3] [36, Prop. II.5.12] for the diagram (7.5), we have established in the proof of lemma (7.3) that all terms of the complexes $\mathcal{K}_{\Delta_S} \otimes_{Y_S \times_{X_S} Y_S} p_{2S}^* Q_S$, $\mathcal{K}_{\Delta_S} \otimes_{Y_S \times_{X_S} Y_S} p_{2S}^* \mathcal{O}_{Y_S}$ are acyclic with respect to push-forward by $p_{1S} : Y_S \times_{X_S} \to Y_S$. Then the base change theorem implies that the morphisms (7.25) are isomorphisms. The evaluation maps are naturally compatible with base change, therefore the diagrams (7.24) follow.

□

7.2. **Admissible Pairs and ADHM Sheaves.** Next we prove that there is a one-to-one correspondence between admissible pairs on Y and stable ADHM sheaves on X with $E_\infty = \mathcal{O}_X$. We start with a preliminary lemma summarizing the relevant properties of the complexes $\mathbf{B}(\mathcal{E}_S)$, $\mathbf{B}'(\mathcal{E}_S)$ defined in (7.4.*ii*).

Lemma 7.6. *Let X be a smooth projective curve over an infinite field k of characteristic 0, and let $\mathcal{E} = (E, \Phi_{1,2}, \psi)$ be a stable ADHM sheaf on X with $E_\infty = \mathcal{O}_X$. Then the following hold*

- (*i*) *$\sigma : \mathcal{B}_{-1}(\mathcal{E}) \to \mathcal{B}_0(\mathcal{E})$ is an injective morphism of $\mathcal{O}_Y$-modules and* Coker(σ) *is a coherent torsion free $\mathcal{O}_Y$-module.*
- (*ii*) *The middle cohomology sheaf $\mathcal{H}^0(\mathbf{B}'(\mathcal{E}))$ is trivial.*
- (*iii*) *The morphism of $\mathcal{O}_Y$-modules $\tau : \mathcal{B}_0(\mathcal{E}) \to \mathcal{B}_1(\mathcal{E})$ is surjective on the complement of a codimension three closed subset of Y.*
- (*iv*) *The restriction $\sigma|_{Y_x} : \mathcal{B}_{-1}(\mathcal{E})|_{Y_x} \to \mathcal{B}_0(\mathcal{E})|_{Y_x}$ to an arbitrary fiber Y_x, $x \in X$ is an injective morphism of $\mathcal{O}_{Y_x}$-modules and* Coker$(\sigma|_{Y_x})$ *is a coherent torsion free $\mathcal{O}_{Y_x}$-module.*
- (*v*) *The middle cohomology sheaf $\mathcal{H}^0(\mathbf{B}'(\mathcal{E})|_{Y_x})$ is trivial for any point $x \in X$.*

Proof. Suppose $\sigma : \mathcal{B}_{-1}(\mathcal{E}) \to \mathcal{B}_0(\mathcal{E})$ is not injective. Then Ker(σ) must be a nontrivial subsheaf of $\mathcal{B}_{-1}(\mathcal{E}) = \mathcal{O}_Y(-2) \otimes_Y \pi^*(E \otimes_X M_{12})$, which is a locally free $\mathcal{O}_Y$-module. Therefore Ker(σ) has to be torsion-free, hence locally free outside a codimension two closed subset of Y. Let Y_σ be the maximal open subset of Y so that Ker$(\sigma)|_{Y_\sigma}$ is a locally free $\mathcal{O}_{Y_\sigma}$-module. Since $Y_\sigma \subset Y$ is an open embedding, we have Ker$(\sigma|_{Y_\sigma}) =$ Ker$(\sigma)|_{Y_\sigma}$, therefore Ker$(\sigma|_{Y_\sigma})$ is a locally free $\mathcal{O}_{Y_\sigma}$-module as well, and we have a short exact sequence of $\mathcal{O}_{Y_\sigma}$-modules

$$0 \to \mathrm{Ker}(\sigma|_{Y_\sigma}) \to \mathcal{O}_{Y_\sigma}(-2) \otimes_{Y_\sigma} \pi^*(E \otimes_X M_{12})|_{Y_\sigma} \to \mathrm{Coker}(\sigma|_{Y_\sigma}) \to 0.$$

in which the first two terms are locally free. Then the associated $\mathcal{T}or^{\mathcal{O}_{Y_\sigma}}$ long exact sequence shows that

$$\mathcal{T}or_1^{\mathcal{O}_{Y_\sigma}}(\mathrm{Coker}(\sigma|_{Y_\sigma}), \mathcal{O}_y) = 0$$

where $\mathcal{O}_y$ is the structure sheaf of any point $y \in Y$. This implies that for any closed point $x \in X$, the induced map of $k(y)$-vector spaces

$$\sigma(y) : E(x) \otimes_{k(x)} k(y) \to (E(x) \oplus E(x) \oplus k(x)) \otimes_{k(x)} k(y)$$

has a nontrivial kernel for any point $y \in Y_\sigma \cap Y_x$. However, as observed in the proof [53, Lemma 2.7], for any $x \in X$, $\sigma(y)$ may have a nontrivial kernel at most at finitely many closed points $y \in Y_x$. Hence σ must be injective in order to avoid a contradiction.

In order to prove that $\mathrm{Coker}(\sigma)$ is torsion free, note that $\mathrm{Coker}(\sigma)_y$ is a free $\mathcal{O}_{Y,y}$-module at all points $y \in Y$ where $\sigma(y)$ is injective, according to lemma (3.10). Therefore $\mathrm{Coker}(\sigma)$ is locally free on the complement of a closed subset of codimension two. Let ω_Y be the dualizing sheaf of Y; since Y is smooth and projective ω_Y is locally free. The long exact $\mathcal{E}xt^\bullet_Y(\_\,,\omega_Y)$ sequence associated to the short exact sequence

$$0 \to \mathcal{B}_{-1}(\mathcal{E}) \xrightarrow{\sigma} \mathcal{B}_0(\mathcal{E}) \to \mathrm{Coker}(\sigma) \to 0$$

yields

$$\mathcal{E}xt^q_Y(\mathrm{Coker}(\sigma), \omega_Y) = 0$$

for all $q \geq 2$ since the first two terms are locally free. Moreover, the local ext sheaf $\mathcal{E}xt^1_Y(\mathrm{Coker}(\sigma), \omega_Y)$ must be supported in codimension 2 since $\mathrm{Coker}(\sigma)$ is locally free outside a codimension 2 closed subset. Then [40, Prop. 1.1.10] implies that $\mathrm{Coker}(\sigma)$ must be torsion free. This proves (7.6.i).

In order to prove that the middle cohomology sheaf $\mathcal{H}^0(\mathbf{B}'(\mathcal{E}))$ is trivial, it suffices to prove that its stalk $\mathcal{H}^0(\mathbf{B}'(\mathcal{E}))_y$ is trivial for any closed point $y \in Y$. This follows from a simple local computation. Given the explicit expressions of the differentials

$$\sigma' = {}^t(-z_0\otimes\pi^*\Phi_2+z_2, z_0\otimes\pi^*\Phi_1-z_1), \qquad \tau' = (-z_0\otimes\pi^*\Phi_1+z_1, -z_0\otimes\pi^*\Phi_2+z_2)$$

where Φ_1, Φ_2 satisfy (7.8), it follows that the complex $\mathbf{B}'(\mathcal{E})_y$ is exact in degree 0 if at least one of the morphisms of $\mathcal{O}_{Y,y}$-modules

$$(z_0 \otimes \pi^*\Phi_1 - z_1)_y, \qquad (z_0 \otimes \pi^*\Phi_2 - z_2)_y$$

is an isomorphism. Therefore $\mathcal{H}^0(\mathbf{B}'(\mathcal{E}))$ must be supported on the subset of Y where both endomorphisms fail to be isomorphisms, which is a codimension two condition on Y. However note that $\mathcal{H}^0(\mathbf{B}'(\mathcal{E}))$ is a subsheaf of $\mathrm{Coker}(\sigma')$. Since the last component of σ is trivial, we have

$$\mathrm{Coker}(\sigma) \simeq \mathrm{Coker}(\sigma') \oplus \mathcal{O}_Y,$$

therefore $\mathrm{Coker}(\sigma')$ must be torsion free since $\mathrm{Coker}(\sigma)$ is torsion-free according to (7.6.i). In conclusion the cohomology sheaf $\mathcal{H}^0(\mathbf{B}'(\mathcal{E}))$ must be trivial.

For the next claim, note that τ_y is surjective on stalks at a point $y \in Y$ if and only if the induced linear map of $k(y)$-vector spaces

$$\tau(y) : (E(x) \oplus E(x) \oplus k(x)) \otimes_{k(x)} k(y) \to E(x) \otimes_{k(x)} k(y)$$

is surjective, where $x = \pi(y)$. If $\tau(y)$ fails to be surjective at some $y \in Y$, it follows that the dual map $\tau(y)^\vee$ will fail to be injective. As observed in the proof of [53, Lemma 2.7], this implies the existence of a nontrivial proper linear subspace $W \subset E(x) \otimes_{k(x)} k(y)$ so that

$$(\Phi_{1,2}(x) \otimes 1_{k(y)})(W) \subseteq W, \qquad \mathrm{Im}(\psi(x) \otimes 1_{k(y)}) \subseteq W$$

Recall that in lemma (2.4) we have constructed a canonical destabilizing subsheaf $E_0 \subset E$. Note that $E_0(x)$ is a linear subspace of $E(x)$ for a generic closed point $x \in X$. Moreover, if a subspace $W \subset E(x) \otimes_{k(x)} k(y)$ as above exists for such a generic closed point $x \in X$, by construction we have $E_0(x) \otimes_{k(x)} k(y) \subset W(x)$. Therefore $\mathrm{rk}(E_0) < \mathrm{rk}(E)$, and E_0 must be a proper subsheaf of E in this case. In conclusion, $\tau(y)$ must be surjective for any point $y \in Y_x$, for generic $x \in X$. This implies that there must exist a closed subset $S_{\mathcal{E}} \subset X$ determined by $\mathcal{E}$ so that $\tau(y)$, hence also τ_y, is surjective for any $y \in \pi^{-1}(X \setminus S_{\mathcal{E}})$.

Now suppose $x \in S_{\mathcal{E}}$. The condition that $\tau|_{Y_x}$ fail to be surjective is a codimension two condition along Y_x. Therefore τ_y may fail to be surjective at most along a codimension three locus in Y. Moreover, it is easy to check that τ is surjective along D_∞, which is cut by $z_0 = 0$, hence the codimension three subset in question must be contained in $Y \setminus D_\infty$.

The proofs of (7.6.*iv*), (7.6.*v*) are analogous to the proofs of (7.6.*i*), (7.6.*ii*) respectively, therefore will be omitted.

□

Proposition 7.7. *Let X be a smooth projective curve over an infinite field k of characteristic 0. Then there is a one-to-one correspondence between admissible pairs (Q, ρ) on X with $\mathrm{ch}_2(Q) = d\beta$ and $\chi(Q) = n$ and stable ADHM sheaves $\mathcal{E}$ on X with $E_\infty = \mathcal{O}_X$ and*

$$r = d, \qquad e = n + d(g-1). \tag{7.26}$$

Moreover, two admissible pairs (Q, ρ), (Q', ρ') are isomorphic if and only if the corresponding ADHM sheaves $\mathcal{E}$, $\mathcal{E}'$ are isomorphic.

Proof. Given an admissible pair (Q, ρ) lemma (7.3) provides a relative monad $\mathbf{B}(Q, \rho)$. The data $(E, \Phi_{1,2}, \psi)$ defines an ADHM sheaf $\mathcal{E}$ with $E_\infty = \mathcal{O}_X$ and $\phi = 0$.

We claim that the resulting ADHM sheaf $\mathcal{E}$ must be stable. Suppose there exists a nontrivial saturated proper subsheaf $E' \subset E$ so that

$$\Phi_i(E' \otimes_X M_i) \subseteq E', \qquad \mathrm{Im}(\psi) \subseteq E' \tag{7.27}$$

for $i = 1, 2$. Note that E' must be torsion free, hence locally free, on X and $\mathrm{rk}(E') < \mathrm{rk}(E)$. Let $\Phi'_i = \Phi_i|_{E' \otimes M_i}$, $i = 1, 2$ and let $\psi' : \mathcal{O}_X \to E'$ denote the factorization of ψ through $E' \subset E$; note that $\mathrm{Im}(\psi') = \mathrm{Im}(\psi)$ as subsheaves of E'. Note that the sheaf inclusions (7.27) yield the following exact sequences of $\mathcal{O}_X$-modules

$$\begin{aligned} &\mathcal{T}or_1^{\mathcal{O}_X}(\mathrm{Coker}(\Phi_i'), \mathcal{O}_x) \to \mathrm{Im}(\Phi_i') \otimes_X \mathcal{O}_x \to E' \otimes_X \mathcal{O}_x \\ &\mathcal{T}or_1^{\mathcal{O}_X}(\mathrm{Coker}(\psi'), \mathcal{O}_x) \to \mathrm{Im}(\psi') \otimes_X \mathcal{O}_x \to E' \otimes_X \mathcal{O}_x \end{aligned} \tag{7.28}$$

where $\mathcal{O}_x$ is the structure sheaf of an arbitrary closed point $x \in X$.

According to [40, Def. 1.1.4], any coherent $\mathcal{O}_X$-module F admits a torsion sub-module $T_0(F)$ so that the quotient $F/T_0(F)$ is torsion free, hence locally free, on X. Then the long exact $\mathcal{T}or^{\mathcal{O}_X}$ sequence associated to the short exact sequence

$$0 \to T_0(F) \to F \to F/T_0(F) \to 0$$

implies that

$$\mathcal{T}or_1^{\mathcal{O}_X}(F, \mathcal{O}_x) \simeq \mathcal{T}or_1^{\mathcal{O}_X}(T_0(F), \mathcal{O}_x).$$

Therefore for any coherent $\mathcal{O}_X$ module F, $\mathcal{T}or_1^{\mathcal{O}_X}(F, \mathcal{O}_x)$ is trivial unless $x \in X$ belongs to the support of $T_0(F)$, which consists of finitely many points on X.

Applying this argument to $F = \mathrm{Coker}(\Phi_i')$, $\mathrm{Coker}(\psi')$ in (7.28) it follows that

$$\mathrm{Im}(\Phi_i'(x)) \subseteq E'(x) \qquad \mathrm{Im}(\psi(x)) \subseteq E'(x) \tag{7.29}$$

for all except finitely many closed points $x \in X$. Now for each point $y \in Y$, the morphism of $\mathcal{O}_{Y,y}$-modules τ_y is surjective if and only if the linear map of $k(y)$-vector spaces

$$\tau(y) : (E(x) \oplus E(x) \oplus k(x)) \otimes_{k(x)} k(y) \to E(x) \otimes_{k(x)} k(y)$$

is surjective. This is equivalent to the dual linear map $\tau(y)^\vee$ being injective.

However, as observed in the proof of lemma [53, Lemma 2.7], the existence of a proper nontrivial subspace $E'(x) \subset E(x)$ satisfying conditions (7.29) implies that $\tau(y)^\vee$ will fail to be injective at finitely many closed points in Y_x. In conclusion, the existence of a destabilizing subsheaf $E' \subset E$ implies that τ will fail to be surjective at finitely many closed points $y \in Y_x$ for generic $x \in X$.

Now, according to lemma (7.3) the complex (7.7) is quasi-isomorphic to the two term complex $C(Q, \rho)$. Therefore $\mathrm{Coker}(\tau) \simeq \mathrm{Coker}(\mathcal{O}_Y \xrightarrow{\rho} Q)$. This implies that the morphism τ in (7.7) must be surjective everywhere on Y except the support of $\mathrm{Coker}(\mathcal{O}_Y \xrightarrow{\rho} Q)$, which is dimension zero by definition. This contradicts the conclusion reached in the previous paragraph. Therefore a destabilizing proper subsheaf $E' \subset E$ as above cannot exist.

The relations (7.26) follow by a straightforward Grothendieck-Riemann-Roch computation using lemma (6.1).

Conversely, suppose we have a stable ADHM sheaf $(E, \Phi_{1,2}, \phi, \psi)$ of type $(r, e) = (d, n + d(g-1))$ with $E_\infty = \mathcal{O}_X$. According to (2.5) we must have $\phi = 0$. Then we construct the complex $\mathbf{B}(\mathcal{E}_S)$ of the form according to definition (7.4.*ii*). Note that we have a commutative diagram of $\mathcal{O}_Y$-modules with exact rows of the form

$$\begin{tikzcd}
0 \arrow[r] & \mathrm{Coker}(\sigma') \arrow[r] \arrow[d, "\overline{\tau}'"] & \mathrm{Coker}(\sigma) \arrow[r] \arrow[d, "\overline{\tau}"] & \mathcal{O}_Y \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & \mathcal{B}_1(\mathcal{E}) \arrow[r, "1"] & \mathcal{B}_1(\mathcal{E}) \arrow[r] & 0 \arrow[r] & 0
\end{tikzcd}$$

where $\overline{\tau}', \overline{\tau}$ are morphisms induced by τ', τ. Applying the snake lemma we obtain an exact sequence of $\mathcal{O}_Y$-modules of the form

$$0 \to \mathrm{Ker}(\overline{\tau}') \to \mathrm{Ker}(\overline{\tau}) \to \mathcal{O}_Y \to \mathrm{Coker}(\overline{\tau}') \to \mathrm{Coker}(\overline{\tau}) \to 0 \tag{7.30}$$

However lemma (7.6.*ii*) implies that $\mathrm{Ker}(\overline{\tau}') = 0$ since the complex $\mathbf{B}'(\mathcal{E})$ is exact in degree 0. Therefore it follows that

$$\mathrm{Ker}(\overline{\tau}) \simeq \mathrm{Ker}(\tau)/\mathrm{Im}(\sigma)$$

is a subsheaf of $\mathcal{O}_Y$. Moreover, it is easy to check that the morphism

$$\mathrm{Ker}(\tau)/\mathrm{Im}(\sigma)|_{D_\infty} \to \mathcal{O}_Y|_{D_\infty}$$

is an isomorphism. This implies that $\mathrm{Ker}(\overline{\tau})$ is a torsion-free rank one sheaf on Y with trivial determinant, therefore it must be the ideal sheaf of a closed subscheme $Z \subset Y$. Moreover the support of Z is disjoint from D_∞.

Set $Q = \mathrm{Coker}(\overline{\tau}')$ and let $\rho : \mathcal{O}_Y \to Q$ be the morphism obtained from (7.30). Since Q is the only nontrivial cohomology sheaf of the complex $\mathbf{B}'(\mathcal{E})$, , a straightforward computation using the first relation in lemma (6.1) yields

$$\mathrm{ch}_0(Q) = 0, \qquad \mathrm{ch}_1(Q) = 0, \qquad \mathrm{ch}_2(Q) = d\beta. \tag{7.31}$$

Hence Q satisfies condition (*iii*) of definition (6.2). Moreover, we claim that the morphism ρ is not identically zero. If it were trivial, we would obtain $\mathrm{Coker}(\overline{\tau}') \simeq \mathrm{Coker}(\overline{\tau})$ from (7.30). However $\mathrm{Coker}(\overline{\tau})$ is zero dimensional according to lemma (7.6.*iii*), which would contradict (7.31).

We also have to prove that Q is flat over X. Note that according to lemma (7.6.*v*) the restriction

$$\overline{\tau}'|_{Y_x} : \mathrm{Coker}(\sigma')|_{Y_x} \to \mathcal{B}_1(\mathcal{E})|_{Y_x}$$

is injective for all $x \in X$. Therefore $Q = \mathrm{Coker}(\overline{\tau}')$ is flat over X according to lemma (3.10).

The relations (7.26) follow again by a standard Grothendieck-Riemann-Roch computation using lemma (6.1).

Finally, compatibility with isomorphisms follows from a routine verification.

□

7.3. **Isomorphism of Moduli Spaces.** In this section we prove theorem (1.11). We first prove that the relative Beilinson spectral sequence yields a $\mathbf{T}$-equivariant isomorphism of algebraic spaces

$$\mathfrak{f} : \mathfrak{M}_{Adm}(Y, d, n) \simeq \mathfrak{M}_{ADHM}(\mathcal{X}, d, n + d(g-1)).$$

According to lemma (7.3), a flat family of admissible pairs (Q_S, ρ_S) parameterized by a scheme S of finite type over $\mathbb{C}$ determines an ADHM sheaf $\mathcal{E}_S = (E_S, \Phi_{S,1,2}, 0, \psi_S)$ on X_S. E_S is a locally free $\mathcal{O}_{X_S}$-module, hence it is flat over S. Lemma (7.5) and proposition (7.7) imply that $\mathcal{E}_S|_{X_s}$ is a stable ADHM sheaf for any point $s \in S$. Moreover, lemma (7.5) and proposition (7.7) further imply that this correspondence is compatible with isomorphisms and base change, therefore we obtain a functor $\mathfrak{f} : \mathfrak{M}_{Adm}(Y, d, n) \to \mathfrak{M}_{ADHM}(\mathcal{X}, d, n + d(g-1))$.

Conversely, suppose $\mathcal{E}_S = (E_S, \Phi_{S,1,2}, \phi_S, \psi_S)$ is a flat family of stable ADHM sheaves on X parameterized by S. According to lemma (2.5), the restriction $\phi_S|_{X_s}$ must be trivial for each closed point $s \in S$. Therefore $\phi_S = 0$.

Consider the three-term complex $\mathbf{B}(\mathcal{E}_S)$ defined in (7.4. *ii*). The restriction of $\mathbf{B}(\mathcal{E}_S)$ to each fiber Y_s, $s \in S$ is isomorphic to $\mathbf{B}(\mathcal{E}_s)$, where $\mathcal{E}_s = \mathcal{E}_S|_{Y_s}$. Therefore lemma (7.6.*i*) and lemma (3.10) imply that $\sigma_S : \mathcal{B}_{-1}(\mathcal{E}_S) \to \mathcal{B}_0(\mathcal{E}_S)$ is an injective morphism of $\mathcal{O}_{Y_S}$-modules, and $\mathrm{Coker}(\sigma_S)$ is flat over S. The same holds for $\sigma'_S : \mathcal{B}'_{-1}(\mathcal{E}_S) \to \mathcal{B}'_0(\mathcal{E}_S)$ by a similar argument. Let

$$\overline{\tau}'_S : \mathrm{Coker}(\sigma'_S) \to \mathcal{B}_1(\mathcal{E}_S)$$

be the morphism of $\mathcal{O}_{Y_S}$-modules induced by τ'_S. Then applying again (7.6.*v*) and lemma (3.10) it follows that $\mathrm{Ker}(\overline{\tau}'_S) = 0$ and $Q_S = \mathrm{Coker}(\overline{\tau}'_S)$ is flat over S. Furthermore, since

$$Q_S|_{Y_s} \simeq \mathrm{Coker}(\overline{\tau}'_S|_{Y_s}) \tag{7.32}$$

proposition (7.7) and lemma (7.2) imply that Q_S is flat over X_S and

$$\mathrm{ch}_0(Q_S|_{Y_s}) = 0, \qquad \mathrm{ch}_1(Q_S|_{Y_s}) = 0, \qquad \mathrm{ch}_2(Q_S|_{Y_s}) = d\beta, \qquad \chi(Q_S|_{Y_s}) = n$$

for any $s \in S$.

Now, applying the snake lemma as in the proof of the inverse implication of proposition (7.7) (above equation (7.30)) we obtain an exact sequence of $\mathcal{O}_{Y_S}$-modules of the form

$$0 \to \mathrm{Ker}(\overline{\tau}_S) \to \mathcal{O}_{Y_S} \xrightarrow{\rho_S} Q_S \to \mathrm{Coker}(\overline{\tau}_S) \to 0. \tag{7.33}$$

Using (7.33) and the fact that tensor product is right exact, we have isomorphisms of $\mathcal{O}_{Y_s}$-modules

$$\mathrm{Coker}(\rho_S|_{Y_s}) \simeq \mathrm{Coker}(\rho_S)|_{Y_s} \simeq \mathrm{Coker}(\overline{\tau}_S)|_{Y_s} \simeq \mathrm{Coker}(\overline{\tau}_S|_{Y_s}) \simeq \mathrm{Coker}(\tau_S|_{Y_s})$$

for any $s \in S$. Then proposition (7.7) implies that $\mathrm{Coker}(\rho_S|_{Y_s})$ is a sheaf of pure dimension 0 for any $s \in Y$. Therefore we conclude that (Q_S, ρ_S) is indeed a flat family of admissible pairs of type (d, n) parameterized by S.

It is again routine to check that this construction is compatible with isomorphisms and base change, therefore we obtain an inverse functor $\mathfrak{g} : \mathfrak{M}_{ADHM}(\mathcal{X}, d, n+d(g-1)) \to \mathfrak{M}_{Adm}(Y, d, n)$. By construction, the functors $\mathfrak{f}$, $\mathfrak{g}$ are inverse to each other. Therefore we have proven that the two stacks are indeed isomorphic. It is also easy to check that the functors $\mathfrak{f}$, $\mathfrak{g}$ are $\mathbf{T}$-equivariant.

7.4. **Comparison of Tangent-Obstruction Theories.** Next we prove that the perfect tangent-obstruction theories of the isomorphic moduli spaces $\mathfrak{M}_{Adm}(Y, d, n)$, $\mathfrak{M}_{ADHM}(\mathcal{X}, d, n + d(g-1))$, are compatible with the respect to the isomorphism $\mathfrak{f}$ in the sense of definition [47, Def 4.1]. More precisely, let $(\iota_\alpha : S_\alpha \to \mathfrak{M}_{Adm}(Y, d, n))$, $\alpha \in \Lambda$, be a finite cover of $\mathfrak{M}_{Adm}(Y, d, n)$ consisting of affine étale $\mathbf{T}$-equivariant charts. Note that $(\mathfrak{f} \circ \iota_\alpha : S_\alpha \to \mathfrak{M}_{ADHM}(\mathcal{X}, d, n + d(g-1)))$, $\alpha \in \Lambda$, is a similar cover of $\mathfrak{M}_{ADHM}(\mathcal{X}, d, n + d(g-1))$. Let $\mathcal{Q}_\alpha = (Q_\alpha, \rho_\alpha)$, respectively $\mathcal{E}_\alpha$ denote corresponding flat families of admissible pairs, respectively stable ADHM sheaves for all $\alpha \in \Lambda$. Then we will prove the following

(I) For each $\alpha \in \Lambda$ and any $\mathbf{T}$-equivariant coherent $\mathcal{O}_{S_\alpha}$-module I, there is a natural $\mathbf{T}$-equivariant isomorphism of tangent-obstruction theories associated to the data $(\mathcal{Q}_\alpha, I)$, $(\mathcal{E}_\alpha, I)$. These isomorphisms are furthermore compatible with base change morphisms of the form $f_\alpha : T_\alpha \to S_\alpha$.

(II) There exist $\mathbf{T}$-equivariant perfect tangent-obstruction data $(\mathbb{F}^\bullet_\alpha)$, $(\mathbb{E}^\bullet_\alpha)$ and $\mathbf{T}$-equivariant quasi-isomorphisms

$$\mathsf{q}_\alpha : \mathbb{F}^\bullet_\alpha \to \mathbb{E}^\bullet_\alpha$$

of complexes of $\mathcal{O}_{S_\alpha}$-modules so that

- (a) The induced isomorphisms in cohomology

  $$\mathcal{H}^k(\mathsf{q}_\alpha)(I) : \mathcal{H}^k(\mathbb{F}^\bullet_\alpha \otimes_{S_\alpha} I) \xrightarrow{\sim} \mathcal{H}^k(\mathbb{E}^\bullet_\alpha \otimes_{S_\alpha} I)$$

  $k = 1, 2$ agree with the natural isomorphisms constructed at point (I) above for all $\alpha \in \Lambda$.
- (b) For $k = 2$, the isomorphisms

  $$\mathcal{H}^2(\mathsf{q}_\alpha) : \mathcal{H}^2(\mathbb{F}^\bullet_\alpha) \xrightarrow{\sim} \mathcal{H}^k(\mathbb{E}^\bullet_\alpha)$$

  determine a global isomorphism of obstruction sheaves.
- (c) Given any deformation situation $S_\alpha \subset S'_\alpha \subset S''_\alpha$ as in (4.16), the obstructions $\mathfrak{o}b_\alpha(\mathcal{Q}_\alpha, S'_\alpha, S''_\alpha)$, $\mathfrak{o}b_\alpha(\mathcal{E}_\alpha, S'_\alpha, S''_\alpha)$ agree i.e.

  $$\mathcal{H}^2(\mathsf{q}_\alpha)(I_{S'_\alpha \subset S''_\alpha})(\mathfrak{o}b_\alpha(\mathcal{Q}_\alpha, S'_\alpha, S''_\alpha)) = \mathfrak{o}b_\alpha(\mathcal{E}_\alpha, S'_\alpha, S''_\alpha)$$

  for all $\alpha \in \Lambda$.

Lemma 7.8. (i) *Let (Q_S, ρ_S) be a flat family of admissible pairs parameterized by an affine scheme S of finite type over $\mathbb{C}$, and let I be a coherent $\mathcal{O}_S$-module. Then we have canonical isomorphisms*

(7.34)
$$\mathcal{E}xt^k_{q_S}(C(Q_S, \rho_S), C(Q_S, \rho_S) \otimes_{Y_S} q_S^* I)_0 \simeq \mathcal{E}xt^k_{q_S}(C(Q_S, \rho_S), C(Q_S, \rho_S)(-1) \otimes_{Y_S} q_S^* I)$$

for any $k \in \mathbb{Z}$, *where* $q_S : Y_S \to S$ *is the natural projection.*

(ii) *Given a base change morphism* $f : T \to S$, *with* T *affine of finite type over* $\mathbb{C}$, *a coherent* $\mathcal{O}_T$*-module* J *and a morphism of* $\mathcal{O}_T$*-modules* $\xi : f^*I \to J$, *we have a commuting diagram*

(7.35)

$$\begin{tikzcd}
f^*\mathcal{E}xt^k_{q_S}(C(Q_S,\rho_S), C(Q_S,\rho_S) \otimes_{Y_S} q_S^*I)_0 \arrow[r, phantom, "\simeq"] \arrow[d, "b^k(f,\xi)_0"'] & f^*\mathcal{E}xt^k_{q_S}(C(Q_S,\rho_S), C(Q_S,\rho_S)(-1) \otimes_{Y_S} q_S^*I) \arrow[d, "b^k(f,\xi)"] \\
\mathcal{E}xt^k_{q_T}(C(Q_T,\rho_T), C(Q_T,\rho_T) \otimes_{Y_T} q_T^*J)_0 \arrow[r, phantom, "\simeq"] & \mathcal{E}xt^k_{q_T}(C(Q_T,\rho_T), C(Q_T,\rho_T)(-1) \otimes_{Y_T} q_T^*J)
\end{tikzcd}$$

where $(Q_T, \rho_T) = f_Y^*(Q_S, \rho_S)$, *and the vertical arrows are natural base change morphisms.*

(iii) *Under the assumptions of* (i), (ii) *above, suppose moreover that the base schemes* S, T *are equipped with torus* $\mathbf{T}$ *actions so that* (Q_S, ρ_S), (Q_T, ρ_T), *as well as* I, J, $f : T \to S$ *and* $\xi : f^*I \to J$ *are* $\mathbf{T}$*-equivariant. Then the isomorphisms* (7.34), *as well as the base change diagrams* (7.35) *are* $\mathbf{T}$*-equivariant.*

Proof. According to lemma (7.3), the relative Beilinson monad $\mathbf{B}(Q_S, \rho_S)$ is a locally free resolution of $C(Q_S, \rho_S)$. Moreover we have shown in the proof of lemma (7.3) that there is a canonical quasi-isomorphism

$$(0, 1_{\mathcal{O}_{Y_S}}, \mathrm{ev}_S) : \mathbf{B}(Q_S, \rho_S) \to C(Q_S, \rho_S).$$

Therefore we have canonical isomorphisms

(7.36)

$$\mathcal{E}xt^k_{q_S}(C(Q_S,\rho_S), C(Q_S,\rho_S) \otimes_{Y_S} q_S^*I)_0 \simeq \mathcal{E}xt^k_{q_S}(\mathbf{B}(Q_S,\rho_S), \mathbf{B}(Q_S,\rho_S) \otimes_{Y_T} q_S^*I)_0$$
$$\mathcal{E}xt^k_{q_S}(C(Q_S,\rho_S), C(Q_S,\rho_S)(-1) \otimes_{Y_S} q_S^*I) \simeq \mathcal{E}xt^k_{q_S}(\mathbf{B}(Q_S,\rho_S), \mathbf{B}(Q_S,\rho_S)(-1) \otimes_{Y_S} q_S^*I)$$

Next, since the complex $\mathbf{B}(Q_S, \rho_S)$ is locally free, the trace map yields a morphism of complexes of $\mathcal{O}_{Y_S}$-modules

(7.37)

$$\begin{tikzcd}
0 \arrow[d] & 0 \arrow[d] \\
\mathcal{H}om_{Y_S}(\mathbf{B}(Q_S,\rho_S), \mathbf{B}(Q_S,\rho_S)(-1) \otimes_{Y_S} q_S^*I) \arrow[r, "\mathrm{tr}(-1)"] \arrow[d] & q_S^*I(-1) \arrow[d] \\
\mathcal{H}om_{Y_S}(\mathbf{B}(Q_S,\rho_S), \mathbf{B}(Q_S,\rho_S) \otimes_{Y_S} q_S^*I) \arrow[r, "\mathrm{tr}"] \arrow[d] & q_S^*I \arrow[d] \\
\mathcal{H}om_{Y_S}(\mathbf{B}(Q_S,\rho_S), \mathbf{B}(Q_S,\rho_S) \otimes_{Y_S} q_S^*I) \otimes_{Y_S} \mathcal{O}_{(D_\infty)_S} \arrow[r, "\mathrm{tr}_\infty"] \arrow[d] & q_S^*I \otimes_{Y_S} \mathcal{O}_{(D_\infty)_S} \arrow[d] \\
0 & 0
\end{tikzcd}$$

A straightforward calculation shows that there is a canonical quasi-isomorphism

$$\mathbf{B}(Q_S,\rho_S)\otimes_{Y_S}\mathcal{O}_{(D_\infty)_S}\to\mathcal{O}_{(D_\infty)_S}$$

and the bottom horizontal row of (7.37) is a quasi-isomorphism of complexes as well. In order to keep the notation short we will denote by $\mathcal{H}(-1),\mathcal{H},\mathcal{H}_\infty$ the terms in the left column of (7.37), starting from the top. Then note that the diagram (7.37) yields a commutative diagram of $\mathcal{O}_S$-modules of the form

$$\begin{tikzcd}
\mathbf{R}^{k-1}_{q_{S*}}(\mathcal{H}_\infty) \arrow[r, "\mathbf{R}^{k-1}q_{S*}(\mathrm{tr}_\infty)"] \arrow[d] & \mathbf{R}^{k-1}q_{S*}(q_S^*I\otimes_{Y_S}\mathcal{O}_{(D_\infty)_S}) \arrow[d] \\
\mathbf{R}^{k}q_{S*}(\mathcal{H}(-1)) \arrow[r, "\mathbf{R}^{k}q_{S*}(\mathrm{tr}(-1))"] \arrow[d] & \mathbf{R}^{k}q_{S*}(q_S^*I(-1)) \arrow[d] \\
\mathbf{R}^{k}q_{S*}(\mathcal{H}) \arrow[r, "\mathbf{R}^{k}q_{S*}(\mathrm{tr})"] \arrow[d] & \mathbf{R}^{k}q_{S*}(q_S^*I) \arrow[d] \\
\mathbf{R}^{k}_{q_{S*}}(\mathcal{H}_\infty) \arrow[r, "\mathbf{R}^{k}q_{S*}(\mathrm{tr}_\infty)"] & \mathbf{R}^{k}q_{S*}(q_S^*I\otimes_{Y_S}\mathcal{O}_{(D_\infty)_S})
\end{tikzcd}\tag{7.38}$$

in which the columns are exact and the horizontal morphisms of the form $\mathbf{R}^kq_{S*}(q_S^*I\otimes_{Y_S}\mathcal{O}_{(D_\infty)_S})$ are isomorphisms. Furthermore we have $q_S=p_S\circ\pi_S$, where $p_S:X_S\to S$, $\pi_S:Y_S\to X_S$ are the canonical projections. Then we have a Grothendieck spectral sequence

$$\mathbf{R}^mp_{S*}\mathbf{R}^l\pi_{S*}(q_S^*I(-1))\Rightarrow\mathbf{R}^{m+l}q_S\ast(q_S^*I(-1)).$$

Since $q_S^*I\simeq\pi_S^*p_S^*I$, and $\mathbf{R}^l\pi_{S*}\mathcal{O}_{Y_S}(-1)=0$ for all $l\in\mathbb{Z}$, the projection formula for the flat morphism π_S implies that all terms in the above Grothendieck spectral sequence vanish, therefore we have

$$\mathbf{R}^kq_{S*}(q_S^*I(-1))=0$$

for all $k\in\mathbb{Z}$. Then applying the snake lemma to the commutative diagram (7.38), we obtain isomorphisms

$$\mathrm{Ker}(\mathbf{R}^kq_{S*}(\mathrm{tr}))\simeq\mathbf{R}^kq_{S*}\mathcal{H}om_{Y_S}(\mathbf{B}(Q_S,\rho_S),\mathbf{B}(Q_S,\rho_S)(-1)\otimes_{Y_S}q_S^*I)$$

for all $k\in\mathbb{Z}$. Since $\mathbf{B}(Q_S,\rho_S)$ is a locally free resolution of $C(Q_S,\rho_S)$, this proves lemma (7.8.*i*).

Lemma (7.8.*ii*) follows from lemma (7.5) and lemma (A.1). The third part follows from the fact that all above calculations are also valid in the equivariant setting since $\mathbf{T}$ has trivial cohomology.

□

Lemma 7.9. (*i*) *Let* (Q_S,ρ_S), (Q'_S,ρ'_S) *be flat families of admissible pairs parameterized by a scheme* S *of finite type over* $\mathbb{C}$. *Let* $\mathcal{E}_S$, $\mathcal{E}'_S$ *be the corresponding flat families of stable ADHM sheaves. Then we have canonical* *isomorphisms*

$$\mathcal{E}xt^k_{q_S}(C(Q_S,\rho_S), C(Q'_S,\rho'_S)\otimes_{Y_S} q_S^*I) \simeq \mathbf{R}^k p_{S*}\widetilde{\mathcal{C}}(\mathcal{E}_S,\mathcal{E}'_S, I) \tag{7.39}$$

for any $k\in\mathbb{Z}$ and for any coherent $\mathcal{O}_S$-module I, where $q_S : Y_S \to S$ is the projection morphism $q_S = p_S\circ\pi_S : Y_S\to S$ and $\widetilde{\mathcal{C}}(\mathcal{E}_S,\mathcal{E}'_S, I)$ is the complex defined in (4.22).

(*ii*) *Under the assumptions of* (*i*) *above, let $f : T\to S$ be a base change morphism with T affine of finite type over $\mathbb{C}$, J be a coherent $\mathcal{O}_T$-module, and $\xi : f^*I\to J$ be a morphism of $\mathcal{O}_T$-modules. Let $(Q_T,\rho_T) = f_Y^*(Q_S,\rho_S)$, $(Q'_T,\rho'_T) = f_Y^*(Q'_S,\rho'_S)$ and let $\mathcal{E}_T,\mathcal{E}'_T$ be the flat families of stable ADHM sheaves corresponding to (Q_T,ρ_T), (Q'_T,ρ'_T) respectively. Then for any $k\in\mathbb{Z}$ we have a commutative diagram of $\mathcal{O}_T$-modules*

(7.40)

$$\begin{tikzcd}
f^*\mathcal{E}xt^k_{q_S}(C(Q_S,\rho_S), C(Q'_S,\rho'_S)\otimes_{Y_S} q_S^*I) \arrow[r, phantom, "\simeq"] \arrow[d, "b^k(f,\xi)"'] & f^*\mathbf{R}^k p_{S*}\widetilde{\mathcal{C}}(\mathcal{E}_S,\mathcal{E}'_S, I) \arrow[d, "c^k(f,\xi)"] \\
\mathcal{E}xt^k_{q_T}(C(Q_T,\rho_T), C(Q'_T,\rho'_T)\otimes_{Y_T} q_T^*J) \arrow[r, phantom, "\simeq"] & \mathbf{R}^k p_{T*}\widetilde{\mathcal{C}}(\mathcal{E}_T,\mathcal{E}'_T, J)
\end{tikzcd}$$

where the vertical arrows are natural morphism determined by (f,ξ).

(*iii*) *Under the assumptions of* (*i*), (*ii*) *above, suppose moreover that the base schemes S, T are equipped with torus $\mathbf{T}$ actions so that (Q_S,ρ_S), (Q'_S,ρ'_S) as well as I, J, $f : T\to S$ and $\xi : f^*I\to J$ are $\mathbf{T}$-equivariant. Then the isomorphisms* (7.39), *as well as the base change diagrams* (7.40) *are $\mathbf{T}$-equivariant.*

Proof. We will use again the canonical quasi-isomorphism

$$(0, 1_{\mathcal{O}_{Y_S}}, \mathrm{ev}_S) : \mathbf{B}(Q_S,\rho_S)\to C(Q_S,\rho_S)$$

constructed in the proof of lemma (7.3). This yields canonical isomorphisms

$$\begin{aligned}\mathcal{E}xt^k_{q_S}(C(Q_S,\rho_S), C(Q'_S,\rho'_S)\otimes_{Y_S} q_S^*I) \simeq{}& \\ \mathbf{R}^k q_{S*}(C(Q'_S,\rho'_S)\otimes_{Y_S}\mathbf{B}(Q_S,\rho_S)^\vee\otimes_{Y_S} q_S^*I)&\end{aligned} \tag{7.41}$$

for all $k\in\mathbb{Z}$. Next note that

$$C(Q'_S,\rho'_S)\otimes_{Y_S} q_S^*I = \mathsf{Cone}\left(q_S^*I \xrightarrow{-\rho'_S\otimes 1_{q_S^*I}} Q'_S\otimes_{Y_S} q_S^*I\right)[-1]$$

Therefore

$$\begin{gathered} C(Q'_S,\rho'_S)\otimes_{Y_S}\mathbf{B}(Q_S,\rho_S)^\vee\otimes_{Y_S} q_S^*I = \\ \mathsf{Cone}\left(\begin{tikzcd} \overline{\mathbf{B}(Q_S,\rho_S)^\vee}\otimes_{Y_S} q_S^*I \arrow[d, "-1_{\mathbf{B}(Q_S,\rho_S)}\otimes\rho'_S\otimes 1_{q_S^*I}"] \\ \mathbf{B}(Q_S,\rho_S)^\vee\otimes_{Y_S} Q'_S\otimes_{Y_S} q_S^*I \end{tikzcd}\right)[-1] \end{gathered} \tag{7.42}$$

where $\overline{\mathbf{B}(Q_S,\rho_S)^\vee}$ is the complex obtained by flipping the sign of all differentials in $\mathbf{B}(Q_S,\rho_S)^\vee$. Here we are using the sign conventions of [20, Ch. 1.3] as stated at the end of the introduction.

Using equation (7.7) and the projection formula for the flat morphism $\pi_S : Y_S \to X_S$, it is straightforward to check that all terms in the complex (7.42) are acyclic with respect to pushforward by π_S. This implies that the Grothendieck spectral sequence for the composition $q_S = p_S \circ \pi_S$ collapses to an isomorphism

$$\begin{aligned} &\mathbf{R}^k q_{S*}(C(Q'_S, \rho'_S) \otimes_{Y_S} \mathbf{B}(Q_S, \rho_S)^\vee \otimes_{Y_S} q_S^* I) \simeq \\ &\mathbf{R}^k p_{S*}\pi_{S*}(C(Q'_S, \rho'_S) \otimes_{Y_S} \mathbf{B}(Q_S, \rho_S)^\vee \otimes_{Y_S} q_S^* I) \end{aligned} \tag{7.43}$$

for each $k \in \mathbb{Z}$. Moreover, using (7.42) we obtain

$$\begin{aligned} &\pi_{S*}(\mathbf{B}(Q_S, \rho_S)^\vee \otimes_{Y_S} C(Q'_S, \rho'_S) \otimes_{Y_S} q_S^* I) = \\ &\mathsf{Cone}\left(\begin{tikzcd} \pi_{S*}\overline{\mathbf{B}(Q_S, \rho_S)^\vee} \otimes_{Y_S} q_S^* I \arrow[d, "\pi_{S*}(-1_{\mathbf{B}(Q_S,\rho_S)} \otimes \rho'_S \otimes 1_{q_S^* I})"] \\ \pi_{S*}\mathbf{B}(Q_S, \rho_S)^\vee \otimes_{Y_S} Q'_S \otimes_{Y_S} q_S^* I \end{tikzcd}\right)[-1] \end{aligned} \tag{7.44}$$

Next, we claim there is a quasi-isomorphism of complexes of $\mathcal{O}_{X_S}$-modules

$$\mathcal{O}_{X_S} \oplus \mathcal{H}om_{X_S}(E_S \otimes_{X_S} (M_{12})_S, p_S^* I)[-1] \to \pi_{S*}(\overline{\mathbf{B}(Q_S, \rho_S)^\vee} \otimes_{Y_S} q_S^* I) \tag{7.45}$$

This will be proven by a direct computation. Note that using the projection formula for the flat morphism $\pi_S : Y_S \to X_S$, it suffices to prove this claim for $I = \mathcal{O}_S$. $\overline{\mathbf{B}(Q_S, \rho_S)^\vee}$ is the following complex of locally free $\mathcal{O}_{Y_S}$-modules (7.46)

$$\begin{aligned} 0 \to \pi_S^* E_S^\vee &\xrightarrow{-\tau_S^\vee} \pi_S^*(E_S^\vee \otimes_{X_S} ((M_1^{-1})_S \oplus (M_2^{-1})_S))(1) \oplus \mathcal{O}_{Y_S} \\ &\xrightarrow{\sigma_S^\vee} \pi_S^*(E_S^\vee \otimes_{X_S} (M_{12}^{-1})_S)(2) \to 0 \end{aligned}$$

where

$$\begin{aligned} \tau_S^\vee &= {}^t(z_{S,0} \otimes \pi_S^*\Phi_{S,1}^\vee - z_{S,1}, z_{S,0} \otimes \pi_S^*\Phi_{S,2}^\vee - z_{S,2}, -\pi_S^*\psi_S^\vee) \\ \sigma_S^\vee &= (z_{S,0} \otimes \pi_S^*\Phi_{S,2}^\vee - z_{S,2}, -z_{S,0} \otimes \pi_S^*\Phi_{S,1}^\vee + z_{S,1}, 0). \end{aligned}$$

We also have

$$\begin{aligned} \pi_{S*}\mathcal{O}_{Y_S} &= \mathcal{O}_{X_S} \\ \pi_{S*}\mathcal{O}_{Y_S}(1) &= \mathcal{O}_{X_S} \oplus (M_1)_S \oplus (M_2)_S. \\ \pi_{S*}\mathcal{O}_{Y_S}(2) &= \mathcal{O}_{X_S} \oplus (M_1^{\otimes 2})_S \oplus (M_2^{\otimes 2})_S \oplus (M_1)_S \oplus (M_2)_S \oplus (M_{12})_S \end{aligned}$$

Therefore, applying π_{S*} to the complex (7.46) we obtain a complex of the form

$$\begin{tikzcd}
E_S^\vee \arrow[d, "-\pi_{S*}(\tau_S^\vee)"] \\
\begin{array}{c} E_S^\vee \oplus E_S^\vee \oplus E_S^\vee \otimes_{X_S} ((M_1^{-1})_S \oplus (M_2)_S^{-1}) \\ \oplus E_S^\vee \otimes_{X_S} ((M_1 \otimes_X M_2^{-1})_S \oplus (M_1^{-1} \otimes_X M_2)_S) \oplus \mathcal{O}_{X_S} \end{array} \arrow[d, "\pi_{S*}(\sigma_S^\vee)"] \\
\begin{array}{c} E_S^\vee \oplus E_S^\vee \otimes_{X_S} ((M_1^{-1})_S \oplus (M_2^{-1})_S) \\ \oplus E_S^\vee \otimes_{X_S} ((M_1 \otimes_X M_2^{-1})_S \oplus (M_1^{-1} \otimes_X M_2)_S) \oplus E_S^\vee \otimes_{X_S} (M_{12}^{-1})_S \end{array}
\end{tikzcd} \tag{7.47}$$

with degrees $-1, 0, 1$ respectively. The computation of the morphism $\pi_{S*}(\sigma_S^\vee(-1))$ is straightforward, taking into account that $\pi_{S*}(z_{S,0}) = \pi_{S*}(z_{S,1}) = \pi_{S*}(z_{S,2}) = 1$ by construction. In block form with respect to the above direct sum decompositions, we obtain

$$-\pi_{S*}(\tau_S^\vee) = {}^t\left(1_{E_S^\vee}, 1_{E_S^\vee}, -\Phi_{S,1}^\vee, -\Phi_{S,2}^\vee, 0, 0, \psi_S^\vee\right)$$

$\pi_{S*}(\sigma^\vee) =$

$$\begin{pmatrix}
-1_{E_S^\vee} & 1_{E_S^\vee} & 0 & 0 & 0 & 0 & 0 \\
0 & -\Phi_{S,1}^\vee & -1_{E_S^\vee \otimes (M_1^{-1})_S} & 0 & 0 & \Phi_{S,2}^\vee & 0 \\
\Phi_{S,2}^\vee & 0 & 0 & 1_{E_S^\vee \otimes (M_1^{-1})_S} & \Phi_{S,1}^\vee & 0 & 0 \\
0 & 0 & 0 & 0 & 1_{E_S^\vee \otimes (M_1 \otimes M_2^{-1})_S} & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 1_{E_S^\vee \otimes (M_1^{-1} \otimes M_2)_S} & 0 \\
0 & 0 & \Phi_{S,2}^\vee & \Phi_{S,1}^\vee & 0 & 0 & 0
\end{pmatrix}$$

Note that there is a morphism of complexes from the two term complex

$$\mathcal{O}_{X_S} \xrightarrow{0} E_S^\vee \otimes_{X_S} (M_{12}^{-1})_S$$

with degrees $0, 1$ to the complex (7.47) given by the obvious inclusions in degrees $0, 1$. Given the above explicit expression for the morphisms $\pi_{S*}(\sigma_S^\vee)$ a straightforward computation shows that this morphism is a quasi-isomorphism. Since E_S is a locally free $\mathcal{O}_{X_S}$-module, this proves the claim.

In order to determine the right hand side of equation (7.44), we also need to compute $\pi_{S*}(\mathbf{B}(Q_S, \rho_S)^\vee \otimes_{Y_S} Q'_S \otimes_{Y_S} q_S^* I)$. Again, since all terms of this complex are acyclic with respect to π_{S*} the computation reduces to a term-by-term application of π_{S*}. This is very similar to the derivation of the complex (7.19) in the proof of lemma (7.3), therefore the details will be omitted. The final result is a complex of the form

(7.48)

$$0 \to \mathcal{H}om_{X_S}(E_S, E'_S) \otimes_{X_S} p_S^*I \xrightarrow{d'_1} \begin{array}{c} \mathcal{H}om_{X_S}(E_S \otimes_{X_S} (M_1)_S, E'_S) \otimes_{X_S} p_S^*I \\ \oplus \\ \mathcal{H}om_{X_S}(E_S \otimes_{X_S} (M_2)_S, E'_S) \otimes_{X_S} p_S^*I \\ \oplus \\ \mathcal{H}om_{X_S}(\mathcal{O}_{X_S}, E'_S) \otimes_{X_S} p_S^*I \end{array}$$

$$\xrightarrow{d'_2} \mathcal{H}om_{X_S}(E_S \otimes_{X_S} (M_{12})_S, E'_S) \otimes_{X_S} p_S^*I \to 0$$

with degrees $-1, 0, 1$, where

$$\begin{aligned} d'_1(\alpha) =& (\alpha \circ \Phi_{S,1} - (\Phi'_{S,1} \otimes 1_I) \circ (\alpha \otimes 1_{(M_1)_S}), \\ & \alpha \circ \Phi_{S,2} - (\Phi'_{S,2} \otimes 1_I) \circ (\alpha \otimes 1_{(M_2)_S}), \alpha \circ \psi_S) \\ d'_2(\beta_1, \beta_2, \delta) =& -\beta_1 \circ (\Phi_{S,2} \otimes 1_{(M_1)_S}) + (\Phi'_{S,2} \otimes 1_I) \circ (\beta_1 \otimes 1_{(M_2)_S})) \\ & + \beta_2 \circ (\Phi_{S,1} \otimes 1_{(M_2)_S}) - (\Phi'_1 \otimes 1_I) \circ (\beta_2 \otimes 1_{(M_1)_S}) \end{aligned} \tag{7.49}$$

for any local section α of $\mathcal{H}om_{X_S}(E_S, E'_S) \otimes_{X_S} p_S^*I$ and any local sections $(\beta_1, \beta_2, \delta)$ of

$$\begin{array}{c} \mathcal{H}om_{X_S}(E_S \otimes_{X_S} (M_1)_S, E'_S) \otimes_{X_S} p_S^*I \\ \oplus \\ \mathcal{H}om_{X_S}(E_S \otimes_{X_S} (M_2)_S, E'_S) \otimes_{X_S} p_S^*I \\ \oplus \\ \mathcal{H}om_{X_S}(\mathcal{O}_{X_S}, E'_S) \otimes_{X_S} p_S^*I \end{array}$$

Taking into account the quasi-isomorphism (7.45) and equation (7.49), it follows that the right hand side of equation (7.44) is canonically quasi-isomorphic to the following complex of $\mathcal{O}_{X_S}$-modules

(7.50)

$$0 \to \begin{array}{c} \mathcal{H}om_{X_S}(E_S, E'_S) \otimes_{X_S} p_S^*I \\ \oplus \\ p_S^*I \end{array} \xrightarrow{d''{}_1} \begin{array}{c} \mathcal{H}om_{X_S}(E_S \otimes_{X_S} (M_1)_S, E'_S) \otimes_{X_S} p_S^*I \\ \oplus \\ \mathcal{H}om_{X_S}(E_S \otimes_{X_S} (M_2)_S, E'_S) \otimes_{X_S} p_S^*I \\ \oplus \\ \mathcal{H}om_{X_S}(E_S \otimes_{X_S} (M_{12})_S, \mathcal{O}_{X_S}) \otimes_{X_S} p_S^*I \\ \oplus \\ \mathcal{H}om_{X_S}(\mathcal{O}_{X_S}, E'_S) \otimes_{X_S} p_S^*I \end{array}$$

$$\xrightarrow{d''{}_2} \mathcal{H}om_{X_S}(E_S \otimes_{X_S} (M_{12})_S, E'_S) \otimes_{X_S} p_S^*I \to 0$$

where

(7.51)

$$\begin{aligned} d''{}_1(\alpha) =& (-\alpha \circ \Phi_{S,1} + (\Phi'_{S,1} \otimes 1_I) \circ (\alpha \otimes 1_{(M_1)_S}), \\ & -\alpha \circ \Phi_{S,2} + (\Phi'_{S,2} \otimes 1_I) \circ (\alpha \otimes 1_{(M_2)_S}), -\alpha \circ \psi_S + \psi'_S \circ \alpha_\infty) \\ d''{}_2(\beta_1, \beta_2, \delta) =& \beta_1 \circ (\Phi_{S,2} \otimes 1_{(M_1)_S}) - (\Phi'_{S,2} \otimes 1_I) \circ (\beta_1 \otimes 1_{(M_2)_S})) \\ & - \beta_2 \circ (\Phi_{S,1}, 1_{(M_2)_S}) + (\Phi'_1 \otimes 1_I) \circ (\beta_2 \otimes 1_{(M_1)_S}) + (\psi'_S \otimes 1_I) \circ \gamma \end{aligned}$$

for any local sections $(\alpha, \beta_1, \beta_1, \delta)$ as above and any local sections α_∞, γ of p_S^*I, $\mathcal{H}om_{X_S}(E_S \otimes_{X_S} (M_{12})_S, \mathcal{O}_{X_S}) \otimes_{X_S} p_S^*I$ respectively. Taking into account that $\phi_S = 0, \phi'_S = 0$ for families of stable ADHM sheaves with $E_\infty = \mathcal{O}_X$ the complex (7.50) is identical to the complex $\widetilde{\mathcal{C}}(\mathcal{E}_S, \mathcal{E}'_S, I)$ defined in ((4.22)).

The base change property follows again from the base change theorem and lemmas (7.5) and (A.1) by a routine argument. Finally the above computation is valid without any modification for equivariant flat families since the torus $\mathbf{T}$ has trivial group cohomology.

□

Lemma 7.10. *Under the conditions of lemma (7.9) we also have canonical isomorphisms*

$$\mathcal{E}xt^k_{q_S}(C(Q_S, \rho_S), C(Q'_S, \rho'_S)(-1) \otimes_{Y_S} q_S^*I) \simeq \mathbf{R}^k p_{S*}\mathcal{C}(\mathcal{E}_S, \mathcal{E}'_S, I) \tag{7.52}$$

for any $k \in \mathbb{Z}$, where $\mathcal{C}(\mathcal{E}_S, \mathcal{E}'_S, I)$ is the complex defined in (4.2)*. Moreover, the isomorphisms* (7.52) *satisfy compatibility conditions with base change and $\mathbf{T}$-equivariant structure analogous to to (7.9.ii), (7.9.iii).*

Proof. The proof is very similar to the proof of lemma (7.9) and will be omitted.

□

Lemma 7.11. *Consider a deformation situation of the form* (4.16)*. Let (Q_S, ρ_S) be a flat family of admissible pairs parameterized by an affine scheme S of finite type over $\mathbb{C}$, and let $(Q_{S'}, \rho_{S'})$ be an extension to S'. Let $\mathcal{E}_S, \mathcal{E}_{S'}$ respectively be the corresponding flat families of stable ADHM sheaves parameterized by S, S'. Then $\mathcal{E}_{S'}$ is an extension of $\mathcal{E}_S$ to S', and we have commutative diagrams of the form*

(7.53)

$$\begin{tikzcd}
\mathrm{Ext}^0_{Y_S}(C(Q_S, \rho_S), C(Q_S, \rho_S)) \arrow[r, "\simeq"] \arrow[d, "\delta^1"'] & \mathbb{H}^0(X_S, \widetilde{\mathcal{C}}(\mathcal{E}_S, \mathcal{E}_S)) \arrow[d, "\widetilde{\partial}^1"] \\
\mathrm{Ext}^1_{Y_S}(C(Q_S, \rho_S), C(Q_{S'}, \rho_{S'}) \otimes_{Y_S} I_{S\subset S'}) \arrow[r, "\simeq"] & \mathbb{H}^0(X_S, \widetilde{\mathcal{C}}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'}))
\end{tikzcd}$$

(7.54)

$$\begin{tikzcd}
\mathrm{Ext}^1_{Y_S}(C(Q_S, \rho_S), C(Q_{S'}, \rho_{S'}) \otimes_{Y_S} I_{S\subset S'}) \arrow[r, "\simeq"] \arrow[d, "\delta^2"'] & \mathbb{H}^1(X_S, \widetilde{\mathcal{C}}(\mathcal{E}_S, \mathcal{E}_{S'}, I_{S\subset S'})) \arrow[d, "\widetilde{\partial}^2"] \\
\mathrm{Ext}^2_{Y_S}(C(Q_S, \rho_S), C(Q_S, \rho_S) \otimes_{Y_S} I_{S'\subset S''}) \arrow[r, "\simeq"] & \mathbb{H}^2(X_S, \widetilde{\mathcal{C}}(\mathcal{E}_S, I_{S'\subset S''}))
\end{tikzcd}$$

where the horizontal arrows are induced by the isomorphisms (7.52)*, and the vertical arrows are coboundary morphisms.*

Moreover, these diagrams are compatible with base change and $\mathbf{T}$-equivariant structure.

Proof. Using the isomorphisms (7.41), the coboundary morphisms δ^1, δ^2 are determined by the following exact sequences of $\mathcal{O}_{Y_S}$-modules

$$\begin{aligned}
&0 \to C(Q_{S'},\rho_{S'}) \otimes_{Y_S} \mathbf{B}(Q_S,\rho_S)^\vee \otimes_{Y_S} I_{S\subset S'} \to C(Q_{S'},\rho_{S'}) \otimes_{Y_S} \mathbf{B}(Q_S,\rho_S)^\vee \\
&\quad \to C(Q_S,\rho_S) \otimes_{Y_S} \mathbf{B}(Q_S,\rho_S)^\vee \to 0. \\
&0 \to C(Q_S,\rho_S) \otimes_{Y_S} \mathbf{B}(Q_S,\rho_S)^\vee \otimes_{Y_S} I_{S'\subset S''} \to C(Q_{S'},\rho_{S'}) \otimes_{Y_S} \mathbf{B}(Q_S,\rho_S)^\vee \otimes_{Y_S} I_{S\subset S''} \\
&\quad \to C(Q_{S'},\rho_{S'}) \otimes_{Y_S} \mathbf{B}(Q_S,\rho_S)^\vee \otimes_{Y_S} I_{S\subset S'} \to 0.
\end{aligned}$$

As observed in the proof of lemma (7.9) all above complexes consist of acyclic terms with respect to push forward by $\pi_S : Y_S \to X_S$. Therefore applying π_{S*} to the above exact sequences we obtain the exact sequences of $\mathcal{O}_{X_S}$-modules

$$\begin{aligned}
&0 \to \widetilde{\mathcal{C}}(\mathcal{E}_S,\mathcal{E}_{S'},I_{S\subset S'}) \to \widetilde{\mathcal{C}}(\mathcal{E}_S,\mathcal{E}_{S'}) \to \widetilde{\mathcal{C}}(\mathcal{E}_S) \to 0 \\
&0 \to \widetilde{\mathcal{C}}(\mathcal{E}_S,I_{S'\subset S''}) \to \widetilde{\mathcal{C}}(\mathcal{E}_S,\mathcal{E}_{S'},I_{S\subset S''}) \to \widetilde{\mathcal{C}}(\mathcal{E}_S,\mathcal{E}_{S'},I_{S\subset S'}) \to 0
\end{aligned}$$

Now the lemma (7.11) follows from the fact that there is an isomorphism of derived functors

$$\mathbf{R}\Gamma_{Y_S} \simeq \mathbf{R}\Gamma_S \circ \mathbf{R}q_{S*} \simeq \mathbf{R}\Gamma_S \circ \mathbf{R}p_{S*} \circ \mathbf{R}\pi_{S*}$$

Compatibility with base change and equivariant structure follows again by analogy with lemma (7.9).

□.

Finally, we also have

Lemma 7.12. *Let (Q_S,ρ_S) be a flat family of admissible pairs on Y parameterized by an affine scheme S of finite type over $\mathbb{C}$ and let $\mathcal{E}_S$ be the corresponding family of stable ADHM sheaves. Then there exist two-term complexes of coherent locally free $\mathcal{O}_S$-modules $\mathbb{F}_S^\bullet = (F_S^1 \to F_S^2)$, $\mathbb{E}_S^\bullet = (E_S^1 \to E_S^2)$ so that*

$$\begin{aligned}
&\mathcal{H}^1(\mathbb{F}_S^\bullet \otimes_S I) \simeq \mathcal{E}xt^1_{q_S}(C(Q_S,\rho_S), C(Q_S,\rho_S)(-1)\otimes_{Y_S} I) \\
&\mathcal{H}^2(\mathbb{F}_S^\bullet \otimes_S I) \simeq \mathcal{E}xt^2_{q_S}(C(Q_S,\rho_S), C(Q_S,\rho_S)(-1)) \otimes_S I
\end{aligned} \tag{7.55}$$

$$\begin{aligned}
&\mathcal{H}^1(\mathbb{E}_S^\bullet \otimes_S I) \simeq \mathbf{R}^1 p_{S*}\mathcal{C}(\mathcal{E}_S, I) \\
&\mathcal{H}^2(\mathbb{E}_S^\bullet \otimes_S I) \simeq \mathbf{R}^2 p_{S*}\mathcal{C}(\mathcal{E}_S) \otimes_S I
\end{aligned} \tag{7.56}$$

for any coherent $\mathcal{O}_S$-module I, and a quasi-isomorphism $\mathsf{q} : \mathbb{F}_S^\bullet \to \mathbb{E}_S^\bullet$ so that the $\mathsf{q}\otimes 1_I$ induces the isomorphisms (7.52) in cohomology, for $k = 1,2$.

Moreover, if there is a $\mathbf{T}$-action on S so that (Q_S,ρ_S) is $\mathbf{T}$-equivariant, we can choose the complexes $\mathbb{F}_S^\bullet = (F_S^1 \to F_S^2)$, $\mathbb{E}_S^\bullet = (E_S^1 \to E_S^2)$ and the quasi-isomorphism q to be $\mathbf{T}$-equivariant.

Proof. Lemmas (7.3), (7.10) imply that we have an isomorphism

$$\begin{aligned}
&\mathbf{R}q_{S*}\mathbf{R}\mathcal{H}om_{Y_S}(C(Q_S,\rho_S), C(Q_S,\rho_S)(-1)) \simeq \\
&\mathbf{R}q_{S*}(B(Q_S,\rho_S)\otimes_{Y_S} B(Q_S,\rho_S)^\vee \otimes_{Y_S} I) \simeq \mathbf{R}p_{S*}\mathcal{C}(\mathcal{E}_S, I)
\end{aligned}$$

in the derived category $D^-(S)$, for any coherent $\mathcal{O}_S$-module I. In particular, there exist complexes of $\mathcal{O}_S$-modules $\widetilde{\mathbb{F}}^\bullet_S$, $\widetilde{\mathbb{E}}^\bullet_S$ bounded above so that

$$\begin{aligned} &\mathcal{H}^k(\widetilde{\mathbb{F}}^\bullet_S \otimes_S I) \simeq \mathcal{E}xt^k_{q_S}(C(Q_S,\rho_S), C(Q_S,\rho_S)(-1)\otimes_{Y_S} I) \\ &\mathcal{H}^k(\mathbb{E}^\bullet_S \otimes_S I) \simeq \mathbf{R}^k p_{S*}\mathcal{C}(\mathcal{E}_S, I) \end{aligned}$$

for $k = 1, 2$, and a quasi-isomorphism $\widetilde{\mathsf{q}} : \widetilde{\mathbb{F}}^\bullet_S \to \widetilde{\mathbb{E}}^\bullet_S$ which induces the isomorphisms (7.52). Then, employing the inductive construction of [37, Lemma III.12.3], we obtain complexes $\mathbb{F}'^\bullet_S$, $\mathbb{E}'^\bullet_S$ of coherent locally free $\mathcal{O}_S$-modules, bounded above, and a natural quasi-isomorphism $\mathsf{q}' : \mathbb{F}'^\bullet_S \to \mathbb{E}'^\bullet_S$ with the same properties. Moreover, using the same argument as in the proof of lemma (5.3), the complexes $\mathbb{F}'^\bullet_S$, $\mathbb{E}'^\bullet_S$ can be simultaneously truncated to two-term complexes of coherent locally-free modules $\mathbb{F}_S$, $\mathbb{E}_S$. The quasi-isomorphism q' also yields by truncation a quasi-isomorphism $\mathsf{q} : \mathbb{F}_S \to \mathbb{E}_S$.

All above considerations are valid without modification in the presence of a torus action.

□

Now we can conclude the proof of theorem (1.11). The tangent-obstruction theory of the moduli space of ADHM sheaves has been constructed in (5.1), (5.2), the relevant deformation theory results being proven in proposition (4.5), corollary (4.7) and proposition (4.9). The tangent-obstruction theory of the moduli space of admissible pairs has been presented in the proof of theorem (1.9) (see section (6).) Then statement (I) in section (7.4) follows from lemmas (7.8), (7.9), (7.10), (7.11). Statements $(II.a) - (II.c)$ follow from lemmas (7.10) and (7.12).

□

Appendix A. Base Change Properties of Direct Images

For completeness we summarize some standard results on functorial and base change properties of direct images which are used repeatedly in the main text.

Lemma A.1. (*i*) *Let*

$$\begin{tikzcd} Z' \arrow[r, "v'"] \arrow[d, "p'"'] & Z \arrow[d, "p"] \\ S' \arrow[r, "u'"] & S \end{tikzcd}$$

be a commutative diagram of morphisms of Noetherian schemes over $\mathbb{C}$, where $p : Z \to S$ is separated and of finite type. Let $F^\bullet$ be a complex of quasi-coherent $\mathcal{O}_X$ modules. Then we have a natural morphism of quasi-coherent $\mathcal{O}_{S'}$-modules

$$c(u', F^\bullet) : u'^* \mathbf{R}^k p_* F^\bullet \to \mathbf{R}^k p'_*(v'^* F^\bullet) \tag{A.1}$$

for any $k \in \mathbb{Z}$.

(ii) The morphism (A.1) *is functorial in* $F^\bullet$ *i.e. if* $\varphi : F^\bullet \to G^\bullet$ *is a morphism of complexes of quasi-coherent* $\mathcal{O}_Z$*-modules, there is a natural morphism of* $\mathcal{O}_S$*-modules*

$$\mathbf{R}^k p_*(\varphi) : \mathbf{R}^k p_* F^\bullet \to \mathbf{R}^k p_* G^\bullet \tag{A.2}$$

so that the diagram

$$\begin{tikzcd}
u'^*\mathbf{R}^k p_* F^\bullet \arrow[r, "u'^*\mathbf{R}^k p_*(\varphi)"] \arrow[d, "c(u'{,}F^\bullet)"'] & u'^*\mathbf{R}^k p_* G^\bullet \arrow[d, "c(u'{,}G^\bullet)"] \\
\mathbf{R}^k p'_* v'^* F^\bullet \arrow[r, "\mathbf{R}^k p'_*(v'^*\varphi)"] & \mathbf{R}^k p'_* v'^* G^\bullet
\end{tikzcd} \tag{A.3}$$

is commutative for any $k \in \mathbb{Z}$. *(iii) Let*

$$\begin{tikzcd}
Z'' \arrow[r, "v''"] \arrow[d, "p''"] & Z' \arrow[r, "v'"] \arrow[d, "p'"'] & Z \arrow[d, "p"] \\
S'' \arrow[r, "u''"] & S' \arrow[r, "u'"] & S
\end{tikzcd}$$

be a commutative diagram of Noetherian schemes over $\mathbb{C}$, *where* $p : Z \to S$ *is separated and of finite type. Let* $F^\bullet$ *be a complex of quasi-coherent* $\mathcal{O}_Z$*-modules. Then we have a commutative diagram of quasi-coherent* $\mathcal{O}_{S''}$*-modules*

$$\begin{tikzcd}
(u' \circ u'')^*\mathbf{R}^k p_* F^\bullet \arrow[rr, "c(u'\circ u''{,}F^\bullet)"] \arrow[d, "\simeq"'] & & \mathbf{R}^k p''_*(v' \circ v'')^* F^\bullet \arrow[d, "\simeq"] \\
u''^* u'^* \mathbf{R}^k p_* F^\bullet \arrow[dr, "u''^* c(u'{,}F^\bullet)"'] & & \mathbf{R}^k p''_*(v''^* v'^* F^\bullet) \\
& u''^*\mathbf{R}^k p'_*(v'^* F^\bullet) \arrow[ur, "c(v''{,}v'^*F^\bullet)"'] &
\end{tikzcd} \tag{A.4}$$

for any $k \in \mathbb{Z}$.

Proof. The proof follows from the construction of the morphism (A.1) presented in [34, 7.2.2] or alternatively [37, Prop. III.9.3] (see [37, Rem. III.9.3.1].) In particular using Čech cohomology as in the proof of [37, Prop. III.9.3] it reduces to standard properties of modules over Noetherian rings. We will leave the details to the reader.

□

References

[1] *Stack Theory and Applications*. notes taken by H. Clemens at an informal seminar run by A. Bertram, H. Clemens and A. Vistoli; available from www.math.utah.edu/~bertram/lectures.

[2] A. B. Altman and S. L. Kleiman. Compactifying the Picard scheme. *Adv. in Math.*, 35(1):50–112, 1980.

[3] L. Álvarez-Cónsul and O. García-Prada. Dimensional reduction and quiver bundles. *J. Reine Angew. Math.*, 556:1–46, 2003.

[4] L. Álvarez-Cónsul and O. García-Prada. Hitchin-Kobayashi correspondence, quivers, and vortices. *Comm. Math. Phys.*, 238(1-2):1–33, 2003.

[5] L. Álvarez-Cónsul, O. García-Prada, and A. H. W. Schmitt. On the geometry of moduli spaces of holomorphic chains over compact Riemann surfaces. *IMRP Int. Math. Res. Pap.*, pages Art. ID 73597, 82, 2006.

[6] M. Artin. Algebraization of formal moduli i. In *Global Analysis*, pages 21–71. Tokyo, 1969.

[7] M. Artin. Versal deformations and algebraic stacks. *Invent. Math.*, 27:165–189, 1974.

[8] C. Bănică and O. Stănăşilă. *Algebraic methods in the global theory of complex spaces*. Editura Academiei, Bucharest, 1976. Translated from the Romanian.

[9] K. Behrend, B. Conrad, D. Edidin, W. Fulton, B. Fantechi, L. Göttsche, and A. Kresch. *Algebraic Stacks*. available from www.math.lsa.umich.edu/˜wfulton.

[10] K. Behrend and B. Fantechi. The intrinsic normal cone. *Invent. Math.*, 128(1):45–88, 1997.

[11] D. Ben-Zvi and T. Nevins. Perverse bundles and Calogero-Moser spaces. arXiv.org:math/0610097.

[12] J. Bernstein and V. Lunts. *Equivariant sheaves and functors*, volume 1578 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1994.

[13] A. Bertram. Stable pairs and log flips. In *Algebraic geometry—Santa Cruz 1995*, volume 62 of *Proc. Sympos. Pure Math.*, pages 185–201. Amer. Math. Soc., Providence, RI, 1997.

[14] A. Bertram, G. Daskalopoulos, and R. Wentworth. Gromov invariants for holomorphic maps from Riemann surfaces to Grassmannians. *J. Amer. Math. Soc.*, 9(2):529–571, 1996.

[15] S. B. Bradlow and G. D. Daskalopoulos. Moduli of stable pairs for holomorphic bundles over Riemann surfaces. *Internat. J. Math.*, 2(5):477–513, 1991.

[16] S. B. Bradlow, O. García-Prada, and P. B. Gothen. Surface group representations and $\mathrm{U}(p,q)$-Higgs bundles. *J. Differential Geom.*, 64(1):111–170, 2003.

[17] S. B. Bradlow, O. García-Prada, and P. B. Gothen. Moduli spaces of holomorphic triples over compact Riemann surfaces. *Math. Ann.*, 328(1-2):299–351, 2004.

[18] J. Bryan and R. Pandharipande. The local Gromov-Witten theory of curves. math.AG/0411037.

[19] I. Ciocan-Fontanine and M. Kapranov. Virtual fundamental classes via dg-manifolds. arXiv.org:math/0703214.

[20] B. Conrad. *Grothendieck duality and base change*, volume 1750 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2000.

[21] M. Demazure. Structures algébriques, cohomologie des groupes. In *SGA 3: Schémas en groupes*, volume 151 of *Lecture Notes in Mathematics*, pages 1–37. Springer Verlag, Berlin, 1970.

[22] D. E. Diaconescu. Moduli of ADHM sheaves in string theory. unpublished manuscript.

[23] D. Edidin and W. Graham. Algebraic cycles and completions of equivariant K-theory. math.AG/0702671.

[24] D. Eisenbud. *Commutative algebra*, volume 150 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995. With a view toward algebraic geometry.

[25] B. Fantechi and Göttsche. Riemann-Roch theorems and elliptic genus for virtually smooth schemes.

[26] B. Fantechi, L. Göttsche, L. Illusie, S. L. Kleiman, N. Nitsure, and A. Vistoli. *Fundamental algebraic geometry*, volume 123 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2005. Grothendieck's FGA explained.

[27] W. Fulton. *Intersection theory*, volume 2 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, second edition, 1998.

[28] T. L. Gómez. Algebraic stacks. *Proc. Indian Acad. Sci. Math. Sci.*, 111(1):1–31, 2001.

[29] T. L. Gomez and I. Sols. Stable tensors and moduli space of orthogonal sheaves. math.AG/0103150.

[30] P. B. Gothen and A. D. King. Homological algebra of twisted quiver bundles. *J. London Math. Soc. (2)*, 71(1):85–99, 2005.

[31] T. Graber and R. Pandharipande. Localization of virtual classes. *Invent. Math.*, 135(2):487–518, 1999.

[32] A. Grothendieck. Éléments de géométrie algébrique. I. Le langage des schémas. *Inst. Hautes Études Sci. Publ. Math.*, (4):228, 1960.

[33] A. Grothendieck. Éléments de géométrie algébrique. III. Étude cohomologique des faisceaux cohérents. I. *Inst. Hautes Études Sci. Publ. Math.*, (11):167, 1961.

[34] A. Grothendieck. Éléments de géométrie algébrique. III. Étude cohomologique des faisceaux cohérents. II. *Inst. Hautes Études Sci. Publ. Math.*, (17):91, 1963.

[35] A. Grothendieck. Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. II. *Inst. Hautes Études Sci. Publ. Math.*, (24):231, 1965.

[36] R. Hartshorne. *Residues and duality*. Lecture notes of a seminar on the work of A. Grothendieck, given at Harvard 1963/64. With an appendix by P. Deligne. Lecture Notes in Mathematics, No. 20. Springer-Verlag, Berlin, 1966.

[37] R. Hartshorne. *Algebraic geometry*. Springer-Verlag, New York, 1977. Graduate Texts in Mathematics, No. 52.

[38] D. Huybrechts and M. Lehn. Framed modules and their moduli. *Internat. J. Math.*, 6(2):297–324, 1995.

[39] D. Huybrechts and M. Lehn. Stable pairs on curves and surfaces. *J. Algebraic Geom.*, 4(1):67–104, 1995.

[40] D. Huybrechts and M. Lehn. *The geometry of moduli spaces of sheaves*. Aspects of Mathematics, E31. Friedr. Vieweg & Sohn, Braunschweig, 1997.

[41] M. Jardim. Moduli spaces of framed instanton sheaves on projective spaces. arXiv.org:math.AG/0801.2550.

[42] S. Keel and S. Mori. Quotients by groupoids. *Ann. of Math. (2)*, 145(1):193–213, 1997.

[43] B. Kim, A. Kresch, and T. Pantev. Functoriality in intersection theory and a conjecture of Cox, Katz, and Lee. *J. Pure Appl. Algebra*, 179(1-2):127–136, 2003.

[44] G. Laumon and L. Moret-Bailly. *Champs algébriques*, volume 39 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 2000.

[45] J. Le Poitier. Systèmes coh 'erents et structures de niveau. *Astérisque*, 214:143, 1993.

[46] J. Le Poitier. Faisceaux semi-stable et systèmes coh 'erents. In *Vector Bundles in Algebraic Geometry (Durham, 1993)*, volume 208 of *London Math. Soc. Lecture Notes Ser.*, pages 179–239. Cambridge University Press, Cambridge, 1995.

[47] J. Li. A degeneration formula of GW-invariants. *J. Differential Geom.*, 60(2):199–293, 2002.

[48] J. Li and G. Tian. Virtual moduli cycles and Gromov-Witten invariants of algebraic varieties. *J. Amer. Math. Soc.*, 11(1):119–174, 1998.

[49] M. Maruyama. On boundedness of families of torsion free sheaves. *J. Math. Kyoto Univ.*, 21:673–701, 1981.

[50] D. Maulik, N. Nekrasov, A. Okounkov, and R. Pandharipande. Gromov-Witten theory and Donaldson-Thomas theory. I. *Compos. Math.*, 142(5):1263–1285, 2006.

[51] D. Maulik, N. Nekrasov, A. Okounkov, and R. Pandharipande. Gromov-Witten theory and Donaldson-Thomas theory. II. *Compos. Math.*, 142(5):1286–1304, 2006.

[52] T. Mochizuki. A theory of the invariants obtained from the moduli stacks of stable objects on a smooth polarized surface. arXiv:math/0210211.

[53] H. Nakajima. *Lectures on Hilbert schemes of points on surfaces*, volume 18 of *University Lecture Series*. American Mathematical Society, Providence, RI, 1999.

[54] H. Nakajima and K. Yoshioka. Instanton counting on blowup. I. 4-dimensional pure gauge theory. *Invent. Math.*, 162(2):313–355, 2005.

[55] H. Nakajima and K. Yoshioka. Instanton counting on blowup. II. K-theoretic partition function. *Transform. Groups*, 10(3-4):489–519, 2005.

[56] N. Nitsure. Moduli space of semistable pairs on a curve. *Proc. London Math. Soc. (3)*, 62(2):275–300, 1991.

[57] C. Okonek and A. Teleman. Comparing virtual fundamental classes: gauge theoretical Gromov-Witten invariants for toric varieties. *Asian J. Math.*, 7(2):167–198, 2003.

[58] A. Okounkov and R. Pandharipande. The local Donaldson-Thomas theory of curves. math.AG/0512573.

[59] D. O. Orlov. Projective bundles, monoidal transformations, and derived categories of coherent sheaves. *Izv. Ross. Akad. Nauk Ser. Mat.*, 56(4):852–862, 1992.

[60] R. Pandharipande and R. P. Thomas. The 3-fold vertex via stable pairs. arXiv.org:0709.3823.

[61] R. Pandharipande and R. P. Thomas. Curve counting via stable pairs in the derived category. arXiv.org:0707.2348.

[62] R. Pandharipande and R. P. Thomas. Stable pairs and BPS invariants. arXiv.org:0711.3899.

[63] M. Romagny. Group actions on stacks and applications. *Michigan Math. J.*, 53(1):209–236, 2005.

[64] A. Schmitt. Projective moduli for Hitchin pairs. *Internat. J. Math.*, 9(1):107–118, 1998.

[65] A. Schmitt. A universal construction for moduli spaces of decorated vector bundles over curves. *Transform. Groups*, 9(2):167–209, 2004.

[66] A. Schmitt. Moduli for decorated tuples of sheaves and representation spaces for quivers. *Proc. Indian Acad. Sci. Math. Sci.*, 115(1):15–49, 2005.

[67] B. Szendroi. Sheaves on fibered threefolds and quiver sheaves. arXiv.org:math/0506301.

[68] M. Thaddeus. Stable pairs, linear systems and the Verlinde formula. *Invent. Math.*, 117(2):317–353, 1994.

[69] R. P. Thomas. A holomorphic Casson invariant for Calabi-Yau 3-folds, and bundles on $K3$ fibrations. *J. Differential Geom.*, 54(2):367–438, 2000.

[70] R. W. Thomason. Equivariant resolution, linearization, and Hilbert's fourteenth problem over arbitrary base schemes. *Adv. in Math.*, 65(1):16–34, 1987.

NHETC, Rutgers University, 126 Frelinghuysen Rd, Piscataway NJ 08854